

CS 107, Probability, Spring 2019

Lecture 33

Michael Poghosyan

AUA

10 April 2019

- Joint Distribution of two R.V.s, Continuous Case

Example:

Example: Assume (X, Y) is a continuous r.vector with a Joint PDF

$$f(x, y) = \begin{cases} K \cdot (x + y), & x, y \in [0, 2] \\ 0, & \text{otherwise} \end{cases}$$

- Find K
- Find the Marginal PDFs of X and Y
- Find the CDFs of (X, Y) , and Marginal CDFs
- Calculate $\mathbb{P}(X^2 + Y^2 \leq 1)$
- Calculate the PDF of the r.v. $Z = Y - X$

Distribution of more than 2 r.v.s

The notions concerning the Distribution of 2 r.v.s can be generalized to more than 2 r.v.s.

Distribution of more than 2 r.v.s

The notions concerning the Distribution of 2 r.v.s can be generalized to more than 2 r.v.s. Say, if X , Y , Z are r.v. in the same Experiment, then

Distribution of more than 2 r.v.s

The notions concerning the Distribution of 2 r.v.s can be generalized to more than 2 r.v.s. Say, if X, Y, Z are r.v. in the same Experiment, then

- Their Joint CDF is:

$$F(x, y, z) = \mathbb{P}(X \leq x, Y \leq y, Z \leq z), \quad \forall (x, y, z) \in \mathbb{R}^3;$$

Distribution of more than 2 r.v.s

The notions concerning the Distribution of 2 r.v.s can be generalized to more than 2 r.v.s. Say, if X, Y, Z are r.v. in the same Experiment, then

- Their Joint CDF is:

$$F(x, y, z) = \mathbb{P}(X \leq x, Y \leq y, Z \leq z), \quad \forall (x, y, z) \in \mathbb{R}^3;$$

- If X, Y, Z are Discrete, then their Joint PMF is given by

$$p_{i,j,k} = \mathbb{P}(X = x_i, Y = y_j, Z = z_k)$$

Distribution of more than 2 r.v.s

The notions concerning the Distribution of 2 r.v.s can be generalized to more than 2 r.v.s. Say, if X, Y, Z are r.v. in the same Experiment, then

- Their Joint CDF is:

$$F(x, y, z) = \mathbb{P}(X \leq x, Y \leq y, Z \leq z), \quad \forall (x, y, z) \in \mathbb{R}^3;$$

- If X, Y, Z are Discrete, then their Joint PMF is given by

$$p_{i,j,k} = \mathbb{P}(X = x_i, Y = y_j, Z = z_k)$$

- And, in the latter case, say, the Distribution of X can be found by:

$$\mathbb{P}(X = x_i) =$$

Distribution of more than 2 r.v.s

The notions concerning the Distribution of 2 r.v.s can be generalized to more than 2 r.v.s. Say, if X, Y, Z are r.v. in the same Experiment, then

- Their Joint CDF is:

$$F(x, y, z) = \mathbb{P}(X \leq x, Y \leq y, Z \leq z), \quad \forall (x, y, z) \in \mathbb{R}^3;$$

- If X, Y, Z are Discrete, then their Joint PMF is given by

$$p_{i,j,k} = \mathbb{P}(X = x_i, Y = y_j, Z = z_k)$$

- And, in the latter case, say, the Distribution of X can be found by:

$$\mathbb{P}(X = x_i) = \sum_{j,k} \mathbb{P}(X = x_i, Y = y_j, Z = z_k)$$

Distribution of more than 2 r.v.s

- And if X, Y, Z are Jointly Continuous, then their Joint PDF $f(x, y, z)$ and Joint CDF $F(x, y, z)$ satisfy

$$F(x, y, z) = \int_{-\infty}^x \int_{-\infty}^y \int_{-\infty}^z f(u, v, w) du dv dw$$

Distribution of more than 2 r.v.s

- And if X, Y, Z are Jointly Continuous, then their Joint PDF $f(x, y, z)$ and Joint CDF $F(x, y, z)$ satisfy

$$F(x, y, z) = \int_{-\infty}^x \int_{-\infty}^y \int_{-\infty}^z f(u, v, w) du dv dw$$

- And, in the latter case, say, the PDF of X can be found by:

$$f_X(x) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y, z) dy dz$$

This is the "integrating y and z out" technique.

Distribution of more than 2 r.v.s

- And if X, Y, Z are Jointly Continuous, then their Joint PDF $f(x, y, z)$ and Joint CDF $F(x, y, z)$ satisfy

$$F(x, y, z) = \int_{-\infty}^x \int_{-\infty}^y \int_{-\infty}^z f(u, v, w) du dv dw$$

- And, in the latter case, say, the PDF of X can be found by:

$$f_X(x) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y, z) dy dz$$

This is the "integrating y and z out" technique.

Exercise: Can you express the Joint PDF of, say, X and Z , in terms of $f(x, y, z)$?

Important Multivariate Distributions

Discrete Distribution: Multinomial

One of the famous Discrete Multivariate Distributions is the Multinomial Distribution:

Discrete Distribution: Multinomial

One of the famous Discrete Multivariate Distributions is the Multinomial Distribution:

Multinomial Distribution

Assume $n, m \in \mathbb{N}$, and $p_k \in [0, 1]$, $k = 1, 2, \dots, m$ with $p_1 + \dots + p_m = 1$. We say that the r.vector $\mathbf{X} = (X_1, X_2, \dots, X_m)$ has a Multinomial Distribution with probabilities $\mathbf{p} = (p_1, p_2, \dots, p_m)$, and we write

$$\mathbf{X} = (X_1, X_2, \dots, X_m) \sim \text{Multinomial}(n, p_1, p_2, \dots, p_m),$$

if its PMF is given by:

$$\mathbb{P}(X_1 = k_1, X_2 = k_2, \dots, X_m = k_m) = \binom{n}{k_1, k_2, \dots, k_m} \cdot p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_m^{k_m}$$

for any $k_1, \dots, k_m \in \mathbb{N} \cup \{0\}$, with $k_1 + k_2 + \dots + k_m = n$.

Multinomial Distribution

Multinomial Distribution is used when modeling the following type of Experiments:

Multinomial Distribution

Multinomial Distribution is used when modeling the following type of Experiments:

- Our Experiment consists of n times independent repetition of the same Simple Experiment;

Multinomial Distribution

Multinomial Distribution is used when modeling the following type of Experiments:

- Our Experiment consists of n times independent repetition of the same Simple Experiment;
- In our Simple Experiment, there are m possible outcomes, and the probability to have the outcome k is p_k for $k = 1, 2, \dots, m$;

Multinomial Distribution

Multinomial Distribution is used when modeling the following type of Experiments:

- Our Experiment consists of n times independent repetition of the same Simple Experiment;
- In our Simple Experiment, there are m possible outcomes, and the probability to have the outcome k is p_k for $k = 1, 2, \dots, m$;
- The r.v. X_k shows the number of outcome k obtained in n repetitions.

Multinomial Distribution

Multinomial Distribution is used when modeling the following type of Experiments:

- Our Experiment consists of n times independent repetition of the same Simple Experiment;
- In our Simple Experiment, there are m possible outcomes, and the probability to have the outcome k is p_k for $k = 1, 2, \dots, m$;
- The r.v. X_k shows the number of outcome k obtained in n repetitions.

Exercise: Prove that the Marginal Distributions of r.v. X_k are Bernoulli, particularly, $X_k \sim \text{Bernoulli}(n, p_k)$

Multinomial Distribution

Multinomial Distribution is used when modeling the following type of Experiments:

- Our Experiment consists of n times independent repetition of the same Simple Experiment;
- In our Simple Experiment, there are m possible outcomes, and the probability to have the outcome k is p_k for $k = 1, 2, \dots, m$;
- The r.v. X_k shows the number of outcome k obtained in n repetitions.

Exercise: Prove that the Marginal Distributions of r.v. X_k are Bernoulli, particularly, $X_k \sim \text{Bernoulli}(n, p_k)$

Exercise: Find the Joint Distribution of, say, (X_1, X_2) .

Example

Example: 10 AUA instructors are choosing (independently) at random one AUA student for some committee. We know that the relationship between the number of Bus/CSE/EC students is $8 : 3 : 1$. What is the Probability that among 10 chosen students, we will have exactly 3 Bus, 5 CSE and 2 EC students?