

CS 107, Probability, Spring 2019

Lecture 11

Michael Poghosyan

AUA

11 February 2019

Content

- Total Probability Formula
- Bayes Formula

Medical Test gives a correct answer in 95% of cases: if the person is ill, it is saying that he/she is ill with 95% probability, and if the person is healthy, it is saying that he/she is healthy with 95% probability (i.e., in 95% of cases).

Medical Test gives a correct answer in 95% of cases: if the person is ill, it is saying that he/she is ill with 95% probability, and if the person is healthy, it is saying that he/she is healthy with 95% probability (i.e., in 95% of cases).

Now assume I am taking that test, and it says that I am ill. What is the probability that I am really ill?

The Multiplication Rule

From the Conditional Probability definition, we obtain the following Multiplication (or Chain) Rule:

$$\mathbb{P}(A \cap B) = \mathbb{P}(A|B) \cdot \mathbb{P}(B).$$

Example, from the last lecture

Problem: We have 36 participants in our Prob class, from which 26 are male.

Example, from the last lecture

Problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz.

Example, from the last lecture

Problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

Example, from the last lecture

Problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

We know that

$$\mathbb{P}(\text{chosen person is a female}) = \frac{10}{36} = \frac{5}{18}$$

Also,

$$\mathbb{P}(\text{ person loves Jazz} | \text{ person is a female}) =$$

Example, from the last lecture

Problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

We know that

$$\mathbb{P}(\text{chosen person is a female}) = \frac{10}{36} = \frac{5}{18}$$

Also,

$$\mathbb{P}(\text{ person loves Jazz} | \text{ person is a female}) = 0.15.$$

So

$$\mathbb{P}(\text{chosen person is a female and she loves Jazz}) = 0.15 \cdot \frac{5}{18}.$$

Example, from the last lecture

Problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

We know that

$$\mathbb{P}(\text{chosen person is a female}) = \frac{10}{36} = \frac{5}{18}$$

Also,

$$\mathbb{P}(\text{ person loves Jazz} | \text{ person is a female}) = 0.15.$$

So

$$\mathbb{P}(\text{chosen person is a female and she loves Jazz}) = 0.15 \cdot \frac{5}{18}.$$

Now, let us give this in the tree form: look at the Board!

Jazz Problem, Cont'd

Now, let us complete the tree!

Jazz Problem, Cont'd

Now, let us complete the tree!
Recall the above problem:

Jazz Problem, Cont'd

Now, let us complete the tree!

Recall the above problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

Jazz Problem, Cont'd

Now, let us complete the tree!

Recall the above problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

Assume, that in the problem above we know additionally that only 20% of males love Jazz.

Jazz Problem, Cont'd

Now, let us complete the tree!

Recall the above problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

Assume, that in the problem above we know additionally that only 20% of males love Jazz.

Then we can calculate the probability that the chosen person will be a Jazz lover male student.

Jazz Problem, Cont'd

Now, let us complete the tree!

Recall the above problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

Assume, that in the problem above we know additionally that only 20% of males love Jazz.

Then we can calculate the probability that the chosen person will be a Jazz lover male student.

$$\mathbb{P}(\text{chosen person is a male and he loves Jazz}) =$$

Jazz Problem, Cont'd

Now, let us complete the tree!

Recall the above problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

Assume, that in the problem above we know additionally that only 20% of males love Jazz.

Then we can calculate the probability that the chosen person will be a Jazz lover male student.

$$\mathbb{P}(\text{chosen person is a male and he loves Jazz}) =$$

Jazz Problem, Cont'd

Now, let us complete the tree!

Recall the above problem: We have 36 participants in our Prob class, from which 26 are male. Assume we know that only 15% of females love Jazz. We choose a participant from our class at random. What is the Probability that the chosen person will be a Jazz lover female student?

Assume, that in the problem above we know additionally that only 20% of males love Jazz.

Then we can calculate the probability that the chosen person will be a Jazz lover male student.

$$\mathbb{P}(\text{chosen person is a male and he loves Jazz}) =$$

Important is that now we can calculate the probability that the chosen person will like Jazz:

Jazz Problem, Cont'd

$$\begin{aligned}\mathbb{P}(\text{chosen person likes Jazz}) &= \\ &= \mathbb{P}(\text{chosen person is a male and he loves Jazz}) + \\ &+ \mathbb{P}(\text{chosen person is a female and she loves Jazz}) =\end{aligned}$$

Jazz Problem, Cont'd

$$\begin{aligned}\mathbb{P}(\text{chosen person likes Jazz}) &= \\ &= \mathbb{P}(\text{chosen person is a male and he loves Jazz}) + \\ &+ \mathbb{P}(\text{chosen person is a female and she loves Jazz}) =\end{aligned}$$

Jazz Problem, Cont'd

$$\begin{aligned}\mathbb{P}(\text{chosen person likes Jazz}) &= \\ &= \mathbb{P}(\text{chosen person is a male and he loves Jazz}) + \\ &+ \mathbb{P}(\text{chosen person is a female and she loves Jazz}) =\end{aligned}$$

because the two events above (being a male Jazz lover and female jazz lover) are disjoint.

Jazz Problem, Cont'd

$$\begin{aligned}\mathbb{P}(\text{chosen person likes Jazz}) &= \\ &= \mathbb{P}(\text{chosen person is a male and he loves Jazz}) + \\ &+ \mathbb{P}(\text{chosen person is a female and she loves Jazz}) =\end{aligned}$$

because the two events above (being a male Jazz lover and female jazz lover) are disjoint.

Now, let us generalize the formula:

The Total Probability Formula

Assume A and B are events in some Experiment. Then

The Total Probability Formula

Assume A and B are events in some Experiment. Then

Total Probability Formula

$$\mathbb{P}(A) = \mathbb{P}(B) \cdot \mathbb{P}(A|B) + \mathbb{P}(\overline{B}) \cdot \mathbb{P}(A|\overline{B})$$

The Total Probability Formula

Assume A and B are events in some Experiment. Then

Total Probability Formula

$$\mathbb{P}(A) = \mathbb{P}(B) \cdot \mathbb{P}(A|B) + \mathbb{P}(\bar{B}) \cdot \mathbb{P}(A|\bar{B})$$

Here B and \bar{B} are called Hypotheses, and satisfy

$$B \cap \bar{B} = \emptyset \quad \text{and} \quad B \cup \bar{B} = \Omega,$$

The Total Probability Formula

Assume A and B are events in some Experiment. Then

Total Probability Formula

$$\mathbb{P}(A) = \mathbb{P}(B) \cdot \mathbb{P}(A|B) + \mathbb{P}(\overline{B}) \cdot \mathbb{P}(A|\overline{B})$$

Here B and \overline{B} are called Hypotheses, and satisfy

$$B \cap \overline{B} = \emptyset \quad \text{and} \quad B \cup \overline{B} = \Omega,$$

that is, **we take a partition of our Sample Space into disjoint Hypotheses (Events), calculate the probability of our event A under each Hypothesis, and then add the results.**

The Total Probability Formula

The general form of the TPF is: assume we want to calculate $\mathbb{P}(A)$.

The Total Probability Formula

The general form of the TPF is: assume we want to calculate $\mathbb{P}(A)$. We consider Hypotheses B_1, B_2, \dots, B_n such that

$$B_i \cap B_j = \emptyset \quad \text{and} \quad \bigcup_{k=1}^n B_k = \Omega$$

The Total Probability Formula

The general form of the TPF is: assume we want to calculate $\mathbb{P}(A)$. We consider Hypotheses B_1, B_2, \dots, B_n such that

$$B_i \cap B_j = \emptyset \quad \text{and} \quad \bigcup_{k=1}^n B_k = \Omega$$

Then

TPF

$$\mathbb{P}(A) = \mathbb{P}(B_1) \cdot \mathbb{P}(A|B_1) + \mathbb{P}(B_2) \cdot \mathbb{P}(A|B_2) + \dots + \mathbb{P}(B_n) \cdot \mathbb{P}(A|B_n).$$

Example

Problem: We have an urn full of 100 balls, from which 15 are white, 25 are red, and the rest are black.

Example

Problem: We have an urn full of 100 balls, from which 15 are white, 25 are red, and the rest are black. Our balls are of two type: round and cubic ☺. The proportion of round balls in whites is 20%, in reds - 40%, and in blacks - 55%.

Example

Problem: We have an urn full of 100 balls, from which 15 are white, 25 are red, and the rest are black. Our balls are of two type: round and cubic 😊. The proportion of round balls in whites is 20%, in reds - 40%, and in blacks - 55%. We take a ball at random. What is the probability of drawing a cubic ball?

Example

Problem: We have an urn full of 100 balls, from which 15 are white, 25 are red, and the rest are black. Our balls are of two type: round and cubic 😊. The proportion of round balls in whites is 20%, in reds - 40%, and in blacks - 55%. We take a ball at random. What is the probability of drawing a cubic ball?

Well, Probabilists love urns and balls, and dice and coins. The problem may seem very artificial. So let me give a problem of this type in a more realistic terms:

Example

Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30.

Example

Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30. The proportion of smokers in the young-age population is 20%, in mid-age-population - 40%, and in older-age-population - 55%.

Example

Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30. The proportion of smokers in the young-age population is 20%, in mid-age-population - 40%, and in older-age-population - 55%. What is the probability of that a person in that city will be a cubic ball?

Example

Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30. The proportion of smokers in the young-age population is 20%, in mid-age-population - 40%, and in older-age-population - 55%. What is the probability of that a person in that city will be a cubic ball? I mean, the person will be a smoker? 😊

Example

Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30. The proportion of smokers in the young-age population is 20%, in mid-age-population - 40%, and in older-age-population - 55%. What is the probability of that a person in that city will be a cubic ball? I mean, the person will be a smoker? 😊

Solution: OTB

Bayes Formula

Now, let us talk about one of the most important formulas, the Bayes Formula.

Bayes Formula

Now, let us talk about one of the most important formulas, the Bayes Formula. It is used a lot in Statistics, Machine Learning, Signal Processing, Finance, Bioinformatics etc.

Bayes Formula

Now, let us talk about one of the most important formulas, the Bayes Formula. It is used a lot in Statistics, Machine Learning, Signal Processing, Finance, Bioinformatics etc.

The formula is simple: we just write the property of the Conditional Probabilities:

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B|A) = \mathbb{P}(B) \cdot \mathbb{P}(A|B).$$

Bayes Formula

Now, let us talk about one of the most important formulas, the Bayes Formula. It is used a lot in Statistics, Machine Learning, Signal Processing, Finance, Bioinformatics etc.

The formula is simple: we just write the property of the Conditional Probabilities:

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B|A) = \mathbb{P}(B) \cdot \mathbb{P}(A|B).$$

Hence,

Bayes Formula

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A|B) \cdot \mathbb{P}(B)}{\mathbb{P}(A)}$$

Bayes Formula, Cont'd

In the Bayes formula,

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A|B) \cdot \mathbb{P}(B)}{\mathbb{P}(A)},$$

Bayes Formula, Cont'd

In the Bayes formula,

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A|B) \cdot \mathbb{P}(B)}{\mathbb{P}(A)},$$

- $\mathbb{P}(B)$ is the **Prior Probability** of B

Bayes Formula, Cont'd

In the Bayes formula,

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A|B) \cdot \mathbb{P}(B)}{\mathbb{P}(A)},$$

- $\mathbb{P}(B)$ is the **Prior Probability** of B
- $\mathbb{P}(B|A)$ is the **Posterior Probability** of B , after observing A

Bayes Formula, Cont'd

In the Bayes formula,

$$\mathbb{P}(B|A) = \frac{\mathbb{P}(A|B) \cdot \mathbb{P}(B)}{\mathbb{P}(A)},$$

- $\mathbb{P}(B)$ is the **Prior Probability** of B
- $\mathbb{P}(B|A)$ is the **Posterior Probability** of B , after observing A
- $\mathbb{P}(A|B)$ is called the **Likelihood** of A under B .

Bayes Formula, General Form

Now, the general form of the Bayes Formula, and the interpretation of that Formula as an Inverse Probability Calculation Formula.

Bayes Formula, General Form

Now, the general form of the Bayes Formula, and the interpretation of that Formula as an Inverse Probability Calculation Formula.

Assume we have an event A . We have also some Hypotheses B_1, B_2, \dots, B_n .

Bayes Formula, General Form

Now, the general form of the Bayes Formula, and the interpretation of that Formula as an Inverse Probability Calculation Formula.

Assume we have an event A . We have also some Hypotheses B_1, B_2, \dots, B_n .

- The Direct Problem:

Bayes Formula, General Form

Now, the general form of the Bayes Formula, and the interpretation of that Formula as an Inverse Probability Calculation Formula.

Assume we have an event A . We have also some Hypotheses B_1, B_2, \dots, B_n .

- The Direct Problem: Calculate the Probability of A , given the Hypotheses.

Bayes Formula, General Form

Now, the general form of the Bayes Formula, and the interpretation of that Formula as an Inverse Probability Calculation Formula.

Assume we have an event A . We have also some Hypotheses B_1, B_2, \dots, B_n .

- The Direct Problem: Calculate the Probability of A , given the Hypotheses. Solution: TPF

$$\mathbb{P}(A) = \mathbb{P}(B_1) \cdot \mathbb{P}(A|B_1) + \mathbb{P}(B_2) \cdot \mathbb{P}(A|B_2) + \dots + \mathbb{P}(B_n) \cdot \mathbb{P}(A|B_n)$$

Bayes Formula, General Form

Now, the general form of the Bayes Formula, and the interpretation of that Formula as an Inverse Probability Calculation Formula.

Assume we have an event A . We have also some Hypotheses B_1, B_2, \dots, B_n .

- The Direct Problem: Calculate the Probability of A , given the Hypotheses. Solution: TPF

$$\mathbb{P}(A) = \mathbb{P}(B_1) \cdot \mathbb{P}(A|B_1) + \mathbb{P}(B_2) \cdot \mathbb{P}(A|B_2) + \dots + \mathbb{P}(B_n) \cdot \mathbb{P}(A|B_n)$$

- The Inverse Problem:

Bayes Formula, General Form

Now, the general form of the Bayes Formula, and the interpretation of that Formula as an Inverse Probability Calculation Formula.

Assume we have an event A . We have also some Hypotheses B_1, B_2, \dots, B_n .

- The Direct Problem: Calculate the Probability of A , given the Hypotheses. Solution: TPF

$$\mathbb{P}(A) = \mathbb{P}(B_1) \cdot \mathbb{P}(A|B_1) + \mathbb{P}(B_2) \cdot \mathbb{P}(A|B_2) + \dots + \mathbb{P}(B_n) \cdot \mathbb{P}(A|B_n)$$

- The Inverse Problem: Calculate the Probability that B_k happened, given that we have observed A .

Bayes Formula, General Form

Now, the general form of the Bayes Formula, and the interpretation of that Formula as an Inverse Probability Calculation Formula.

Assume we have an event A . We have also some Hypotheses B_1, B_2, \dots, B_n .

- The Direct Problem: Calculate the Probability of A , given the Hypotheses. Solution: TPF

$$\mathbb{P}(A) = \mathbb{P}(B_1) \cdot \mathbb{P}(A|B_1) + \mathbb{P}(B_2) \cdot \mathbb{P}(A|B_2) + \dots + \mathbb{P}(B_n) \cdot \mathbb{P}(A|B_n)$$

- The Inverse Problem: Calculate the Probability that B_k happened, given that we have observed A . Solution:
Bayes Formula:

$$\mathbb{P}(B_k|A) = \frac{\mathbb{P}(A|B_k) \cdot \mathbb{P}(B_k)}{\mathbb{P}(A)} =$$

Bayes Formula, General Form

Now, the general form of the Bayes Formula, and the interpretation of that Formula as an Inverse Probability Calculation Formula.

Assume we have an event A . We have also some Hypotheses B_1, B_2, \dots, B_n .

- The Direct Problem: Calculate the Probability of A , given the Hypotheses. Solution: TPF

$$\mathbb{P}(A) = \mathbb{P}(B_1) \cdot \mathbb{P}(A|B_1) + \mathbb{P}(B_2) \cdot \mathbb{P}(A|B_2) + \dots + \mathbb{P}(B_n) \cdot \mathbb{P}(A|B_n)$$

- The Inverse Problem: Calculate the Probability that B_k happened, given that we have observed A . Solution: Bayes Formula:

$$\mathbb{P}(B_k|A) = \frac{\mathbb{P}(A|B_k) \cdot \mathbb{P}(B_k)}{\mathbb{P}(A)} = \frac{\mathbb{P}(A|B_k) \cdot \mathbb{P}(B_k)}{\sum_{i=1}^n \mathbb{P}(A|B_i) \cdot \mathbb{P}(B_i)}.$$

Example

Let us recall our Non-Smoking City Problem:

Example

Let us recall our Non-Smoking City Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30.

Example

Let us recall our Non-Smoking City Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30. The proportion of smokers in the younger-age population is 20%, in mid-age-population - 40%, and in older-age-population - 55%.

Example

Let us recall our Non-Smoking City Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30. The proportion of smokers in the younger-age population is 20%, in mid-age-population - 40%, and in older-age-population - 55%. What is the probability of that a person in that city is a smoker?

Example

Let us recall our Non-Smoking City Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30. The proportion of smokers in the younger-age population is 20%, in mid-age-population - 40%, and in older-age-population - 55%. What is the probability of that a person in that city is a smoker?

We know the solution.

Example

Let us recall our Non-Smoking City Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30. The proportion of smokers in the younger-age population is 20%, in mid-age-population - 40%, and in older-age-population - 55%. What is the probability of that a person in that city is a smoker?

We know the solution. Now, the Inverse Probability question: Assume that the randomly chosen person is a smoker. What is the probability that he/she is of younger-age-group?

Example

Let us recall our Non-Smoking City Problem: We know that in some city, 15% of persons are below the age 15, 25% are between 15 and 30, and the rest of population in that city are above 30. The proportion of smokers in the younger-age population is 20%, in mid-age-population - 40%, and in older-age-population - 55%. What is the probability of that a person in that city is a smoker?

We know the solution. Now, the Inverse Probability question: Assume that the randomly chosen person is a smoker. What is the probability that he/she is of younger-age-group?

Solution: OTB

Some Questions to Answer

- Give the TPF.

Some Questions to Answer

- Give the TPF.
- Give the Bayes Formula.

Some Questions to Answer

- Give the TPF.
- Give the Bayes Formula.
- Give the idea of the Bayes Formula.