

# CS 107, Probability, Spring 2019

## Lecture 04

Michael Poghosyan

AUA

23 January 2019

- Properties of the Probability Measure
- Classical Probability Models: Discrete Sample Spaces

Theorem: You can decompose a solid ball into finitely many pieces (5 pieces will suffice) and reassemble them using only rigid motions (rotations + translations) to get two solid balls of the same size.

Theorem: You can decompose a solid ball into finitely many pieces (5 pieces will suffice) and reassemble them using only rigid motions (rotations + translations) to get two solid balls of the same size.

In fact, you can decompose an orange into finitely many pieces, and then reassemble that pieces to obtain the Earth.

Theorem: You can decompose a solid ball into finitely many pieces (5 pieces will suffice) and reassemble them using only rigid motions (rotations + translations) to get two solid balls of the same size.

In fact, you can decompose an orange into finitely many pieces, and then reassemble that pieces to obtain the Earth. Well, the material will be the orange one, of course, but the size will be the Earth one.

Theorem: You can decompose a solid ball into finitely many pieces (5 pieces will suffice) and reassemble them using only rigid motions (rotations + translations) to get two solid balls of the same size.

In fact, you can decompose an orange into finitely many pieces, and then reassemble that pieces to obtain the Earth. Well, the material will be the orange one, of course, but the size will be the Earth one.

This is the famous Banach-Tarski Paradox. See <https://www.youtube.com/watch?v=s86-Z-CbaHA>

# Properties of the Probability Measure (from our last lecture)

The Probability Measure satisfies the following properties:

1.  $\mathbb{P}(\emptyset) = 0$ ;
2. If  $A \cap B = \emptyset$ , then  $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$ ;
3.  $\mathbb{P}(\overline{A}) = 1 - \mathbb{P}(A)$  for any Event  $A$ ;
4. If  $A_i \cap A_j = \emptyset$  for  $i \neq j$ , then  $\mathbb{P}(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n \mathbb{P}(A_i)$ ;
5. for any events  $A, B \in \mathcal{F}$ ,

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B);$$

# Properties of the Probability Measure

6. If  $A_1, A_2, \dots, A_n \in \mathcal{F}$  are events, not necessarily disjoint, then

$$\begin{aligned}\mathbb{P}\left(\bigcup_{i=1}^n A_i\right) &= \mathbb{P}(A_1) + \mathbb{P}(A_2) + \dots + \mathbb{P}(A_n) - \\ &\quad - \mathbb{P}(A_1 \cap A_2) - \mathbb{P}(A_1 \cap A_3) - \dots - \mathbb{P}(A_{n-1} \cap A_n) + \\ &\quad + \mathbb{P}(A_1 \cap A_2 \cap A_3) + \mathbb{P}(A_1 \cap A_2 \cap A_4) + \dots + \mathbb{P}(A_{n-2} \cap A_{n-1} \cap A_n) - \dots \\ &\quad \dots + (-1)^{n-1} \mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_n).\end{aligned}$$



# Properties of the Probability Measure

6. If  $A_1, A_2, \dots, A_n \in \mathcal{F}$  are events, not necessarily disjoint, then

$$\begin{aligned}\mathbb{P}\left(\bigcup_{i=1}^n A_i\right) &= \mathbb{P}(A_1) + \mathbb{P}(A_2) + \dots + \mathbb{P}(A_n) - \\ &\quad - \mathbb{P}(A_1 \cap A_2) - \mathbb{P}(A_1 \cap A_3) - \dots - \mathbb{P}(A_{n-1} \cap A_n) + \\ &\quad + \mathbb{P}(A_1 \cap A_2 \cap A_3) + \mathbb{P}(A_1 \cap A_2 \cap A_4) + \dots + \mathbb{P}(A_{n-2} \cap A_{n-1} \cap A_n) - \dots \\ &\quad \dots + (-1)^{n-1} \mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_n).\end{aligned}$$

This is the general version of the previous property, and is called the inclusion-exclusion principle.

# Properties of the Probability Measure

7. for any events  $A, B \in \mathcal{F}$

$$\mathbb{P}(B \setminus A) = \mathbb{P}(B) - \mathbb{P}(A \cap B);$$

# Properties of the Probability Measure

7. for any events  $A, B \in \mathcal{F}$

$$\mathbb{P}(B \setminus A) = \mathbb{P}(B) - \mathbb{P}(A \cap B);$$

8. if  $A, B \in \mathcal{F}$  are two events such that  $A \subset B$ , then

$$\mathbb{P}(B \setminus A) = \mathbb{P}(B) - \mathbb{P}(A);$$

# Properties of the Probability Measure

7. for any events  $A, B \in \mathcal{F}$

$$\mathbb{P}(B \setminus A) = \mathbb{P}(B) - \mathbb{P}(A \cap B);$$

8. if  $A, B \in \mathcal{F}$  are two events such that  $A \subset B$ , then

$$\mathbb{P}(B \setminus A) = \mathbb{P}(B) - \mathbb{P}(A);$$

9. if  $A$  and  $B$  are two events such that  $A \subset B$ , then

$$\mathbb{P}(A) \leq \mathbb{P}(B);$$

# Properties of the Probability Measure

7. for any events  $A, B \in \mathcal{F}$

$$\mathbb{P}(B \setminus A) = \mathbb{P}(B) - \mathbb{P}(A \cap B);$$

8. if  $A, B \in \mathcal{F}$  are two events such that  $A \subset B$ , then

$$\mathbb{P}(B \setminus A) = \mathbb{P}(B) - \mathbb{P}(A);$$

9. if  $A$  and  $B$  are two events such that  $A \subset B$ , then

$$\mathbb{P}(A) \leq \mathbb{P}(B);$$

10. for any event  $A \in \mathcal{F}$ ,

$$0 \leq \mathbb{P}(A) \leq 1;$$

# Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?

# Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?

# Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?
- For the coin toss experiment, what is  $\mathbb{P}(H)$ ?



# Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?
- For the coin toss experiment, what is  $\mathbb{P}(H)$ ?
- Is it true that  $\mathbb{P}(\{H\}) = \mathbb{P}(\{T\})$  in the coin toss experiment?

# Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?
- For the coin toss experiment, what is  $\mathbb{P}(H)$ ?
- Is it true that  $\mathbb{P}(\{H\}) = \mathbb{P}(\{T\})$  in the coin toss experiment?
- What is  $\mathbb{P}(\overline{A})$ ?

# Some Questions to Answer

- What is the Domain of Definition of the Probability Measure?
- What is a Probability Measure?
- For the coin toss experiment, what is  $\mathbb{P}(H)$ ?
- Is it true that  $\mathbb{P}(\{H\}) = \mathbb{P}(\{T\})$  in the coin toss experiment?
- What is  $\mathbb{P}(\overline{A})$ ?
- What is  $\mathbb{P}(A \cup B)$ ?

# Example

Assume that in some university, for the CS department, the probability that a sophomore student knows Java programming is 50%, and the probability that he/she knows C++ programming is 60%.

# Example

Assume that in some university, for the CS department, the probability that a sophomore student knows Java programming is 50%, and the probability that he/she knows C++ programming is 60%. We consider the following Events:

$Java$  = Student knows Java programming

$C++$  = Student knows C++ programming.

Then

$$\mathbb{P}(Java) = 0.5 \quad \text{and} \quad \mathbb{P}(C++) = 0.6.$$

# Example

Assume that in some university, for the CS department, the probability that a sophomore student knows Java programming is 50%, and the probability that he/she knows C++ programming is 60%. We consider the following Events:

$Java$  = Student knows Java programming

$C++$  = Student knows C++ programming.

Then

$$\mathbb{P}(Java) = 0.5 \quad \text{and} \quad \mathbb{P}(C++) = 0.6.$$

- Is it possible that no student with Java knowledge knows C++ programming?

# Example

Assume that in some university, for the CS department, the probability that a sophomore student knows Java programming is 50%, and the probability that he/she knows C++ programming is 60%. We consider the following Events:

$Java$  = Student knows Java programming

$C++$  = Student knows C++ programming.

Then

$$\mathbb{P}(Java) = 0.5 \quad \text{and} \quad \mathbb{P}(C++) = 0.6.$$

- Is it possible that no student with Java knowledge knows C++ programming? I.e., can  $Java$  and  $C++$  be mutually exclusive?

# Example, cont'd

Recall that

$$\mathbb{P}(\textit{Java}) = 0.5 \quad \text{and} \quad \mathbb{P}(\textit{C}++) = 0.6.$$



# Example, cont'd

Recall that

$$\mathbb{P}(\text{Java}) = 0.5 \quad \text{and} \quad \mathbb{P}(C++) = 0.6.$$

- What is the probability that a student knows Java or C++ programming?

# Example, cont'd

Recall that

$$\mathbb{P}(\text{Java}) = 0.5 \quad \text{and} \quad \mathbb{P}(\text{C}++) = 0.6.$$

- What is the probability that a student knows Java or C++ programming?
- If we assume that the probability of knowing both Java and C++ programming is 25% for the student, what is the probability that the student knows Java or C++ programming?

# Example, cont'd

Recall that

$$\mathbb{P}(\text{Java}) = 0.5 \quad \text{and} \quad \mathbb{P}(\text{C++}) = 0.6.$$

- What is the probability that a student knows Java or C++ programming?
- If we assume that the probability of knowing both Java and C++ programming is 25% for the student, what is the probability that the student knows Java or C++ programming?
- And what is the probability that a student doesn't know neither Java nor C++?

# Example, cont'd

Recall that

$$\mathbb{P}(\text{Java}) = 0.5 \quad \text{and} \quad \mathbb{P}(\text{C++}) = 0.6.$$

- What is the probability that a student knows Java or C++ programming?
- If we assume that the probability of knowing both Java and C++ programming is 25% for the student, what is the probability that the student knows Java or C++ programming?
- And what is the probability that a student doesn't know neither Java nor C++?
- What is the probability that the student knows exactly one of these languages?

# Example, cont'd

Recall that

$$\mathbb{P}(\textit{Java}) = 0.5 \quad \text{and} \quad \mathbb{P}(C++ ) = 0.6.$$

# Example, cont'd

Recall that

$$\mathbb{P}(\text{Java}) = 0.5 \quad \text{and} \quad \mathbb{P}(C++) = 0.6.$$

- In our initial problem, what is the highest possible probability for a student to know Java and C++ programming?

# Example, cont'd

Recall that

$$\mathbb{P}(\text{Java}) = 0.5 \quad \text{and} \quad \mathbb{P}(C++) = 0.6.$$

- In our initial problem, what is the highest possible probability for a student to know Java and C++ programming?
- In our initial problem, what is the smallest possible probability for a student to know Java and C++ programming?

# Classical Probability Models: Discrete Sample Spaces

Discrete = Finite or Countably Infinite



# Classical Probability Models: Discrete Sample Spaces

Discrete = Finite or Countably Infinite

Here we give the general construction of the Probability Models for the Discrete Sample Spaces.

# Classical Probability Models: Discrete Sample Spaces

Discrete = Finite or Countably Infinite

Here we give the general construction of the Probability Models for the Discrete Sample Spaces. The continuous case is much more complicated.

# Classical Probability Models: Discrete Sample Spaces

Discrete = Finite or Countably Infinite

Here we give the general construction of the Probability Models for the Discrete Sample Spaces. The continuous case is much more complicated.

To give Probability Models, Probability Spaces, we need to give:

# Classical Probability Models: Discrete Sample Spaces

Discrete = Finite or Countably Infinite

Here we give the general construction of the Probability Models for the Discrete Sample Spaces. The continuous case is much more complicated.

To give Probability Models, Probability Spaces, we need to give:

- The Sample Space  $\Omega$ ;

# Classical Probability Models: Discrete Sample Spaces

Discrete = Finite or Countably Infinite

Here we give the general construction of the Probability Models for the Discrete Sample Spaces. The continuous case is much more complicated.

To give Probability Models, Probability Spaces, we need to give:

- The Sample Space  $\Omega$ ;
- The set of Events  $\mathcal{F}$ ;

# Classical Probability Models: Discrete Sample Spaces

Discrete = Finite or Countably Infinite

Here we give the general construction of the Probability Models for the Discrete Sample Spaces. The continuous case is much more complicated.

To give Probability Models, Probability Spaces, we need to give:

- The Sample Space  $\Omega$ ;
- The set of Events  $\mathcal{F}$ ;
- The Probability Measure  $\mathbb{P}$ .

# Classical Probability Models: Finite Sample Spaces

Assume the Sample Space  $\Omega$  is finite:

# Classical Probability Models: Finite Sample Spaces

Assume the Sample Space  $\Omega$  is finite:

- Our Sample Space is  $\Omega = \{\omega_1, \omega_2, \dots, \omega_n\}$ .



# Classical Probability Models: Finite Sample Spaces

Assume the Sample Space  $\Omega$  is finite:

- Our Sample Space is  $\Omega = \{\omega_1, \omega_2, \dots, \omega_n\}$ .
- Every subset of  $\Omega$  is an Event, i.e.,  $\mathcal{F} = 2^\Omega$ , the power set of  $\Omega$ .

# Classical Probability Models: Finite Sample Spaces

Assume the Sample Space  $\Omega$  is finite:

- Our Sample Space is  $\Omega = \{\omega_1, \omega_2, \dots, \omega_n\}$ .
- Every subset of  $\Omega$  is an Event, i.e.,  $\mathcal{F} = 2^\Omega$ , the power set of  $\Omega$ .
- We take any real numbers  $p_1, p_2, \dots, p_n$  with

$$p_1 \geq 0, \dots, p_n \geq 0, \quad p_1 + p_2 + \dots + p_n = 1,$$

and define

$$\mathbb{P}(\{\omega_1\}) = p_1, \quad \mathbb{P}(\{\omega_2\}) = p_2, \quad \dots, \quad \mathbb{P}(\{\omega_n\}) = p_n.$$

# Classical Probability Models: Finite Sample Spaces

We write this in a more convenient table form:

Outcome	$\omega_1$	$\omega_2$	$\dots$	$\omega_n$
$\mathbb{P}(\{\omega_k\})$	$p_1$	$p_2$	$\dots$	$p_n$

# Classical Probability Models: Finite Sample Spaces

We write this in a more convenient table form:

Outcome	$\omega_1$	$\omega_2$	$\dots$	$\omega_n$
$\mathbb{P}(\{\omega_k\})$	$p_1$	$p_2$	$\dots$	$p_n$

We are not done yet!

# Classical Probability Models: Finite Sample Spaces

We write this in a more convenient table form:

Outcome	$\omega_1$	$\omega_2$	$\dots$	$\omega_n$
$\mathbb{P}(\{\omega_k\})$	$p_1$	$p_2$	$\dots$	$p_n$

We are not done yet! We define, for any event  $A$ ,

$$\mathbb{P}(A) = \sum_{\omega_i \in A} p_i,$$

and also add  $\mathbb{P}(\emptyset) = 0$ .

# Classical Probability Models: Finite Sample Spaces

We write this in a more convenient table form:

Outcome	$\omega_1$	$\omega_2$	$\dots$	$\omega_n$
$\mathbb{P}(\{\omega_k\})$	$p_1$	$p_2$	$\dots$	$p_n$

We are not done yet! We define, for any event  $A$ ,

$$\mathbb{P}(A) = \sum_{\omega_i \in A} p_i,$$

and also add  $\mathbb{P}(\emptyset) = 0$ .

I FSYO!

# Some Questions to Answer

- What is the Inclusion-Exclusion Principle?

# Some Questions to Answer

- What is the Inclusion-Exclusion Principle?
- If we have  $\mathbb{P}(A)$  and  $\mathbb{P}(B)$ , what is the lower bound for  $\mathbb{P}(A \cup B)$ ?



# Some Questions to Answer

- What is the Inclusion-Exclusion Principle?
- If we have  $\mathbb{P}(A)$  and  $\mathbb{P}(B)$ , what is the lower bound for  $\mathbb{P}(A \cup B)$ ?
- Express  $\mathbb{P}(\overline{A} \cup \overline{B})$  in terms of  $\mathbb{P}(A)$ ,  $\mathbb{P}(B)$  and  $\mathbb{P}(A \cup B)$

# Some Questions to Answer

- What is the Inclusion-Exclusion Principle?
- If we have  $\mathbb{P}(A)$  and  $\mathbb{P}(B)$ , what is the lower bound for  $\mathbb{P}(A \cup B)$ ?
- Express  $\mathbb{P}(\overline{A} \cup \overline{B})$  in terms of  $\mathbb{P}(A)$ ,  $\mathbb{P}(B)$  and  $\mathbb{P}(A \cup B)$
- What we need to give (to describe) a Probability Model?

# Some Questions to Answer

- What is the Inclusion-Exclusion Principle?
- If we have  $\mathbb{P}(A)$  and  $\mathbb{P}(B)$ , what is the lower bound for  $\mathbb{P}(A \cup B)$ ?
- Express  $\mathbb{P}(\overline{A} \cup \overline{B})$  in terms of  $\mathbb{P}(A)$ ,  $\mathbb{P}(B)$  and  $\mathbb{P}(A \cup B)$
- What we need to give (to describe) a Probability Model?
- How to define the Probability Measure in the Finite Sample Space Case?

# Some Questions to Answer

- What is the Inclusion-Exclusion Principle?
- If we have  $\mathbb{P}(A)$  and  $\mathbb{P}(B)$ , what is the lower bound for  $\mathbb{P}(A \cup B)$ ?
- Express  $\mathbb{P}(\overline{A} \cup \overline{B})$  in terms of  $\mathbb{P}(A)$ ,  $\mathbb{P}(B)$  and  $\mathbb{P}(A \cup B)$
- What we need to give (to describe) a Probability Model?
- How to define the Probability Measure in the Finite Sample Space Case?
- How many Probability Models we can have for the Experiment with the Sample Space  $\Omega = \{a, b, c\}$  ?

# Some Questions to Answer

- What is the Inclusion-Exclusion Principle?
- If we have  $\mathbb{P}(A)$  and  $\mathbb{P}(B)$ , what is the lower bound for  $\mathbb{P}(A \cup B)$ ?
- Express  $\mathbb{P}(\overline{A} \cup \overline{B})$  in terms of  $\mathbb{P}(A)$ ,  $\mathbb{P}(B)$  and  $\mathbb{P}(A \cup B)$
- What we need to give (to describe) a Probability Model?
- How to define the Probability Measure in the Finite Sample Space Case?
- How many Probability Models we can have for the Experiment with the Sample Space  $\Omega = \{a, b, c\}$  ? Describe them!