

CS 107, Probability, Spring 2019

Lecture 44

Michael Poghosyan

AUA

- 2 lectures

- Intro to Markov Chains

Intro to Markov Chains

Definition of the Markov Chain

Recall the Definition of a Markov Chain:

Definition of the Markov Chain

Recall the Definition of a Markov Chain:

Markov Chain

Assume X_0, X_1, X_2, \dots is a sequence of r.v.s (Discrete Stochastic Process), which take values from $S = \{1, 2, \dots, N\}$. We say that $X_n, n = 0, 1, 2, \dots$ is a (Finite State Discrete Time) **Markov Chain**, if

$$\mathbb{P}(X_{t+1} = j | X_t = i, X_{t-1} = k, \dots, X_0 = m) = \mathbb{P}(X_{t+1} = j | X_t = i)$$

for any time t , for any state j, i, k, \dots, m .

Definition of the Markov Chain

Recall the Definition of a Markov Chain:

Markov Chain

Assume X_0, X_1, X_2, \dots is a sequence of r.v.s (Discrete Stochastic Process), which take values from $S = \{1, 2, \dots, N\}$. We say that $X_n, n = 0, 1, 2, \dots$ is a (Finite State Discrete Time) **Markov Chain**, if

$$\mathbb{P}(X_{t+1} = j | X_t = i, X_{t-1} = k, \dots, X_0 = m) = \mathbb{P}(X_{t+1} = j | X_t = i)$$

for any time t , for any state j, i, k, \dots, m .

Interpretation: Given today's State, tomorrow's State is independent of the past States.

Definition of the Markov Chain

Recall the Definition of a Markov Chain:

Markov Chain

Assume X_0, X_1, X_2, \dots is a sequence of r.v.s (Discrete Stochastic Process), which take values from $S = \{1, 2, \dots, N\}$. We say that $X_n, n = 0, 1, 2, \dots$ is a (Finite State Discrete Time) **Markov Chain**, if

$$\mathbb{P}(X_{t+1} = j | X_t = i, X_{t-1} = k, \dots, X_0 = m) = \mathbb{P}(X_{t+1} = j | X_t = i)$$

for any time t , for any state j, i, k, \dots, m .

Interpretation: Given today's State, tomorrow's State is independent of the past States. Or, in other words, today's information is enough to completely determine the probabilities of Tomorrow's states.

Definition of the Markov Chain

We will consider only Time-Homogeneous Markov Chains (MC).

We will say that our MC is Time-Homogeneous, if

$$\mathbb{P}(X_{t+1} = j | X_t = i) \text{ does't depend on } t.$$

Definition of the Markov Chain

We will consider only Time-Homogeneous Markov Chains (MC).
We will say that our MC is Time-Homogeneous, if

$$\mathbb{P}(X_{t+1} = j | X_t = i) \text{ does't depend on } t.$$

In this case we denote

$$\mathbb{P}(X_{t+1} = j | X_t = i) = p_{ij}, \quad i, j \in S,$$

and call p_{ij} the Transition Probability from the State i to State j .

Definition of the Markov Chain

We will consider only Time-Homogeneous Markov Chains (MC).
We will say that our MC is Time-Homogeneous, if

$$\mathbb{P}(X_{t+1} = j | X_t = i) \text{ does't depend on } t.$$

In this case we denote

$$\mathbb{P}(X_{t+1} = j | X_t = i) = p_{ij}, \quad i, j \in S,$$

and call p_{ij} the Transition Probability from the State i to State j . So in the Time-Homogeneous MC, the Probability to move from the State i to State j does not depend on the time.

Definition of the Markov Chain

We will consider only Time-Homogeneous Markov Chains (MC).
We will say that our MC is Time-Homogeneous, if

$$\mathbb{P}(X_{t+1} = j | X_t = i) \text{ doesn't depend on } t.$$

In this case we denote

$$\mathbb{P}(X_{t+1} = j | X_t = i) = p_{ij}, \quad i, j \in S,$$

and call p_{ij} the Transition Probability from the State i to State j . So in the Time-Homogeneous MC, the Probability to move from the State i to State j does not depend on the time.

Clearly,

$$p_{ij} \geq 0, \quad \text{and} \quad \sum_{j=1}^N p_{ij} = 1.$$

Definition of the Markov Chain

We will consider only Time-Homogeneous Markov Chains (MC).
We will say that our MC is Time-Homogeneous, if

$$\mathbb{P}(X_{t+1} = j | X_t = i) \text{ doesn't depend on } t.$$

In this case we denote

$$\mathbb{P}(X_{t+1} = j | X_t = i) = p_{ij}, \quad i, j \in S,$$

and call p_{ij} the Transition Probability from the State i to State j . So in the Time-Homogeneous MC, the Probability to move from the State i to State j does not depend on the time.

Clearly,

$$p_{ij} \geq 0, \quad \text{and} \quad \sum_{j=1}^N p_{ij} = 1.$$

The $N \times N$ Matrix $P = [p_{ij}]$ is called the Transition Probability Matrix. Row sums of P are equal to 1.

Describing the Markov Chain/Model

So to give the Markov Chain (Markov Model), one needs to specify:

Describing the Markov Chain/Model

So to give the Markov Chain (Markov Model), one needs to specify:

- The State Space of the Model, $S = \{1, 2, \dots, N\}$;

Describing the Markov Chain/Model

So to give the Markov Chain (Markov Model), one needs to specify:

- The State Space of the Model, $S = \{1, 2, \dots, N\}$;
- The Transition Probability Matrix, $P = [p_{ij}]_{i,j=1,\dots,N}$.

Describing the Markov Chain/Model

So to give the Markov Chain (Markov Model), one needs to specify:

- The State Space of the Model, $S = \{1, 2, \dots, N\}$;
- The Transition Probability Matrix, $P = [p_{ij}]_{i,j=1,\dots,N}$.

Note: Sometimes one adds to the Model also the Initial Probabilities $\{\pi_i : i = 1, \dots, N\}$, where π_i is the Probability that we start at the state i :

$$\mathbb{P}(X_0 = i) = \pi_i, \quad i \in S.$$

Describing the Markov Chain/Model

So to give the Markov Chain (Markov Model), one needs to specify:

- The State Space of the Model, $S = \{1, 2, \dots, N\}$;
- The Transition Probability Matrix, $P = [p_{ij}]_{i,j=1,\dots,N}$.

Note: Sometimes one adds to the Model also the Initial Probabilities $\{\pi_i : i = 1, \dots, N\}$, where π_i is the Probability that we start at the state i :

$$\mathbb{P}(X_0 = i) = \pi_i, \quad i \in S.$$

Of course, in this case we need to have $\sum_{j=1}^N \pi_i = 1$.

Markov Chain Transition Probability Graph

Sometimes people describe the Markov Chain by drawing the Transition Probability Graph: OTB

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game.

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game.

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative.

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative. So we describe our Markov Chain:

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative.

So we describe our Markov Chain:

- State Space = $S = \{0, 1, 2, \dots\}$;

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative.

So we describe our Markov Chain:

- State Space = $S = \{0, 1, 2, \dots\}$;
- Transition Probabilities:

$$p_{00} =$$

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative.

So we describe our Markov Chain:

- State Space = $S = \{0, 1, 2, \dots\}$;
- Transition Probabilities:

$$p_{00} = 1, \quad p_{i,j}$$

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative.

So we describe our Markov Chain:

- State Space = $S = \{0, 1, 2, \dots\}$;
- Transition Probabilities:

$$p_{00} = 1, \quad p_{i,j} = 0, \text{ if } j \neq i \pm 1, \quad \text{and } (i,j) \neq (0,0)$$

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative.

So we describe our Markov Chain:

- State Space = $S = \{0, 1, 2, \dots\}$;
- Transition Probabilities:

$$p_{00} = 1, \quad p_{i,j} = 0, \text{ if } j \neq i \pm 1, \quad \text{and } (i,j) \neq (0,0)$$

$$p_{i,i-1} =$$

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative.

So we describe our Markov Chain:

- State Space = $S = \{0, 1, 2, \dots\}$;
- Transition Probabilities:

$$p_{00} = 1, \quad p_{i,j} = 0, \text{ if } j \neq i \pm 1, \quad \text{and } (i,j) \neq (0,0)$$

$$p_{i,i-1} = \frac{20}{38}, \quad p_{i,i+1} =$$

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative.

So we describe our Markov Chain:

- State Space = $S = \{0, 1, 2, \dots\}$;
- Transition Probabilities:

$$p_{00} = 1, \quad p_{i,j} = 0, \text{ if } j \neq i \pm 1, \quad \text{and } (i,j) \neq (0,0)$$

$$p_{i,i-1} = \frac{20}{38}, \quad p_{i,i+1} = \frac{18}{38}$$

Example:

Example: American Roulette: Assume I am playing an American Roulette. At each round, I bet \$1 on either red or black. Let X_t be the amount of my money after the t -th game. X_0 is the money amount I have before starting the game. Clearly here, X_t depends on X_{t-1} , and given X_{t-1} , the values of X_{t-2}, X_{t-3}, \dots are not informative.

So we describe our Markov Chain:

- State Space = $S = \{0, 1, 2, \dots\}$;
- Transition Probabilities:

$$p_{00} = 1, \quad p_{i,j} = 0, \text{ if } j \neq i \pm 1, \quad \text{and } (i,j) \neq (0,0)$$

$$p_{i,i-1} = \frac{20}{38}, \quad p_{i,i+1} = \frac{18}{38}$$

Give the Transition Probability Graph OTB!

Example

Example: On/Off System Assume we have some system, which can be in two states, over some discrete time interval: ON and OFF.

Example

Example: On/Off System Assume we have some system, which can be in two states, over some discrete time interval: ON and OFF. If the System is in the State OFF at some time instant, then, at the next instant it will be at the State ON is p .

Example

Example: On/Off System Assume we have some system, which can be in two states, over some discrete time interval: ON and OFF. If the System is in the State OFF at some time instant, then, at the next instant it will be at the State ON is p . Also, if the System is in the State ON at some time instant, then, at the next instant it will be at the State OFF is q .

Example

Example: On/Off System Assume we have some system, which can be in two states, over some discrete time interval: ON and OFF. If the System is in the State OFF at some time instant, then, at the next instant it will be at the State ON is p . Also, if the System is in the State ON at some time instant, then, at the next instant it will be at the State OFF is q . Describe this System's states as a Markov Chain.

Example

Example: On/Off System Assume we have some system, which can be in two states, over some discrete time interval: ON and OFF. If the System is in the State OFF at some time instant, then, at the next instant it will be at the State ON is p . Also, if the System is in the State ON at some time instant, then, at the next instant it will be at the State OFF is q . Describe this System's states as a Markov Chain.

Solution: OTB

Example

Example: Thief and Policemen Assume our policemen are situated at the points 0 and N of the real line, and are not moving.

Example

Example: Thief and Policemen Assume our policemen are situated at the points 0 and N of the real line, and are not moving. Our thief, after the successful "operation", is somewhere at the points $1, 2, 3, \dots, N - 1$.

Example

Example: Thief and Policemen Assume our policemen are situated at the points 0 and N of the real line, and are not moving. Our thief, after the successful "operation", is somewhere at the points $1, 2, 3, \dots, N - 1$. He moves along that line with unit moves. At each time instant, he moves 1 units to the right with probability 0.3, to the left with the same probability, and stays at its place with probability 0.4, independent of the previous steps.

Example

Example: Thief and Policemen Assume our policemen are situated at the points 0 and N of the real line, and are not moving. Our thief, after the successful "operation", is somewhere at the points $1, 2, 3, \dots, N - 1$. He moves along that line with unit moves. At each time instant, he moves 1 units to the right with probability 0.3, to the left with the same probability, and stays at its place with probability 0.4, independent of the previous steps.

Describe the position of the Thief as a Markov Chain.

Example

Example: Thief and Policemen Assume our policemen are situated at the points 0 and N of the real line, and are not moving. Our thief, after the successful "operation", is somewhere at the points $1, 2, 3, \dots, N - 1$. He moves along that line with unit moves. At each time instant, he moves 1 units to the right with probability 0.3, to the left with the same probability, and stays at its place with probability 0.4, independent of the previous steps.

Describe the position of the Thief as a Markov Chain.

Solution: OTB

Example

Other Examples:

Example

Other Examples:

- Monopoly game,

Other Examples:

- Monopoly game, see, e.g., <https://faculty.math.illinois.edu/~bishop/monopoly.pdf> or <http://www.bewersdorff-online.de/amonopoly/> or <http://www.math.yorku.ca/Who/Faculty/Steprans/Courses/2042/Monopoly/Stewart4.html>

Example

Other Examples:

- Monopoly game, see, e.g., <https://faculty.math.illinois.edu/~bishop/monopoly.pdf> or <http://www.bewersdorff-online.de/amonopoly/> or <http://www.math.yorku.ca/Who/Faculty/Steprans/Courses/2042/Monopoly/Stewart4.html>
- Games of type "Snakes and Ladders", see https://en.wikipedia.org/wiki/Snakes_and_Ladders

Example

Other Examples:

- Monopoly game, see, e.g., <https://faculty.math.illinois.edu/~bishop/monopoly.pdf> or <http://www.bewersdorff-online.de/amonopoly/> or <http://www.math.yorku.ca/Who/Faculty/Steprans/Courses/2042/Monopoly/Stewart4.html>
- Games of type "Snakes and Ladders", see https://en.wikipedia.org/wiki/Snakes_and_Ladders
- Baseball game: just Google "baseball analysis Markov Chains"

Example

Other Examples:

- Monopoly game, see, e.g., <https://faculty.math.illinois.edu/~bishop/monopoly.pdf> or <http://www.bewersdorff-online.de/amonopoly/> or <http://www.math.yorku.ca/Who/Faculty/Steprans/Courses/2042/Monopoly/Stewart4.html>
- Games of type "Snakes and Ladders", see https://en.wikipedia.org/wiki/Snakes_and_Ladders
- Baseball game: just Google "baseball analysis Markov Chains"
- And Google itself is based on Markov Chains!

Example

Other Examples:

- Monopoly game, see, e.g., <https://faculty.math.illinois.edu/~bishop/monopoly.pdf> or <http://www.bewersdorff-online.de/amonopoly/> or <http://www.math.yorku.ca/Who/Faculty/Steprans/Courses/2042/Monopoly/Stewart4.html>
- Games of type "Snakes and Ladders", see https://en.wikipedia.org/wiki/Snakes_and_Ladders
- Baseball game: just Google "baseball analysis Markov Chains"
- And Google itself is based on Markov Chains! The heart and idea of the Google is PageRank Algorithm developed by Larry Page and Sergei Brin, who later founded Google. See <https://en.wikipedia.org/wiki/PageRank> or again just Google "PageRank Markov Chain"

Example

Other Examples:

Other Examples:

- Markov used his chains to describe the alternation of consonants and vowels in Pushkin's Eugene Onegin, see <https://www.jstor.org/stable/43707006> or the original at <http://books.e-heritage.ru/book/10086570>

Other Examples:

- Markov used his chains to describe the alternation of consonants and vowels in Pushkin's Eugene Onegin, see <https://www.jstor.org/stable/43707006> or the original at <http://books.e-heritage.ru/book/10086570>
- Random Walks are examples of Markov Chains

Other Examples:

- Markov used his chains to describe the alternation of consonants and vowels in Pushkin's Eugene Onegin, see <https://www.jstor.org/stable/43707006> or the original at <http://books.e-heritage.ru/book/10086570>
- Random Walks are examples of Markov Chains
- Ideas of Markov Chains and Processes are important in Language Models (n -gram Models), Machine Learning (HMM, Hidden Markov Models), Reinforcement Learning (Markov Decision Processes), in Simulations (MCMC, Markov Chain Monte Carlo) etc.

Some Problems Concerning Markov Chains

Now assume we have a Markov Chain (S, P) , where $S = \{1, 2, \dots, N\}$ is the set of possible States, and $P = [p_{ij}]$ is the Transition Matrix of our MC.

Some Problems Concerning Markov Chains

Now assume we have a Markov Chain (S, P) , where $S = \{1, 2, \dots, N\}$ is the set of possible States, and $P = [p_{ij}]$ is the Transition Matrix of our MC.

One of the problems people solve is to find the Probability of some State Path:

$$\mathbb{P}(X_0 = i_0, X_1 = i_1, \dots, X_n = i_n).$$

Some Problems Concerning Markov Chains

Now assume we have a Markov Chain (S, P) , where $S = \{1, 2, \dots, N\}$ is the set of possible States, and $P = [p_{ij}]$ is the Transition Matrix of our MC.

One of the problems people solve is to find the Probability of some State Path:

$$\mathbb{P}(X_0 = i_0, X_1 = i_1, \dots, X_n = i_n).$$

It is easy to see, using the Markov property, that this can be calculated as:

$$\begin{aligned} & \mathbb{P}(X_0 = i_0, X_1 = i_1, \dots, X_n = i_n) = \\ & \mathbb{P}(X_0 = i_0) \cdot \mathbb{P}(X_1 = i_1 | X_0 = i_0) \cdot \mathbb{P}(X_2 = i_2 | X_1 = i_1) \cdot \dots \cdot \mathbb{P}(X_n = i_n | X_{n-1} = i_{n-1}) \\ & = \mathbb{P}(X_0 = i_0) \cdot p_{i_0 i_1} \cdot p_{i_1 i_2} \cdot \dots \cdot p_{i_{n-1} i_n}. \end{aligned}$$

Some Problems Concerning Markov Chains

To calculate using the above formula, we need to either have the initial position, say, our System is at the State i_0 at time 0, and, in that case, $\mathbb{P}(X_0 = i_0) = 1$,

Some Problems Concerning Markov Chains

To calculate using the above formula, we need to either have the initial position, say, our System is at the State i_0 at time 0, and, in that case, $\mathbb{P}(X_0 = i_0) = 1$, or we need to have the initial probabilities $\mathbb{P}(X_0 = i)$ for any i .

Some Problems Concerning Markov Chains

To calculate using the above formula, we need to either have the initial position, say, our System is at the State i_0 at time 0, and, in that case, $\mathbb{P}(X_0 = i_0) = 1$, or we need to have the initial probabilities $\mathbb{P}(X_0 = i)$ for any i .

Having the ability to calculate the Probabilities of Paths

$$\mathbb{P}(X_0 = i_0, X_1 = i_1, \dots, X_n = i_n),$$

we can calculate, say, what is the most probable path of movement, say, for the next 3 time instants or what is the most probable state at time $t = 3$.