

CS 107, Probability, Spring 2019

Lecture 08

Michael Poghosyan

AUA

4 February 2019

- Classical Probability Models: Finite Sample Spaces with Equally Likely Outcomes = Combinatorial Problems, Cont'd
- Classical Probability Models: Geometric Probabilities

The Monty Hall Problem

<https://www.youtube.com/watch?v=mhlc7peG1Gg>

Example: Birthday Problem

Problem: We have 36 participants in our group of Probability class, including the instructor. What is the probability that at least two participants share the same birthday?

Example: Birthday Problem

Problem: We have 36 participants in our group of Probability class, including the instructor. What is the probability that at least two participants share the same birthday?

Assumptions: There are 365 days in a year,

Example: Birthday Problem

Problem: We have 36 participants in our group of Probability class, including the instructor. What is the probability that at least two participants share the same birthday?

Assumptions: There are 365 days in a year, and the probability of being born on each day is the same.

Example: Birthday Problem, another version

Problem: We have 36 participants in our group of Probability class, including the instructor. What is the probability that at least one of our students will share the instructor's (MP's) birthday?

Example: Julius Caesar Problem

Do you remember our Caesar?

Example: Julius Caesar Problem

Do you remember our Caesar?



Figure: Gaius Julius Caesar

Example: Julius Caesar Problem

Do you remember our Caesar?

Example: Julius Caesar Problem

Do you remember our Caesar?

Problem: What are the chances that I have just inhaled a molecule which the great Caesar exhaled when saying his last words?

Example: Julius Caesar Problem

Do you remember our Caesar?

Problem: What are the chances that I have just inhaled a molecule which the great Caesar exhaled when saying his last words?

Assume we have N molecules of air in the world, and Caesar exhaled n of them, and they are uniformly spread over the world.

Example: Julius Caesar Problem

Do you remember our Caesar?

Problem: What are the chances that I have just inhaled a molecule which the great Caesar exhaled when saying his last words?

Assume we have N molecules of air in the world, and Caesar exhaled n of them, and they are uniformly spread over the world. And assume I have just inhaled m molecules.

Example: Julius Caesar Problem

Do you remember our Caesar?

Problem: What are the chances that I have just inhaled a molecule which the great Caesar exhaled when saying his last words?

Assume we have N molecules of air in the world, and Caesar exhaled n of them, and they are uniformly spread over the world. And assume I have just inhaled m molecules.

Physics says that $N \sim 10^{44}$, $n, m \sim 2.2 \cdot 10^{22}$

The rest on the board!

Girls and Boys Problem

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Girls and Boys Problem

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 1: We model the outcome to be the number of b/g child in the family:

$$\Omega = \{3g, 2g1b, 1g2b, 3b\},$$

Girls and Boys Problem

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 1: We model the outcome to be the number of b/g child in the family:

$$\Omega = \{3g, 2g1b, 1g2b, 3b\},$$

The Favorable Event is:

$$A = \{1g2b\}$$

So,

$$\mathbb{P}(A) =$$

Girls and Boys Problem

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 1: We model the outcome to be the number of b/g child in the family:

$$\Omega = \{3g, 2g1b, 1g2b, 3b\},$$

The Favorable Event is:

$$A = \{1g2b\}$$

So,

$$\mathbb{P}(A) = \frac{1}{4}.$$

Girls and Boys Problem

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 1: We model the outcome to be the number of b/g child in the family:

$$\Omega = \{3g, 2g1b, 1g2b, 3b\},$$

The Favorable Event is:

$$A = \{1g2b\}$$

So,

$$\mathbb{P}(A) = \frac{1}{4}.$$

Correct?

Girls and Boys Problem

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 1: We model the outcome to be the number of b/g child in the family:

$$\Omega = \{3g, 2g1b, 1g2b, 3b\},$$

The Favorable Event is:

$$A = \{1g2b\}$$

So,

$$\mathbb{P}(A) = \frac{1}{4}.$$

Correct? No!

Girls and Boys Problem, 2nd Sol'n

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Girls and Boys Problem, 2nd Sol'n

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 2: We model the outcome to be which child is b/g in the family:

$$\Omega = \{ggg, ggb, gb g, bgg, gbb, bgb, bbg, bbb\}.$$

Girls and Boys Problem, 2nd Sol'n

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 2: We model the outcome to be which child is b/g in the family:

$$\Omega = \{ggg, ggb, gbg, bgg, gbb, bgb, bbg, bbb\}.$$

The Favorable Event is:

$$A = \{gbb, bgb, bbg\}.$$

So,

$$\mathbb{P}(A) =$$

Girls and Boys Problem, 2nd Sol'n

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 2: We model the outcome to be which child is b/g in the family:

$$\Omega = \{ggg, ggb, gb g, bgg, gbb, bgb, bbg, bbb\}.$$

The Favorable Event is:

$$A = \{gbb, bgb, bbg\}.$$

So,

$$\mathbb{P}(A) = \frac{3}{8}.$$

Girls and Boys Problem, 2nd Sol'n

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 2: We model the outcome to be which child is b/g in the family:

$$\Omega = \{ggg, ggb, gbg, bgg, gbb, bgb, bbg, bbb\}.$$

The Favorable Event is:

$$A = \{gbb, bgb, bbg\}.$$

So,

$$\mathbb{P}(A) = \frac{3}{8}.$$

Correct?

Girls and Boys Problem, 2nd Sol'n

Problem: We randomly choose a family with three children. Find the probability that they have exactly one daughter.

Sol'n 2: We model the outcome to be which child is b/g in the family:

$$\Omega = \{ggg, ggb, gb g, bgg, gbb, bgb, bbg, bbb\}.$$

The Favorable Event is:

$$A = \{gbb, bgb, bbg\}.$$

So,

$$\mathbb{P}(A) = \frac{3}{8}.$$

Correct? Aha!

Geometric Probabilities

Now let us talk about Experiments, where the Sample Space is some non-countable infinite subset of \mathbb{R}^n for some $n \geq 1$, say, the Sample Space can be

Geometric Probabilities

Now let us talk about Experiments, where the Sample Space is some non-countable infinite subset of \mathbb{R}^n for some $n \geq 1$, say, the Sample Space can be

- an interval in \mathbb{R} ;

Geometric Probabilities

Now let us talk about Experiments, where the Sample Space is some non-countable infinite subset of \mathbb{R}^n for some $n \geq 1$, say, the Sample Space can be

- an interval in \mathbb{R} ; e.g., No. 2 marshutka waiting time when arriving at the bus stop;

Geometric Probabilities

Now let us talk about Experiments, where the Sample Space is some non-countable infinite subset of \mathbb{R}^n for some $n \geq 1$, say, the Sample Space can be

- an interval in \mathbb{R} ; e.g., No. 2 marshutka waiting time when arriving at the bus stop;
- a circle or a rectangle in \mathbb{R}^2 ;

Geometric Probabilities

Now let us talk about Experiments, where the Sample Space is some non-countable infinite subset of \mathbb{R}^n for some $n \geq 1$, say, the Sample Space can be

- an interval in \mathbb{R} ; e.g., No. 2 marshutka waiting time when arriving at the bus stop;
- a circle or a rectangle in \mathbb{R}^2 ; e.g., hitting a Darts board with the missile;

Geometric Probabilities

Now let us talk about Experiments, where the Sample Space is some non-countable infinite subset of \mathbb{R}^n for some $n \geq 1$, say, the Sample Space can be

- an interval in \mathbb{R} ; e.g., No. 2 marshutka waiting time when arriving at the bus stop;
- a circle or a rectangle in \mathbb{R}^2 ; e.g., hitting a Darts board with the missile;
- a cube in \mathbb{R}^3 ;

Geometric Probabilities

Now let us talk about Experiments, where the Sample Space is some non-countable infinite subset of \mathbb{R}^n for some $n \geq 1$, say, the Sample Space can be

- an interval in \mathbb{R} ; e.g., No. 2 marshutka waiting time when arriving at the bus stop;
- a circle or a rectangle in \mathbb{R}^2 ; e.g., hitting a Darts board with the missile;
- a cube in \mathbb{R}^3 ; lifetimes for 3 different parts of the above marshutka;

Geometric Probabilities

Now let us talk about Experiments, where the Sample Space is some non-countable infinite subset of \mathbb{R}^n for some $n \geq 1$, say, the Sample Space can be

- an interval in \mathbb{R} ; e.g., No. 2 marshutka waiting time when arriving at the bus stop;
- a circle or a rectangle in \mathbb{R}^2 ; e.g., hitting a Darts board with the missile;
- a cube in \mathbb{R}^3 ; lifetimes for 3 different parts of the above marshutka;
- an n -dimensional cube in \mathbb{R}^n ,

Geometric Probabilities

Now let us talk about Experiments, where the Sample Space is some non-countable infinite subset of \mathbb{R}^n for some $n \geq 1$, say, the Sample Space can be

- an interval in \mathbb{R} ; e.g., No. 2 marshutka waiting time when arriving at the bus stop;
- a circle or a rectangle in \mathbb{R}^2 ; e.g., hitting a Darts board with the missile;
- a cube in \mathbb{R}^3 ; lifetimes for 3 different parts of the above marshutka;
- an n -dimensional cube in \mathbb{R}^n , arrival times of our Probability Students to our class today (? on Wed?)

We want to give a basic Probabilistic model for our Experiment.

Geometric Probabilities, Cont'd

So assume that our Experiment's Sample Space is $\Omega \subset \mathbb{R}^n$.

Geometric Probabilities, Cont'd

So assume that our Experiment's Sample Space is $\Omega \subset \mathbb{R}^n$.
We need an assumption to define our Classical Geometric Probability Model.

Geometric Probabilities, Cont'd

So assume that our Experiment's Sample Space is $\Omega \subset \mathbb{R}^n$.
We need an assumption to define our Classical Geometric Probability Model.

Assumption

We assume that we have **uniform probability distribution**.

Geometric Probabilities, Cont'd

So assume that our Experiment's Sample Space is $\Omega \subset \mathbb{R}^n$.
We need an assumption to define our Classical Geometric Probability Model.

Assumption

We assume that we have **uniform probability distribution**.

What we mean by that:

- **Incorrect interpretation:** the probability of choosing each point in Ω is the same.

Geometric Probabilities, Cont'd

So assume that our Experiment's Sample Space is $\Omega \subset \mathbb{R}^n$.
We need an assumption to define our Classical Geometric Probability Model.

Assumption

We assume that we have **uniform probability distribution**.

What we mean by that:

- **Incorrect interpretation:** the probability of choosing each point in Ω is the same. Well, in this Geometric Model (non-countably infinite Sample Space!) we will have that the probability of choosing any particular point is 0.
- **Correct Interpretation:** the probability of choosing any equal-measure (length/area/volume) subsets (Events) is the same.

Entracte

Assume we choose a point at random from $[0, 1]$, so $\Omega = [0, 1]$.

Entracte

Assume we choose a point at random from $[0, 1]$, so $\Omega = [0, 1]$. We know that the probability to choose the point 0 is 0. So we will not choose 0 for 100%.

Entracte

Assume we choose a point at random from $[0, 1]$, so $\Omega = [0, 1]$. We know that the probability to choose the point 0 is 0. So we will not choose 0 for 100%. Also, we will not choose the point 1 for 100%.

Entracte

Assume we choose a point at random from $[0, 1]$, so $\Omega = [0, 1]$. We know that the probability to choose the point 0 is 0. So we will not choose 0 for 100%. Also, we will not choose the point 1 for 100%. Similarly, we will not choose the point 0.3241 for sure. The same for 0.6473241321.

Entracte

Assume we choose a point at random from $[0, 1]$, so $\Omega = [0, 1]$. We know that the probability to choose the point 0 is 0. So we will not choose 0 for 100%. Also, we will not choose the point 1 for 100%. Similarly, we will not choose the point 0.3241 for sure. The same for 0.6473241321.

Then which point we will choose if we will not choose any point for sure?

Entracte

Assume we choose a point at random from $[0, 1]$, so $\Omega = [0, 1]$. We know that the probability to choose the point 0 is 0. So we will not choose 0 for 100%. Also, we will not choose the point 1 for 100%. Similarly, we will not choose the point 0.3241 for sure. The same for 0.6473241321.

Then which point we will choose if we will not choose any point for sure?

:D

Geometric Probabilities, Cont'd

Let's continue:

¹In the Lebesgue's sense (18+).

Geometric Probabilities, Cont'd

Let's continue:

- Our Experiment's Sample Space is $\Omega \subset \mathbb{R}^n$;

¹In the Lebesgue's sense (18+).

Geometric Probabilities, Cont'd

Let's continue:

- Our Experiment's Sample Space is $\Omega \subset \mathbb{R}^n$;
- We assume that Ω is measurable¹, i.e., it has a finite length (if $\Omega \subset \mathbb{R}$) or it has a finite area (if $\Omega \subset \mathbb{R}^2$) or it has a finite volume (if $\Omega \subset \mathbb{R}^n$, $n \geq 3$).

¹In the Lebesgue's sense (18+).

Geometric Probabilities, Cont'd

Let's continue:

- Our Experiment's Sample Space is $\Omega \subset \mathbb{R}^n$;
- We assume that Ω is measurable¹, i.e., it has a finite length (if $\Omega \subset \mathbb{R}$) or it has a finite area (if $\Omega \subset \mathbb{R}^2$) or it has a finite volume (if $\Omega \subset \mathbb{R}^n$, $n \geq 3$).
- If $A \subset \Omega$ is an Event (if A has a finite measure), then we define

$$\mathbb{P}(A) = \frac{\text{measure}(A)}{\text{measure}(\Omega)}.$$

¹In the Lebesgue's sense (18+).

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

We (silently) assume that the probabilities are uniformly distributed.

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

We (silently) assume that the probabilities are uniformly distributed.

- What is the probability that I will wait more than 8 min?

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

We (silently) assume that the probabilities are uniformly distributed.

- What is the probability that I will wait more than 8 min?
 - Here our Event is $A = (8, 12]$.

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

We (silently) assume that the probabilities are uniformly distributed.

- What is the probability that I will wait more than 8 min?
 - Here our Event is $A = (8, 12]$.
 - Since Ω is 1D ($\Omega \subset \mathbb{R}$), then *measure* = *length*.

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

We (silently) assume that the probabilities are uniformly distributed.

- What is the probability that I will wait more than 8 min?
 - Here our Event is $A = (8, 12]$.
 - Since Ω is 1D ($\Omega \subset \mathbb{R}$), then *measure* = *length*.
 - So

$$\mathbb{P}(A) =$$

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

We (silently) assume that the probabilities are uniformly distributed.

- What is the probability that I will wait more than 8 min?
 - Here our Event is $A = (8, 12]$.
 - Since Ω is 1D ($\Omega \subset \mathbb{R}$), then *measure* = *length*.
 - So

$$\mathbb{P}(A) = \frac{\text{length}(A)}{\text{length}(\Omega)} = \frac{\text{length}((8, 12])}{\text{length}([0, 12])} = \frac{12 - 8}{12 - 0} = \frac{4}{12} = \frac{1}{3}.$$

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

We (silently) assume that the probabilities are uniformly distributed.

- What is the probability that I will wait more than 8 min?
 - Here our Event is $A = (8, 12]$.
 - Since Ω is 1D ($\Omega \subset \mathbb{R}$), then *measure* = *length*.
 - So

$$\mathbb{P}(A) = \frac{\text{length}(A)}{\text{length}(\Omega)} = \frac{\text{length}((8, 12])}{\text{length}([0, 12])} = \frac{12 - 8}{12 - 0} = \frac{4}{12} = \frac{1}{3}.$$

- What is the probability that I will wait exactly 7.3 min?

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

We (silently) assume that the probabilities are uniformly distributed.

- What is the probability that I will wait more than 8 min?
 - Here our Event is $A = (8, 12]$.
 - Since Ω is 1D ($\Omega \subset \mathbb{R}$), then *measure* = *length*.
 - So

$$\mathbb{P}(A) = \frac{\text{length}(A)}{\text{length}(\Omega)} = \frac{\text{length}((8, 12])}{\text{length}([0, 12])} = \frac{12 - 8}{12 - 0} = \frac{4}{12} = \frac{1}{3}.$$

- What is the probability that I will wait exactly 7.3 min?
 - Here our Event is $A = \{7.3\}$.

Example:

Problem: Assume my waiting time for the train can be anything from $[0, 12]$ (in min), $\Omega = [0, 12]$.

We (silently) assume that the probabilities are uniformly distributed.

- What is the probability that I will wait more than 8 min?
 - Here our Event is $A = (8, 12]$.
 - Since Ω is 1D ($\Omega \subset \mathbb{R}$), then *measure* = *length*.
 - So

$$\mathbb{P}(A) = \frac{\text{length}(A)}{\text{length}(\Omega)} = \frac{\text{length}((8, 12])}{\text{length}([0, 12])} = \frac{12 - 8}{12 - 0} = \frac{4}{12} = \frac{1}{3}.$$

- What is the probability that I will wait exactly 7.3 min?
 - Here our Event is $A = \{7.3\}$. So

$$\mathbb{P}(A) = \frac{\text{length}(A)}{\text{length}(\Omega)} = \frac{0}{12} = 0.$$

Some Questions to Answer

- How to define a Probability on the Finite Sample Space?

Some Questions to Answer

- How to define a Probability on the Finite Sample Space?
- How to calculate the Probability of an Event in the Equiprobable outcomes (Finite Sample Space) case?

Some Questions to Answer

- How to define a Probability on the Finite Sample Space?
- How to calculate the Probability of an Event in the Equiprobable outcomes (Finite Sample Space) case?
- What is the definition of the Geometric Probability Model?