

# AUA CS 108, Statistics, Fall 2019

## Lecture 38

Michael Poghosyan

YSU, AUA

[michael@ysu.am](mailto:michael@ysu.am), [mpoghosyan@aua.am](mailto:mpoghosyan@aua.am)

20 Nov 2019

# Contents

- ▶ HypoTesting and CIs
- ▶  $t$ -Test

## Last Lecture ReCap

- ▶ How many Errors we can do when Testing a Hypo?

## Last Lecture ReCap

- ▶ How many Errors we can do when Testing a Hypo?

Correct Answer:  $0, 1, 2, 3, 4, \dots, +\infty$ . Depends on how well we understand HypoTesting 😊

# Last Lecture ReCap

- ▶ How many Errors we can do when Testing a Hypo?

Correct Answer:  $0, 1, 2, 3, 4, \dots, +\infty$ . Depends on how well we understand HypoTesting 😊

- ▶ What is the Significance of the Test?

# Last Lecture ReCap

- ▶ How many Errors we can do when Testing a Hypo?

Correct Answer:  $0, 1, 2, 3, 4, \dots, +\infty$ . Depends on how well we understand HypoTesting 😊

- ▶ What is the Significance of the Test?
- ▶ What is the Power of the Test?

# Last Lecture ReCap

- ▶ How many Errors we can do when Testing a Hypo?

Correct Answer:  $0, 1, 2, 3, 4, \dots, +\infty$ . Depends on how well we understand HypoTesting 😊

- ▶ What is the Significance of the Test?
- ▶ What is the Power of the Test?
- ▶ Describe the Z-Test.

## Z-Test, relation to the Normal $\mu$ CI

Generally, it can be proven that there is a duality/association between Hypothesis Testing and a CI construction. Let me give the relation for our case, for Normal  $\mathcal{N}(\mu, \sigma^2)$  Model, when  $\sigma$  is known.



## Z-Test, relation to the Normal $\mu$ CI

Generally, it can be proven that there is a duality/association between Hypothesis Testing and a CI construction. Let me give the relation for our case, for Normal  $\mathcal{N}(\mu, \sigma^2)$  Model, when  $\sigma$  is known. Recall that we have obtained the following  $1 - \alpha$ -level CI for  $\mu$ :

$$\left( \bar{X} - z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} ; \bar{X} + z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \right)$$

## Z-Test, relation to the Normal $\mu$ CI

Generally, it can be proven that there is a duality/association between Hypothesis Testing and a CI construction. Let me give the relation for our case, for Normal  $\mathcal{N}(\mu, \sigma^2)$  Model, when  $\sigma$  is known. Recall that we have obtained the following  $1 - \alpha$ -level CI for  $\mu$ :

$$\left( \bar{X} - z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} ; \bar{X} + z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \right)$$

Now, consider the following two-tailed Test:

$$\mathcal{H}_0 : \mu = \mu_0 \quad \text{vs} \quad \mathcal{H}_1 : \mu \neq \mu_0$$

## Z-Test, relation to the Normal $\mu$ CI

Generally, it can be proven that there is a duality/association between Hypothesis Testing and a CI construction. Let me give the relation for our case, for Normal  $\mathcal{N}(\mu, \sigma^2)$  Model, when  $\sigma$  is known. Recall that we have obtained the following  $1 - \alpha$ -level CI for  $\mu$ :

$$\left( \bar{X} - z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} ; \bar{X} + z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \right)$$

Now, consider the following two-tailed Test:

$$\mathcal{H}_0 : \mu = \mu_0 \quad \text{vs} \quad \mathcal{H}_1 : \mu \neq \mu_0$$

Our Test procedure was:

► calculate  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$ .

## Z-Test, relation to the Normal $\mu$ CI

Generally, it can be proven that there is a duality/association between Hypothesis Testing and a CI construction. Let me give the relation for our case, for Normal  $\mathcal{N}(\mu, \sigma^2)$  Model, when  $\sigma$  is known. Recall that we have obtained the following  $1 - \alpha$ -level CI for  $\mu$ :

$$\left( \bar{X} - z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} ; \bar{X} + z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \right)$$

Now, consider the following two-tailed Test:

$$\mathcal{H}_0 : \mu = \mu_0 \quad \text{vs} \quad \mathcal{H}_1 : \mu \neq \mu_0$$

Our Test procedure was:

- ▶ calculate  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$ .
- ▶ Reject Null if  $|Z| > z_{1-\alpha/2}$ , otherwise, Do Not Reject Null.

## Z-Test, relation to the Normal $\mu$ CI

Generally, it can be proven that there is a duality/association between Hypothesis Testing and a CI construction. Let me give the relation for our case, for Normal  $\mathcal{N}(\mu, \sigma^2)$  Model, when  $\sigma$  is known. Recall that we have obtained the following  $1 - \alpha$ -level CI for  $\mu$ :

$$\left( \bar{X} - z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} ; \bar{X} + z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \right)$$

Now, consider the following two-tailed Test:

$$\mathcal{H}_0 : \mu = \mu_0 \quad \text{vs} \quad \mathcal{H}_1 : \mu \neq \mu_0$$

Our Test procedure was:

- ▶ calculate  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$ .
- ▶ Reject Null if  $|Z| > z_{1-\alpha/2}$ , otherwise, Do Not Reject Null.

This is equivalent to: Do Not Reject, if

$$-z_{1-\alpha/2} \leq \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \leq z_{1-\alpha/2},$$

## Z-Test, relation to the Normal $\mu$ CI

i.e., if

$$\mu_0 \in \left[ \bar{X} - z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} ; \bar{X} + z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \right].$$

---

<sup>1</sup>Well, we have obtained Closed CI, but that is OK.

## Z-Test, relation to the Normal $\mu$ CI

i.e., if

$$\mu_0 \in \left[ \bar{X} - z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} ; \bar{X} + z_{1-\alpha/2} \cdot \frac{\sigma}{\sqrt{n}} \right].$$

Hence, the relation: we Reject  $\mathcal{H}_0$ , if  $\mu_0$  is not in the CI, and otherwise, we Fail to Reject<sup>1</sup>.

---

<sup>1</sup>Well, we have obtained Closed CI, but that is OK.

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is known**, the Parameter (our unknown) is  $\mu$ ;



## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is known**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is known**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is known**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z =$

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is known**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is known**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim$

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is known, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim \mathcal{N}(0, 1)$ ;

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is known, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim \mathcal{N}(0, 1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is known, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim \mathcal{N}(0, 1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ Z  > z_{1-\frac{\alpha}{2}}$



## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is known, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim \mathcal{N}(0, 1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ Z  > z_{1-\frac{\alpha}{2}}$
$\mu > \mu_0$	

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is known, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim \mathcal{N}(0, 1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ Z  > z_{1-\frac{\alpha}{2}}$
$\mu > \mu_0$	$Z > z_{1-\alpha}$

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is known, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim \mathcal{N}(0, 1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ Z  > z_{1-\frac{\alpha}{2}}$
$\mu > \mu_0$	$Z > z_{1-\alpha}$
$\mu < \mu_0$	

## Z-Test, Complete Version

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is known, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim \mathcal{N}(0, 1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ Z  > z_{1-\frac{\alpha}{2}}$
$\mu > \mu_0$	$Z > z_{1-\alpha}$
$\mu < \mu_0$	$Z < z_{\alpha}$

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is unknown**, the Parameter (our unknown) is  $\mu$ ;

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**



## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t =$

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ ,

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  **is unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 =$

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is **unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$ .

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is **unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim$

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is **unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n-1)$ ;

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is **unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n-1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is **unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n-1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ t  > t_{n-1, 1-\frac{\alpha}{2}}$



## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is **unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n-1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ t  > t_{n-1, 1-\frac{\alpha}{2}}$
$\mu > \mu_0$	

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is **unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n-1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ t  > t_{n-1, 1-\frac{\alpha}{2}}$
$\mu > \mu_0$	$t > t_{n-1, 1-\alpha}$

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is **unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n-1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ t  > t_{n-1, 1-\frac{\alpha}{2}}$
$\mu > \mu_0$	$t > t_{n-1, 1-\alpha}$
$\mu < \mu_0$	

## $t$ -Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$ ,  $\sigma$  is **unknown**, the Parameter (our unknown) is  $\mu$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$ , where  $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n-1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu \neq \mu_0$	$ t  > t_{n-1, 1-\frac{\alpha}{2}}$
$\mu > \mu_0$	$t > t_{n-1, 1-\alpha}$
$\mu < \mu_0$	$t < t_{n-1, \alpha}$

## t-test Example

**Example:** Again, I have generated in **R** a Sample of Size 20 from  $\mathcal{N}(3.12, 2^2)$  and made some rounding:

```
set.seed(20112019)
s.size <- 20; sigma <- 2
obs <- rnorm(s.size, mean = 3.12, sd = sigma)
obs <- round(obs, digits = 2); obs
```

```
## [1] 1.80 5.60 1.10 3.20 4.91 5.15 1.76 2.47 0.
## [13] 3.98 4.79 1.98 4.50 3.52 4.13 -0.08 3.87
```

## t-test Example

**Example:** Again, I have generated in **R** a Sample of Size 20 from  $\mathcal{N}(3.12, 2^2)$  and made some rounding:

```
set.seed(20112019)
s.size <- 20; sigma <- 2
obs <- rnorm(s.size, mean = 3.12, sd = sigma)
obs <- round(obs, digits = 2); obs
```

```
## [1] 1.80 5.60 1.10 3.20 4.91 5.15 1.76 2.47 0.
## [13] 3.98 4.79 1.98 4.50 3.52 4.13 -0.08 3.87
```

Now, let us forget about the fact that the actual value of  $\mu$  is 3.12 and that  $\sigma = 2$ , and do some Testing, just assuming that our Observation is coming from a Normal Distribution.

## t-test Example

**Example:** Again, I have generated in **R** a Sample of Size 20 from  $\mathcal{N}(3.12, 2^2)$  and made some rounding:

```
set.seed(20112019)
s.size <- 20; sigma <- 2
obs <- rnorm(s.size, mean = 3.12, sd = sigma)
obs <- round(obs, digits = 2); obs
```

```
## [1] 1.80 5.60 1.10 3.20 4.91 5.15 1.76 2.47 0.
## [13] 3.98 4.79 1.98 4.50 3.52 4.13 -0.08 3.87
```

Now, let us forget about the fact that the actual value of  $\mu$  is 3.12 and that  $\sigma = 2$ , and do some Testing, just assuming that our Observation is coming from a Normal Distribution. Say, let us Test, at the 5% Significance Level, the following Hypothesis:

$$\mathcal{H}_0 : \mu = 4 \quad \text{vs} \quad \mathcal{H}_1 : \mu \neq 4.$$

## Example, Cont'd

First, we calculate  $t$ -statistic:

```
mu0 <- 4  
t <- (mean(obs) - mu0)/(sd(obs)/sqrt(s.size)); t  
  
## [1] -1.795358
```



## Example, Cont'd

First, we calculate  $t$ -statistic:

```
mu0 <- 4  
t <- (mean(obs) - mu0)/(sd(obs)/sqrt(s.size)); t  
  
## [1] -1.795358
```

Now, we calculate the critical value, the quantile  $t_{n-1, 1-\alpha/2}$ :

```
a <- 0.05  
c <- qt(1-a/2, df = s.size-1); c  
  
## [1] 2.093024
```

## Example, Cont'd

First, we calculate  $t$ -statistic:

```
mu0 <- 4  
t <- (mean(obs) - mu0)/(sd(obs)/sqrt(s.size)); t
```

```
## [1] -1.795358
```

Now, we calculate the critical value, the quantile  $t_{n-1,1-\alpha/2}$ :

```
a <- 0.05  
c <- qt(1-a/2, df = s.size-1); c
```

```
## [1] 2.093024
```

Finally, we check if  $t$  is in RR, i.e., if  $|t| > t_{n-1,1-\alpha/2}$ :

```
abs(t) > c
```

```
## [1] FALSE
```

## Example, Cont'd

First, we calculate  $t$ -statistic:

```
mu0 <- 4  
t <- (mean(obs) - mu0)/(sd(obs)/sqrt(s.size)); t
```

```
## [1] -1.795358
```

Now, we calculate the critical value, the quantile  $t_{n-1,1-\alpha/2}$ :

```
a <- 0.05  
c <- qt(1-a/2, df = s.size-1); c
```

```
## [1] 2.093024
```

Finally, we check if  $t$  is in RR, i.e., if  $|t| > t_{n-1,1-\alpha/2}$ :

```
abs(t) > c
```

```
## [1] FALSE
```

So the decision is:

## Example, Cont'd

First, we calculate  $t$ -statistic:

```
mu0 <- 4  
t <- (mean(obs) - mu0)/(sd(obs)/sqrt(s.size)); t  
  
## [1] -1.795358
```

Now, we calculate the critical value, the quantile  $t_{n-1,1-\alpha/2}$ :

```
a <- 0.05  
c <- qt(1-a/2, df = s.size-1); c  
  
## [1] 2.093024
```

Finally, we check if  $t$  is in RR, i.e., if  $|t| > t_{n-1,1-\alpha/2}$ :

```
abs(t) > c
```

```
## [1] FALSE
```

So the decision is: **Fail to Reject**  $\mathcal{H}_0$  at 5% level.

## Example, Cont'd

Now, the same, but with an **R** built-in function `t.test`:

```
t.test(obs, mu = mu0, conf.level = 0.95)
```

```
##  
##  One Sample t-test  
##  
## data:  obs  
## t = -1.7954, df = 19, p-value = 0.08852  
## alternative hypothesis: true mean is not equal to 4  
## 95 percent confidence interval:  
##  2.524009 4.112991  
## sample estimates:  
## mean of x  
##      3.3185
```

## Example, Cont'd

Now, for the same Data, let us Test, at the 10% Significance Level, the following Hypothesis:

$$\mathcal{H}_0 : \mu = 3 \quad \text{vs} \quad \mathcal{H}_1 : \mu > 3.$$

## Example, Cont'd

Now, for the same Data, let us Test, at the 10% Significance Level, the following Hypothesis:

$$\mathcal{H}_0 : \mu = 3 \quad \text{vs} \quad \mathcal{H}_1 : \mu > 3.$$

```
t.test(obs, mu=3, alternative="greater", conf.level=0.9)
```

```
##  
## One Sample t-test  
##  
## data: obs  
## t = 0.83906, df = 19, p-value = 0.2059  
## alternative hypothesis: true mean is greater than 3  
## 90 percent confidence interval:  
## 2.814508 Inf  
## sample estimates:  
## mean of x  
## 3.3185
```