

# AUA CS108, Statistics, Fall 2020

## Lecture 39

Michael Poghosyan

25 Nov 2020

# Contents

- ▶ Confidence Intervals
- ▶ Hypothesis Testing

Asymptotic CIs

# Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want to Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ .

# Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want to Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ . Here we assume that the Variance  $\sigma^2 = \text{Var}(X_k)$  is finite, but unknown.

## Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want to Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ . Here we assume that the Variance  $\sigma^2 = \text{Var}(X_k)$  is finite, but unknown.

So our Problem is to construct an Asymptotic CI of level  $1 - \alpha$  for  $\mu$ .

# Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want to Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ . Here we assume that the Variance  $\sigma^2 = \text{Var}(X_k)$  is finite, but unknown.

So our Problem is to construct an Asymptotic CI of level  $1 - \alpha$  for  $\mu$ .

**Answer:** The Random Interval (or, rather, the sequence of Intervals)

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

is a  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ .

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$



## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:

$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:

$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

This is because

- ▶ the same form of CI was obtained for the Normal Model, when  $\sigma^2$  was unknown;

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:

$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

This is because

- ▶ the same form of CI was obtained for the Normal Model, when  $\sigma^2$  was unknown;
- ▶ this interval is a little bit larger than the previous one, so it is also an AsympCI for  $\mu$  of level  $1 - \alpha$ ;

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:

$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

This is because

- ▶ the same form of CI was obtained for the Normal Model, when  $\sigma^2$  was unknown;
- ▶ this interval is a little bit larger than the previous one, so it is also an AsympCI for  $\mu$  of level  $1 - \alpha$ ;
- ▶ when  $n \geq 30$ , these two almost coincide;

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:


$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

This is because

- ▶ the same form of CI was obtained for the Normal Model, when  $\sigma^2$  was unknown;
- ▶ this interval is a little bit larger than the previous one, so it is also an AsympCI for  $\mu$  of level  $1 - \alpha$ ;
- ▶ when  $n \geq 30$ , these two almost coincide;
- ▶ although in the theory these intervals work for large  $n$ , but, in practice, the latter one works also for small  $n$


# Example

Asymptotic CI for the Mean with  $z$ ,  $n = 50$




# Example

Asymptotic CI for the Mean with  $z$ ,  $n = 20$



# Example





## Example

**Example:** There was an ancient civilization called Etruscan Civilization, living in the area corresponding to Italy, around 8th century BC.

## Example

**Example:** There was an ancient civilization called Etruscan Civilization, living in the area corresponding to Italy, around 8th century BC. Scientists have a data about 84 skull sizes (widths) of adult Etruscans, giving in average 143.8mm, with a Sample Standard Deviation 5.97mm.

## Example

**Example:** There was an ancient civilization called Etruscan Civilization, living in the area corresponding to Italy, around 8th century BC. Scientists have a data about 84 skull sizes (widths) of adult Etruscans, giving in average 143.8mm, with a Sample Standard Deviation 5.97mm.

Scientists believe Etruscans were Italians.

## Example

**Example:** There was an ancient civilization called Etruscan Civilization, living in the area corresponding to Italy, around 8th century BC. Scientists have a data about 84 skull sizes (widths) of adult Etruscans, giving in average 143.8mm, with a Sample Standard Deviation 5.97mm.

Scientists believe Etruscans were Italians. They believe that the skull size is not changing much through time.

## Example

**Example:** There was an ancient civilization called Etruscan Civilization, living in the area corresponding to Italy, around 8th century BC. Scientists have a data about 84 skull sizes (widths) of adult Etruscans, giving in average 143.8mm, with a Sample Standard Deviation 5.97mm.

Scientists believe Etruscans were Italians. They believe that the skull size is not changing much through time. Nowadays, adult Italians skull size is in average 132.4mm.

## Example

**Example:** There was an ancient civilization called Etruscan Civilization, living in the area corresponding to Italy, around 8th century BC. Scientists have a data about 84 skull sizes (widths) of adult Etruscans, giving in average 143.8mm, with a Sample Standard Deviation 5.97mm.

Scientists believe Etruscans were Italians. They believe that the skull size is not changing much through time. Nowadays, adult Italians skull size is in average 132.4mm.

We want to construct a 95% Asymptotic CI to see if it is supporting scientists hypothesis.

## Example

**Example:** There was an ancient civilization called Etruscan Civilization, living in the area corresponding to Italy, around 8th century BC. Scientists have a data about 84 skull sizes (widths) of adult Etruscans, giving in average 143.8mm, with a Sample Standard Deviation 5.97mm.

Scientists believe Etruscans were Italians. They believe that the skull size is not changing much through time. Nowadays, adult Italians skull size is in average 132.4mm.

We want to construct a 95% Asymptotic CI to see if it is supporting scientists hypothesis.

We model our problem like this: we assume the skull sizes of Italians are coming from some Distribution with some Mean  $\mu$  and Variance  $\sigma^2$ ,  $\sigma^2$  is unknown.

## Example, Cont'd

If we believe that Etruscans are Italians, then we have a Sample from that Distrib:

$$X_1, X_2, \dots, X_{84}.$$

where  $X_k$  is the skull size of  $k$ -th Etruscan person.



## Example, Cont'd

If we believe that Etruscans are Italians, then we have a Sample from that Distrib:

$$X_1, X_2, \dots, X_{84}.$$

where  $X_k$  is the skull size of  $k$ -th Etruscan person. We have that  $n = 84$ ,  $\bar{X} = 143.8$  and  $S = 5.97$ .

## Example, Cont'd

If we believe that Etruscans are Italians, then we have a Sample from that Distrib:

$$X_1, X_2, \dots, X_{84}.$$

where  $X_k$  is the skull size of  $k$ -th Etruscan person. We have that  $n = 84$ ,  $\bar{X} = 143.8$  and  $S = 5.97$ .

Then, we know how to construct an Asympto 95% CI for  $\mu$ :

$$\bar{X} \pm t_{n-1, 1-\alpha/2} \cdot \frac{S}{\sqrt{n}}$$

## Example, Cont'd

If we believe that Etruscans are Italians, then we have a Sample from that Distrib:

$$X_1, X_2, \dots, X_{84}.$$

where  $X_k$  is the skull size of  $k$ -th Etruscan person. We have that  $n = 84$ ,  $\bar{X} = 143.8$  and  $S = 5.97$ .

Then, we know how to construct an Asympto 95% CI for  $\mu$ :

$$\bar{X} \pm t_{n-1, 1-\alpha/2} \cdot \frac{S}{\sqrt{n}}$$

giving us

```
x.bar <- 143.8; s <- 5.97; n <- 84  
a <- 0.05; t <- qt(1-a/2, df = n-1)  
me <- t*s/sqrt(n)  
c(x.bar - me, x.bar +me)
```

```
## [1] 142.5044 145.0956
```

## Example, Cont'd

If we believe that Etruscans are Italians, then we have a Sample from that Distrib:

$$X_1, X_2, \dots, X_{84}.$$

where  $X_k$  is the skull size of  $k$ -th Etruscan person. We have that  $n = 84$ ,  $\bar{X} = 143.8$  and  $S = 5.97$ .

Then, we know how to construct an Asympto 95% CI for  $\mu$ :

$$\bar{X} \pm t_{n-1, 1-\alpha/2} \cdot \frac{S}{\sqrt{n}}$$

giving us

```
x.bar <- 143.8; s <- 5.97; n <- 84  
a <- 0.05; t <- qt(1-a/2, df = n-1)  
me <- t*s/sqrt(n)  
c(x.bar - me, x.bar +me)
```

```
## [1] 142.5044 145.0956
```

We know the true value of  $\mu$ , it is  $\mu = 132.4$ .

## Example, Cont'd

If we believe that Etruscans are Italians, then we have a Sample from that Distrib:

$$X_1, X_2, \dots, X_{84}.$$

where  $X_k$  is the skull size of  $k$ -th Etruscan person. We have that  $n = 84$ ,  $\bar{X} = 143.8$  and  $S = 5.97$ .

Then, we know how to construct an Asympto 95% CI for  $\mu$ :

$$\bar{X} \pm t_{n-1, 1-\alpha/2} \cdot \frac{S}{\sqrt{n}}$$

giving us

```
x.bar <- 143.8; s <- 5.97; n <- 84  
a <- 0.05; t <- qt(1-a/2, df = n-1)  
me <- t*s/sqrt(n)  
c(x.bar - me, x.bar +me)
```

```
## [1] 142.5044 145.0956
```

We know the true value of  $\mu$ , it is  $\mu = 132.4$ . So we can deduce that ...

## Example, Cont'd

If we believe that Etruscans are Italians, then we have a Sample from that Distrib:

$$X_1, X_2, \dots, X_{84}.$$

where  $X_k$  is the skull size of  $k$ -th Etruscan person. We have that  $n = 84$ ,  $\bar{X} = 143.8$  and  $S = 5.97$ .

Then, we know how to construct an Asympto 95% CI for  $\mu$ :

$$\bar{X} \pm t_{n-1, 1-\alpha/2} \cdot \frac{S}{\sqrt{n}}$$

giving us

```
x.bar <- 143.8; s <- 5.97; n <- 84  
a <- 0.05; t <- qt(1-a/2, df = n-1)  
me <- t*s/sqrt(n)  
c(x.bar - me, x.bar +me)
```

```
## [1] 142.5044 145.0956
```

We know the true value of  $\mu$ , it is  $\mu = 132.4$ . So we can deduce that ... they were not Italians,

## Example, Cont'd

If we believe that Etruscans are Italians, then we have a Sample from that Distrib:

$$X_1, X_2, \dots, X_{84}.$$

where  $X_k$  is the skull size of  $k$ -th Etruscan person. We have that  $n = 84$ ,  $\bar{X} = 143.8$  and  $S = 5.97$ .

Then, we know how to construct an Asympto 95% CI for  $\mu$ :

$$\bar{X} \pm t_{n-1, 1-\alpha/2} \cdot \frac{S}{\sqrt{n}}$$

giving us

```
x.bar <- 143.8; s <- 5.97; n <- 84  
a <- 0.05; t <- qt(1-a/2, df = n-1)  
me <- t*s/sqrt(n)  
c(x.bar - me, x.bar +me)
```

```
## [1] 142.5044 145.0956
```

We know the true value of  $\mu$ , it is  $\mu = 132.4$ . So we can deduce that ... they were not Italians, or were looong skul Italians 😊

## Asymptotic CI for the general case, using MLE

Using the Asymptotic Normality of the MLE, we can obtain a  $(1 - \alpha)$ -level Asymptotic CI for the general case.



## Asymptotic CI for the general case, using MLE

Using the Asymptotic Normality of the MLE, we can obtain a  $(1 - \alpha)$ -level Asymptotic CI for the general case.

Assume we have a (regular) Parametric Model  $\mathcal{F}_\theta$ , and a Random Sample from it

$$X_1, X_2, \dots, X_n, \dots \sim \mathcal{F}_\theta.$$

## Asymptotic CI for the general case, using MLE

Using the Asymptotic Normality of the MLE, we can obtain a  $(1 - \alpha)$ -level Asymptotic CI for the general case.

Assume we have a (regular) Parametric Model  $\mathcal{F}_\theta$ , and a Random Sample from it

$$X_1, X_2, \dots, X_n, \dots \sim \mathcal{F}_\theta.$$

We want to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $\theta$ .

## Asymptotic CI for the general case, using MLE

Using the Asymptotic Normality of the MLE, we can obtain a  $(1 - \alpha)$ -level Asymptotic CI for the general case.

Assume we have a (regular) Parametric Model  $\mathcal{F}_\theta$ , and a Random Sample from it

$$X_1, X_2, \dots, X_n, \dots \sim \mathcal{F}_\theta.$$

We want to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $\theta$ .

Let  $\hat{\theta}_n^{MLE}$  be the ML Estimator for  $\theta$ .

## Asymptotic CI for the general case, using MLE

Using the Asymptotic Normality of the MLE, we can obtain a  $(1 - \alpha)$ -level Asymptotic CI for the general case.

Assume we have a (regular) Parametric Model  $\mathcal{F}_\theta$ , and a Random Sample from it

$$X_1, X_2, \dots, X_n, \dots \sim \mathcal{F}_\theta.$$

We want to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $\theta$ .

Let  $\hat{\theta}_n^{MLE}$  be the ML Estimator for  $\theta$ . Then the following is giving us a  $(1 - \alpha)$ -level Asymptotic CI for  $\theta$ :

$$\left( \hat{\theta}_n^{MLE} - z_{1-\alpha/2} \cdot \sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}}; \quad \hat{\theta}_n^{MLE} + z_{1-\alpha/2} \cdot \sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}} \right).$$

## Asymptotic CI for the Bernoulli Model

We have already constructed a CI for  $p$  in the *Bernoulli*( $p$ ) Model. Here we want to construct an Asymptotic CI for  $p$ , which usually gives smaller length interval.

## Asymptotic CI for the Bernoulli Model

We have already constructed a CI for  $p$  in the *Bernoulli*( $p$ ) Model. Here we want to construct an Asymptotic CI for  $p$ , which usually gives smaller length interval.

The Problem is the following: we have a Random Sample

$$X_1, X_2, \dots, X_n, \dots \sim \text{Bernoulli}(p),$$

and  $\alpha \in (0, 1)$ .

## Asymptotic CI for the Bernoulli Model

We have already constructed a CI for  $p$  in the *Bernoulli*( $p$ ) Model. Here we want to construct an Asymptotic CI for  $p$ , which usually gives smaller length interval.

The Problem is the following: we have a Random Sample

$$X_1, X_2, \dots, X_n, \dots \sim \text{Bernoulli}(p),$$

and  $\alpha \in (0, 1)$ . Our aim is to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $p$ .

## Asymptotic CI for the Bernoulli Model

We have already constructed a CI for  $p$  in the *Bernoulli*( $p$ ) Model. Here we want to construct an Asymptotic CI for  $p$ , which usually gives smaller length interval.

The Problem is the following: we have a Random Sample

$$X_1, X_2, \dots, X_n, \dots \sim \text{Bernoulli}(p),$$

and  $\alpha \in (0, 1)$ . Our aim is to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $p$ .

We know that  $\hat{p}_n^{MLE} =$



## Asymptotic CI for the Bernoulli Model

We have already constructed a CI for  $p$  in the *Bernoulli*( $p$ ) Model. Here we want to construct an Asymptotic CI for  $p$ , which usually gives smaller length interval.

The Problem is the following: we have a Random Sample

$$X_1, X_2, \dots, X_n, \dots \sim \text{Bernoulli}(p),$$

and  $\alpha \in (0, 1)$ . Our aim is to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $p$ .

We know that  $\hat{p}_n^{MLE} = \bar{X}_n$ . Also, we can calculate the Fisher Information for the Bernoulli case:  $\mathcal{I}(p) =$

## Asymptotic CI for the Bernoulli Model

We have already constructed a CI for  $p$  in the *Bernoulli*( $p$ ) Model. Here we want to construct an Asymptotic CI for  $p$ , which usually gives smaller length interval.

The Problem is the following: we have a Random Sample

$$X_1, X_2, \dots, X_n, \dots \sim \text{Bernoulli}(p),$$

and  $\alpha \in (0, 1)$ . Our aim is to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $p$ .

We know that  $\hat{p}_n^{MLE} = \bar{X}_n$ . Also, we can calculate the Fisher Information for the Bernoulli case:  $\mathcal{I}(p) = \frac{1}{p(1-p)}$ .

## Asymptotic CI for the Bernoulli Model

By plugging the values for our case, we'll obtain the following Asymptotic CI of level  $(1 - \alpha)$  for  $p$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \sqrt{\frac{\bar{X}_n(1 - \bar{X}_n)}{n}}; \bar{X}_n + z_{1-\alpha/2} \cdot \sqrt{\frac{\bar{X}_n(1 - \bar{X}_n)}{n}} \right).$$

## Example, Asymptotic CI for Poisson Model

**Example:** We consider the Poisson Model  $Pois(\lambda)$ , and a Random Sample from it:

$$X_1, X_2, \dots, X_n, \dots \sim Pois(\lambda).$$

## Example, Asymptotic CI for Poisson Model

**Example:** We consider the Poisson Model  $Pois(\lambda)$ , and a Random Sample from it:

$$X_1, X_2, \dots, X_n, \dots \sim Pois(\lambda).$$

Taking an  $\alpha \in (0, 1)$ , we want to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $\lambda$ .

## Example, Asymptotic CI for Poisson Model

**Example:** We consider the Poisson Model  $Pois(\lambda)$ , and a Random Sample from it:

$$X_1, X_2, \dots, X_n, \dots \sim Pois(\lambda).$$

Taking an  $\alpha \in (0, 1)$ , we want to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $\lambda$ .

We know that  $\hat{\lambda}_n^{MLE} =$

## Example, Asymptotic CI for Poisson Model

**Example:** We consider the Poisson Model  $Pois(\lambda)$ , and a Random Sample from it:

$$X_1, X_2, \dots, X_n, \dots \sim Pois(\lambda).$$

Taking an  $\alpha \in (0, 1)$ , we want to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $\lambda$ .

We know that  $\hat{\lambda}_n^{MLE} = \bar{X}_n$ , and also,  $\mathcal{I}(\lambda) =$

## Example, Asymptotic CI for Poisson Model

**Example:** We consider the Poisson Model  $Pois(\lambda)$ , and a Random Sample from it:

$$X_1, X_2, \dots, X_n, \dots \sim Pois(\lambda).$$

Taking an  $\alpha \in (0, 1)$ , we want to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $\lambda$ .

We know that  $\hat{\lambda}_n^{MLE} = \bar{X}_n$ , and also,  $\mathcal{I}(\lambda) = \frac{1}{\lambda}$ .



## Example, Asymptotic CI for Poisson Model

**Example:** We consider the Poisson Model  $Pois(\lambda)$ , and a Random Sample from it:

$$X_1, X_2, \dots, X_n, \dots \sim Pois(\lambda).$$


Taking an  $\alpha \in (0, 1)$ , we want to construct a  $(1 - \alpha)$ -level Asymptotic CI for  $\lambda$ .

We know that  $\hat{\lambda}_n^{MLE} = \bar{X}_n$ , and also,  $\mathcal{I}(\lambda) = \frac{1}{\lambda}$ . Then, using the formula above, we'll obtain the following Asymptotic CI of level  $(1 - \alpha)$  for  $\lambda$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \sqrt{\frac{\bar{X}_n}{n}}; \quad \bar{X}_n + z_{1-\alpha/2} \cdot \sqrt{\frac{\bar{X}_n}{n}} \right).$$

# Example, in R

Asymptotic CI for the Pois lambda, n = 50



## Example, in R, Code

```
lambda <- 3.2
conf.level <- 0.95; a = 1 - conf.level
sample.size <- 15; no.of.intervals <- 100
z <- qnorm(1-a/2)

plot.new()
plot.window(xlim=c(0,no.of.intervals),ylim=c(lambda-3,lambda+3))
axis(1); axis(2)
title("Asymptotic CI for the Pois lambda, n = 50")
for(i in 1:no.of.intervals){
  x <- rpois(sample.size, lambda = lambda)
  ME <- z*sqrt(mean(x)/sample.size) #Margin of Error
  lo <- mean(x) - ME; up <- mean(x) + ME
  if(lo > lambda || up < lambda){
    segments(c(i), c(lo), c(i), c(up), col = "red")
  }
  else{
    segments(c(i), c(lo), c(i), c(up))
  }
}
abline(h = lambda, lwd = 2, col = "blue")
```

# Hypothesis Testing

# Hypothesis Testing, Intro

Up to this point, we have considered the Point Estimation and CI construction Problems.

# Hypothesis Testing, Intro

Up to this point, we have considered the Point Estimation and CI construction Problems. Here we will talk about another, very important Problem in Statistics, Hypothesis Testing.

# Hypothesis Testing, Intro

Up to this point, we have considered the Point Estimation and CI construction Problems. Here we will talk about another, very important Problem in Statistics, Hypothesis Testing.

As always, we assume we have a Dataset coming as a realization of a Random Sample from some unknown Parametric Distribution  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta.$$

In this case we want to Test a Hypothesis about  $\theta$ : say, see whether  $\theta = \theta_0$ , a given number, or not.

## Example

**Example:** We have a coin, and we want to see if it is fair or not.



## Example

**Example:** We have a coin, and we want to see if it is fair or not.  
We formulate a Hypothesis: coin is fair.

## Example

**Example:** We have a coin, and we want to see if it is fair or not. We formulate a Hypothesis: coin is fair. To test the hypothesis, we toss that coin 100 times and obtain 56 Heads. Is this supporting our Hypothesis or not?

## Example

**Example:** We have a coin, and we want to see if it is fair or not. We formulate a Hypothesis: coin is fair. To test the hypothesis, we toss that coin 100 times and obtain 56 Heads. Is this supporting our Hypothesis or not?

**The Idea** The idea is the following: even if our coin was fair, the Probability of Heads  $p = 0.5$ , it is possible to have some deviation from the expected number of Heads, 50 (in 100 tosses).

## Example

**Example:** We have a coin, and we want to see if it is fair or not. We formulate a Hypothesis: coin is fair. To test the hypothesis, we toss that coin 100 times and obtain 56 Heads. Is this supporting our Hypothesis or not?

**The Idea** The idea is the following: even if our coin was fair, the Probability of Heads  $p = 0.5$ , it is possible to have some deviation from the expected number of Heads, 50 (in 100 tosses). The question is: how extreme, how unbelievable is that deviation.

## Example

**Example:** We have a coin, and we want to see if it is fair or not. We formulate a Hypothesis: coin is fair. To test the hypothesis, we toss that coin 100 times and obtain 56 Heads. Is this supporting our Hypothesis or not?

**The Idea** The idea is the following: even if our coin was fair, the Probability of Heads  $p = 0.5$ , it is possible to have some deviation from the expected number of Heads, 50 (in 100 tosses). The question is: how extreme, how unbelievable is that deviation. If the deviation from the expected is large, we will not believe that the Hypothesis is True. Otherwise, we will keep believing that it is True.

# Hypothesis Testing: Problem Setting and Formalization

So we have a Parametric Model  $\mathcal{F}_\theta$ , with  $\theta \in \Theta$ .

## Hypothesis Testing: Problem Setting and Formalization

So we have a Parametric Model  $\mathcal{F}_\theta$ , with  $\theta \in \Theta$ . We consider a point  $\theta_0 \in \Theta$ , and consider the following **Null Hypothesis**:

$$\mathcal{H}_0 : \theta = \theta_0.$$

# Hypothesis Testing: Problem Setting and Formalization

So we have a Parametric Model  $\mathcal{F}_\theta$ , with  $\theta \in \Theta$ . We consider a point  $\theta_0 \in \Theta$ , and consider the following **Null Hypothesis**:

$$\mathcal{H}_0 : \theta = \theta_0.$$

We consider three type of **Alternative Hypotheses**:

- ▶  $\mathcal{H}_1 : \theta \neq \theta_0$  (Two-Tailed, Two-Sided Test)



# Hypothesis Testing: Problem Setting and Formalization

So we have a Parametric Model  $\mathcal{F}_\theta$ , with  $\theta \in \Theta$ . We consider a point  $\theta_0 \in \Theta$ , and consider the following **Null Hypothesis**:

$$\mathcal{H}_0 : \theta = \theta_0.$$

We consider three type of **Alternative Hypotheses**:

- ▶  $\mathcal{H}_1 : \theta \neq \theta_0$  (Two-Tailed, Two-Sided Test)
- ▶  $\mathcal{H}_1 : \theta > \theta_0$  (Upper-Tailed Test)

# Hypothesis Testing: Problem Setting and Formalization

So we have a Parametric Model  $\mathcal{F}_\theta$ , with  $\theta \in \Theta$ . We consider a point  $\theta_0 \in \Theta$ , and consider the following **Null Hypothesis**:

$$\mathcal{H}_0 : \theta = \theta_0.$$

We consider three type of **Alternative Hypotheses**:

- ▶  $\mathcal{H}_1 : \theta \neq \theta_0$  (Two-Tailed, Two-Sided Test)
- ▶  $\mathcal{H}_1 : \theta > \theta_0$  (Upper-Tailed Test)
- ▶  $\mathcal{H}_1 : \theta < \theta_0$  (Lower-Tailed Test)

# Hypothesis Testing: Problem Setting and Formalization

So we have a Parametric Model  $\mathcal{F}_\theta$ , with  $\theta \in \Theta$ . We consider a point  $\theta_0 \in \Theta$ , and consider the following **Null Hypothesis**:

$$\mathcal{H}_0 : \theta = \theta_0.$$

We consider three type of **Alternative Hypotheses**:

- ▶  $\mathcal{H}_1 : \theta \neq \theta_0$  (Two-Tailed, Two-Sided Test)
- ▶  $\mathcal{H}_1 : \theta > \theta_0$  (Upper-Tailed Test)
- ▶  $\mathcal{H}_1 : \theta < \theta_0$  (Lower-Tailed Test)

So our Hypothesis will be either

$$\mathcal{H}_0 : \theta = \theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta \neq \theta_0$$

# Hypothesis Testing: Problem Setting and Formalization

So we have a Parametric Model  $\mathcal{F}_\theta$ , with  $\theta \in \Theta$ . We consider a point  $\theta_0 \in \Theta$ , and consider the following **Null Hypothesis**:

$$\mathcal{H}_0 : \theta = \theta_0.$$

We consider three type of **Alternative Hypotheses**:

- ▶  $\mathcal{H}_1 : \theta \neq \theta_0$  (Two-Tailed, Two-Sided Test)
- ▶  $\mathcal{H}_1 : \theta > \theta_0$  (Upper-Tailed Test)
- ▶  $\mathcal{H}_1 : \theta < \theta_0$  (Lower-Tailed Test)

So our Hypothesis will be either

$$\mathcal{H}_0 : \theta = \theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta \neq \theta_0$$

or

$$\mathcal{H}_0 : \theta = \theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta > \theta_0$$

# Hypothesis Testing: Problem Setting and Formalization

So we have a Parametric Model  $\mathcal{F}_\theta$ , with  $\theta \in \Theta$ . We consider a point  $\theta_0 \in \Theta$ , and consider the following **Null Hypothesis**:

$$\mathcal{H}_0 : \theta = \theta_0.$$

We consider three type of **Alternative Hypotheses**:

- ▶  $\mathcal{H}_1 : \theta \neq \theta_0$  (Two-Tailed, Two-Sided Test)
- ▶  $\mathcal{H}_1 : \theta > \theta_0$  (Upper-Tailed Test)
- ▶  $\mathcal{H}_1 : \theta < \theta_0$  (Lower-Tailed Test)

So our Hypothesis will be either

$$\mathcal{H}_0 : \theta = \theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta \neq \theta_0$$

or

$$\mathcal{H}_0 : \theta = \theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta > \theta_0$$

or

$$\mathcal{H}_0 : \theta = \theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta < \theta_0.$$

# Hypothesis Testing: Problem Setting and Formalization

Next, we have a Random Sample from  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and using this Sample, we want to Test the Hypothesis, we want to see if we can **Reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$  or not**, i.e., we want to see if **we have enough evidence in our Data to Reject  $\mathcal{H}_0$** .

# Hypothesis Testing: Problem Setting and Formalization

Next, we have a Random Sample from  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and using this Sample, we want to Test the Hypothesis, we want to see if we can **Reject  $\mathcal{H}_0$  in favor of  $\mathcal{H}_1$  or not**, i.e., we want to see if **we have enough evidence in our Data to Reject  $\mathcal{H}_0$** .

**Example:** In the above example about the coin fairness, if  $p$  is the Probability of a Head, then our Hypotheses are:

$$\mathcal{H}_0 : p = \frac{1}{2} \quad \text{vs} \quad \mathcal{H}_1 : p \neq \frac{1}{2}.$$

## Example

**Example:** Assume that the Mean Stat grade for my courses is 71.



## Example

**Example:** Assume that the Mean Stat grade for my courses is 71. This year we are doing classes online, I am changing the hw, using **R** a lot.

## Example

**Example:** Assume that the Mean Stat grade for my courses is 71. This year we are doing classes online, I am changing the hw, using **R** a lot. And we want to see if this affects positively on the average grade or not.

## Example

**Example:** Assume that the Mean Stat grade for my courses is 71. This year we are doing classes online, I am changing the hw, using **R** a lot. And we want to see if this affects positively on the average grade or not. With high Probability, our average grade will not be exactly 71.

## Example

**Example:** Assume that the Mean Stat grade for my courses is 71. This year we are doing classes online, I am changing the hw, using **R** a lot. And we want to see if this affects positively on the average grade or not. With high Probability, our average grade will not be exactly 71. Say, it will be 73.5.

## Example

**Example:** Assume that the Mean Stat grade for my courses is 71. This year we are doing classes online, I am changing the hw, using **R** a lot. And we want to see if this affects positively on the average grade or not. With high Probability, our average grade will not be exactly 71. Say, it will be 73.5. Will this be a sign that my new approach is working, or that will be just because of randomness?

## Example

**Example:** Assume that the Mean Stat grade for my courses is 71. This year we are doing classes online, I am changing the hw, using **R** a lot. And we want to see if this affects positively on the average grade or not. With high Probability, our average grade will not be exactly 71. Say, it will be 73.5. Will this be a sign that my new approach is working, or that will be just because of randomness? So we want to see if the **difference is significant or not**.

## Example

**Example:** Assume that the Mean Stat grade for my courses is 71. This year we are doing classes online, I am changing the hw, using **R** a lot. And we want to see if this affects positively on the average grade or not. With high Probability, our average grade will not be exactly 71. Say, it will be 73.5. Will this be a sign that my new approach is working, or that will be just because of randomness? So we want to see if the **difference is significant or not**.

To state the Hypotheses, let us denote the Average Stat Grade by  $\mu$ .

## Example

**Example:** Assume that the Mean Stat grade for my courses is 71. This year we are doing classes online, I am changing the hw, using **R** a lot. And we want to see if this affects positively on the average grade or not. With high Probability, our average grade will not be exactly 71. Say, it will be 73.5. Will this be a sign that my new approach is working, or that will be just because of randomness? So we want to see if the **difference is significant or not**.

To state the Hypotheses, let us denote the Average Stat Grade by  $\mu$ . Then our Hypotheses will be:

$$\mathcal{H}_0 : \mu = 71 \quad \text{vs} \quad \mathcal{H}_1 : \mu > 71.$$



## Example

**Example:** Assume that the Mean Stat grade for my courses is 71. This year we are doing classes online, I am changing the hw, using **R** a lot. And we want to see if this affects positively on the average grade or not. With high Probability, our average grade will not be exactly 71. Say, it will be 73.5. Will this be a sign that my new approach is working, or that will be just because of randomness? So we want to see if the **difference is significant or not**.

To state the Hypotheses, let us denote the Average Stat Grade by  $\mu$ . Then our Hypotheses will be:

$$\mathcal{H}_0 : \mu = 71 \quad \text{vs} \quad \mathcal{H}_1 : \mu > 71.$$

So, when we will have our Data, we will see if we can **Reject**  $\mathcal{H}_0$ .

# Notes

**Note:** In Hypothesis Testing, our Strategy is, based on Data, to **try to Reject**  $\mathcal{H}_0$ .

## Notes

**Note:** In Hypothesis Testing, our Strategy is, based on Data, to **try to Reject**  $\mathcal{H}_0$ . If our Data contains enough evidence against  $\mathcal{H}_0$ , we are Rejecting it.

# Notes

**Note:** In Hypothesis Testing, our Strategy is, based on Data, to **try to Reject**  $\mathcal{H}_0$ . If our Data contains enough evidence against  $\mathcal{H}_0$ , we are Rejecting it. **Otherwise, we are Failing to Reject**  $\mathcal{H}_0$ , and we are not talking about we are accepting  $\mathcal{H}_1$ .

# Notes

**Note:** In Hypothesis Testing, our Strategy is, based on Data, to **try to Reject**  $\mathcal{H}_0$ . If our Data contains enough evidence against  $\mathcal{H}_0$ , we are Rejecting it. **Otherwise, we are Failing to Reject**  $\mathcal{H}_0$ , and we are not talking about we are accepting  $\mathcal{H}_1$ .

So the conclusion of the Hypothesis Testing need to be either:

**Reject  $\mathcal{H}_0$    or   Fail to Reject  $\mathcal{H}_0$ .**

## Notes

At this point, Statisticians always compare this with the process in the court:

## Notes

At this point, Statisticians always compare this with the process in the court: if the person is suspected, say, in murder, then, by default, judges assume that he is **not guilty** (anmeghutyan kankhavarkac ☺).

# Notes

At this point, Statisticians always compare this with the process in the court: if the person is suspected, say, in murder, then, by default, judges assume that he is **not guilty** (anmeghutyan kankhavarkac 😊). This is our Null Hypothesis:

$\mathcal{H}_0$  : He is not guilty    vs     $\mathcal{H}_1$  : He is guilty.



# Notes

At this point, Statisticians always compare this with the process in the court: if the person is suspected, say, in murder, then, by default, judges assume that he is **not guilty** (anmeghutyan kankhavarkac ☺). This is our Null Hypothesis:

$\mathcal{H}_0$  : He is not guilty    vs     $\mathcal{H}_1$  : He is guilty.

Now, based on the evidence, Data, the judges either will Reject  $\mathcal{H}_0$ , meaning that, upon the light of a lot of facts, they will decide that our person is guilty,

# Notes

At this point, Statisticians always compare this with the process in the court: if the person is suspected, say, in murder, then, by default, judges assume that he is **not guilty** (anmeghutyan kankhavarkac 😊). This is our Null Hypothesis:

$\mathcal{H}_0$  : He is not guilty    vs     $\mathcal{H}_1$  : He is guilty.

Now, based on the evidence, Data, the judges either will Reject  $\mathcal{H}_0$ , meaning that, upon the light of a lot of facts, they will decide that our person is guilty, or, if the Data will not be enough, the judges will Not Reject  $\mathcal{H}_0$ , so they will free our person.

# Notes

At this point, Statisticians always compare this with the process in the court: if the person is suspected, say, in murder, then, by default, judges assume that he is **not guilty** (anmeghutyan kankhavarkac 😊). This is our Null Hypothesis:

$\mathcal{H}_0$  : He is not guilty    vs     $\mathcal{H}_1$  : He is guilty.

Now, based on the evidence, Data, the judges either will Reject  $\mathcal{H}_0$ , meaning that, upon the light of a lot of facts, they will decide that our person is guilty, or, if the Data will not be enough, the judges will Not Reject  $\mathcal{H}_0$ , so they will free our person. **But this does not mean that the murder was not of our person's hands affair!**

# Notes

At this point, Statisticians always compare this with the process in the court: if the person is suspected, say, in murder, then, by default, judges assume that he is **not guilty** (anmeghutyan kankhavarkac 😊). This is our Null Hypothesis:

$\mathcal{H}_0$  : He is not guilty    vs     $\mathcal{H}_1$  : He is guilty.

Now, based on the evidence, Data, the judges either will Reject  $\mathcal{H}_0$ , meaning that, upon the light of a lot of facts, they will decide that our person is guilty, or, if the Data will not be enough, the judges will Not Reject  $\mathcal{H}_0$ , so they will free our person. **But this does not mean that the murder was not of our person's hands affair!** This just means that the **Data was not showing strong Evidence against that he is not guilty** - maybe this is because we do not have enough Data.

# Notes

At this point, Statisticians always compare this with the process in the court: if the person is suspected, say, in murder, then, by default, judges assume that he is **not guilty** (anmeghutyan kankhavarkac 😊). This is our Null Hypothesis:

$\mathcal{H}_0$  : He is not guilty    vs     $\mathcal{H}_1$  : He is guilty.

Now, based on the evidence, Data, the judges either will Reject  $\mathcal{H}_0$ , meaning that, upon the light of a lot of facts, they will decide that our person is guilty, or, if the Data will not be enough, the judges will Not Reject  $\mathcal{H}_0$ , so they will free our person. **But this does not mean that the murder was not of our person's hands affair!** This just means that the **Data was not showing strong Evidence against that he is not guilty** - maybe this is because we do not have enough Data.