

# AUA CS 108, Statistics, Fall 2019

## Lecture 34

Michael Poghosyan

YSU, AUA

[michael@ysu.am](mailto:michael@ysu.am), [mpoghosyan@aua.am](mailto:mpoghosyan@aua.am)

11 Nov 2019

# Contents

- ▶ AsympTotic CI-s

## Last Lecture ReCap

- ▶ Give the definition of the  $\chi^2(n)$  Distribution.

## Last Lecture ReCap

- ▶ Give the definition of the  $\chi^2(n)$  Distribution.
- ▶ Give the  $(1 - \alpha)$ -level CI for  $\sigma^2$  in the Normal Model.

## CI for $\sigma^2$ , $\mathcal{N}(\mu, \sigma^2)$ Model, $\mu$ is **unknown**

Recall that, for the Model  $\mathcal{N}(\mu, \sigma^2)$ , in the case when  $\mu$  was unknown, we have obtained the following  $(1 - \alpha)$ -level CI for  $\sigma^2$ :

## CI for $\sigma^2$ , $\mathcal{N}(\mu, \sigma^2)$ Model, $\mu$ is **unknown**

Recall that, for the Model  $\mathcal{N}(\mu, \sigma^2)$ , in the case when  $\mu$  was unknown, we have obtained the following  $(1 - \alpha)$ -level CI for  $\sigma^2$ :

$$\left( \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{\chi_{n-1, 1-\frac{\alpha}{2}}^2}, \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{\chi_{n-1, \frac{\alpha}{2}}^2} \right).$$

## CI for $\sigma^2$ , $\mathcal{N}(\mu, \sigma^2)$ Model, $\mu$ is **unknown**

Recall that, for the Model  $\mathcal{N}(\mu, \sigma^2)$ , in the case when  $\mu$  was unknown, we have obtained the following  $(1 - \alpha)$ -level CI for  $\sigma^2$ :

$$\left( \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{\chi_{n-1, 1-\frac{\alpha}{2}}^2}, \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{\chi_{n-1, \frac{\alpha}{2}}^2} \right).$$

This is the same as

$$\left( \frac{(n-1) \cdot S^2}{\chi_{n-1, 1-\frac{\alpha}{2}}^2}, \frac{(n-1) \cdot S^2}{\chi_{n-1, \frac{\alpha}{2}}^2} \right),$$

where  $S$  is the Sample Standard Deviation given by

$$S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}.$$

## Example

**Example:** Assume we have scale of very high precision, and we want to measure the precision of our scale. Say, we weight a 50AMD coin, using that scale, 10 times, and we obtain the following measurements (in grams):

[1] 3.449243 3.450802 3.453054 3.448778 3.452541 3.448835 3.448

[9] 3.450314 3.449047

Our aim is to Estimate the Precision of the Scale.



## Example

**Example:** Assume we have scale of very high precision, and we want to measure the precision of our scale. Say, we weight a 50AMD coin, using that scale, 10 times, and we obtain the following measurements (in grams):

```
[1] 3.449243 3.450802 3.453054 3.448778 3.452541 3.448835 3.448  
[9] 3.450314 3.449047
```

Our aim is to Estimate the Precision of the Scale.

We model this as: say, we weight, by using our scale, something with **exact** weight  $\mu$ , but we do not know  $\mu$ .

## Example

**Example:** Assume we have scale of very high precision, and we want to measure the precision of our scale. Say, we weight a 50AMD coin, using that scale, 10 times, and we obtain the following measurements (in grams):

```
[1] 3.449243 3.450802 3.453054 3.448778 3.452541 3.448835 3.448  
[9] 3.450314 3.449047
```

Our aim is to Estimate the Precision of the Scale.

We model this as: say, we weight, by using our scale, something with **exact** weight  $\mu$ , but we do not know  $\mu$ . Our scale will show a number around, very close to  $\mu$ , let it be, let it be  $W$ .

## Example

**Example:** Assume we have scale of very high precision, and we want to measure the precision of our scale. Say, we weight a 50AMD coin, using that scale, 10 times, and we obtain the following measurements (in grams):

```
[1] 3.449243 3.450802 3.453054 3.448778 3.452541 3.448835 3.448  
[9] 3.450314 3.449047
```

Our aim is to Estimate the Precision of the Scale.

We model this as: say, we weight, by using our scale, something with **exact** weight  $\mu$ , but we do not know  $\mu$ . Our scale will show a number around, very close to  $\mu$ , let it be, let it be  $W$ . So we will assume

$$W \sim$$

## Example

**Example:** Assume we have scale of very high precision, and we want to measure the precision of our scale. Say, we weight a 50AMD coin, using that scale, 10 times, and we obtain the following measurements (in grams):

[1] 3.449243 3.450802 3.453054 3.448778 3.452541 3.448835 3.448  
[9] 3.450314 3.449047

Our aim is to Estimate the Precision of the Scale.

We model this as: say, we weight, by using our scale, something with **exact** weight  $\mu$ , but we do not know  $\mu$ . Our scale will show a number around, very close to  $\mu$ , let it be, let it be  $W$ . So we will assume

$$W \sim \mathcal{N}(\mu, \sigma^2),$$

where  $\sigma^2$  (or, better,  $\sigma$ ) is measuring our scale Precision.

## Example

**Example:** Assume we have scale of very high precision, and we want to measure the precision of our scale. Say, we weight a 50AMD coin, using that scale, 10 times, and we obtain the following measurements (in grams):

```
[1] 3.449243 3.450802 3.453054 3.448778 3.452541 3.448835 3.448  
[9] 3.450314 3.449047
```

Our aim is to Estimate the Precision of the Scale.

We model this as: say, we weight, by using our scale, something with **exact** weight  $\mu$ , but we do not know  $\mu$ . Our scale will show a number around, very close to  $\mu$ , let it be, let it be  $W$ . So we will assume

$$W \sim \mathcal{N}(\mu, \sigma^2),$$

where  $\sigma^2$  (or, better,  $\sigma$ ) is measuring our scale Precision.

So now, using the above observations (weighting results), we will construct a 90% CI for  $\sigma^2$ .

## Example, Cont'd

Recall the  $(1 - \alpha)$ -level CI for  $\sigma^2$ :

$$\left( \frac{(n-1) \cdot S^2}{\chi_{n-1, 1-\frac{\alpha}{2}}^2}, \frac{(n-1) \cdot S^2}{\chi_{n-1, \frac{\alpha}{2}}^2} \right).$$

## Example, Cont'd

Recall the  $(1 - \alpha)$ -level CI for  $\sigma^2$ :

$$\left( \frac{(n-1) \cdot S^2}{\chi^2_{n-1, 1-\frac{\alpha}{2}}}, \frac{(n-1) \cdot S^2}{\chi^2_{n-1, \frac{\alpha}{2}}} \right).$$

Now, we use **R** to do the rest:

## Example, Cont'd

Recall the  $(1 - \alpha)$ -level CI for  $\sigma^2$ :

$$\left( \frac{(n-1) \cdot S^2}{\chi_{n-1, 1-\frac{\alpha}{2}}^2}, \frac{(n-1) \cdot S^2}{\chi_{n-1, \frac{\alpha}{2}}^2} \right).$$

Now, we use **R** to do the rest: we want to construct 90% CI, so our  $\alpha =$



## Example, Cont'd

Recall the  $(1 - \alpha)$ -level CI for  $\sigma^2$ :

$$\left( \frac{(n-1) \cdot S^2}{\chi_{n-1, 1-\frac{\alpha}{2}}^2}, \frac{(n-1) \cdot S^2}{\chi_{n-1, \frac{\alpha}{2}}^2} \right).$$

Now, we use **R** to do the rest: we want to construct 90% CI, so our  $\alpha = 0.1$ .

## Example, Cont'd

Recall the  $(1 - \alpha)$ -level CI for  $\sigma^2$ :

$$\left( \frac{(n-1) \cdot S^2}{\chi_{n-1, 1-\frac{\alpha}{2}}^2}, \frac{(n-1) \cdot S^2}{\chi_{n-1, \frac{\alpha}{2}}^2} \right).$$

Now, we use **R** to do the rest: we want to construct 90% CI, so our  $\alpha = 0.1$ . We have 10 observations, so  $n = 10$ .

## Example, Cont'd

Recall the  $(1 - \alpha)$ -level CI for  $\sigma^2$ :

$$\left( \frac{(n-1) \cdot S^2}{\chi_{n-1, 1-\frac{\alpha}{2}}^2}, \frac{(n-1) \cdot S^2}{\chi_{n-1, \frac{\alpha}{2}}^2} \right).$$

Now, we use **R** to do the rest: we want to construct 90% CI, so our  $\alpha = 0.1$ . We have 10 observations, so  $n = 10$ . We calculate  $S^2$ :

```
w <- c(3.449243, 3.450802, 3.453054, 3.448778, 3.452541, 3.451234, 3.449876, 3.450123, 3.451567, 3.449987)
s2 <- var(w)
s2

## [1] 4.605341e-06
```

## Example, Cont'd

The Degrees of Freedom in our case, since we do not know  $\mu$ , is

## Example, Cont'd

The Degrees of Freedom in our case, since we do not know  $\mu$ , is  $n - 1 = 9$ , and we calculate the corresponding quantiles for the  $\chi^2(9)$ :

```
alpha <- 0.1  
lq<-qchisq(alpha/2, df=9); uq<-qchisq(1-alpha/2, df=9)  
c(lq,uq)  
  
## [1] 3.325113 16.918978
```

## Example, Cont'd

The Degrees of Freedom in our case, since we do not know  $\mu$ , is  $n - 1 = 9$ , and we calculate the corresponding quantiles for the  $\chi^2(9)$ :

```
alpha <- 0.1  
lq<-qchisq(alpha/2, df=9); uq<-qchisq(1-alpha/2, df=9)  
c(lq,uq)
```

```
## [1] 3.325113 16.918978
```

Finally, we calculate our CI endpoints:

```
n <- 10  
c((n-1)*s2/uq, (n-1)*s2/lq)
```

```
## [1] 2.449797e-06 1.246516e-05
```

## Example, Cont'd

The Degrees of Freedom in our case, since we do not know  $\mu$ , is  $n - 1 = 9$ , and we calculate the corresponding quantiles for the  $\chi^2(9)$ :

```
alpha <- 0.1  
lq<-qchisq(alpha/2, df=9); uq<-qchisq(1-alpha/2, df=9)  
c(lq,uq)
```

```
## [1] 3.325113 16.918978
```

Finally, we calculate our CI endpoints:

```
n <- 10  
c((n-1)*s2/uq, (n-1)*s2/lq)
```

```
## [1] 2.449797e-06 1.246516e-05
```

**Note:** The actual value of  $sd$  I was using was:  $sd = 0.002$ , so the true value of my  $\sigma^2$  was

$$\sigma^2 = 4 \cdot 10^{-6}.$$

## CI for $\sigma^2$ , $\mathcal{N}(\mu, \sigma^2)$ Model, Summary

Again, as above, let us summarize what we have obtained for this model. The problem is: given a Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2),$$

and  $\alpha \in (0, 1)$ , we want to construct an  $1 - \alpha$ -level CI for the unknown parameter  $\sigma^2$ .



## CI for $\sigma^2$ , $\mathcal{N}(\mu, \sigma^2)$ Model, Summary

Again, as above, let us summarize what we have obtained for this model. The problem is: given a Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2),$$

and  $\alpha \in (0, 1)$ , we want to construct an  $1 - \alpha$ -level CI for the unknown parameter  $\sigma^2$ .

---

	$\mu$ is known	$\mu$ is unknown
--	----------------	------------------

Piv:

## CI for $\sigma^2$ , $\mathcal{N}(\mu, \sigma^2)$ Model, Summary

Again, as above, let us summarize what we have obtained for this model. The problem is: given a Random Sample

$$X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2),$$

and  $\alpha \in (0, 1)$ , we want to construct an  $1 - \alpha$ -level CI for the unknown parameter  $\sigma^2$ .

	$\mu$ is known	$\mu$ is unknown
Piv:	$\chi^2 = \sum_{k=1}^n \left( \frac{X_k - \mu}{\sigma} \right)^2$	$\chi^2 = \sum_{k=1}^n \left( \frac{X_k - \bar{X}}{\sigma} \right)^2$
Dis of Piv		

## CI for $\sigma^2$ , $\mathcal{N}(\mu, \sigma^2)$ Model, Summary

Again, as above, let us summarize what we have obtained for this model. The problem is: given a Random Sample

$$X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2),$$

and  $\alpha \in (0, 1)$ , we want to construct an  $1 - \alpha$ -level CI for the unknown parameter  $\sigma^2$ .

	$\mu$ is known	$\mu$ is unknown
Piv:	$\chi^2 = \sum_{k=1}^n \left( \frac{X_k - \mu}{\sigma} \right)^2$	$\chi^2 = \sum_{k=1}^n \left( \frac{X_k - \bar{X}}{\sigma} \right)^2$
Dis of Piv	$\chi^2 \sim \chi^2(n)$	$\chi^2 \sim \chi^2(n-1)$
Reg w/ $\mathbb{P}$ $1-\alpha$		

## CI for $\sigma^2$ , $\mathcal{N}(\mu, \sigma^2)$ Model, Summary

Again, as above, let us summarize what we have obtained for this model. The problem is: given a Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2),$$

and  $\alpha \in (0, 1)$ , we want to construct an  $1 - \alpha$ -level CI for the unknown parameter  $\sigma^2$ .

	$\mu$ is known	$\mu$ is unknown
Piv:	$\chi^2 = \sum_{k=1}^n \left( \frac{X_k - \mu}{\sigma} \right)^2$	$\chi^2 = \sum_{k=1}^n \left( \frac{X_k - \bar{X}}{\sigma} \right)^2$
Dis of Piv	$\chi^2 \sim \chi^2(n)$	$\chi^2 \sim \chi^2(n-1)$
Reg w/ $\mathbb{P}$ $1-\alpha$	$\chi_{n,\alpha/2}^2 < \chi^2 < \chi_{n,1-\alpha/2}^2$	$\chi_{n-1,\alpha/2}^2 < \chi^2 < \chi_{n-1,1-\alpha/2}^2$
CI for $\sigma^2$ :		

## CI for $\sigma^2$ , $\mathcal{N}(\mu, \sigma^2)$ Model, Summary

Again, as above, let us summarize what we have obtained for this model. The problem is: given a Random Sample

$$X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2),$$

and  $\alpha \in (0, 1)$ , we want to construct an  $1 - \alpha$ -level CI for the unknown parameter  $\sigma^2$ .

	$\mu$ is known	$\mu$ is unknown
Piv:	$\chi^2 = \sum_{k=1}^n \left( \frac{X_k - \mu}{\sigma} \right)^2$	$\chi^2 = \sum_{k=1}^n \left( \frac{X_k - \bar{X}}{\sigma} \right)^2$
Dis of Piv	$\chi^2 \sim \chi^2(n)$	$\chi^2 \sim \chi^2(n-1)$
Reg w/ $\mathbb{P}$ $1-\alpha$	$\chi_{n,\alpha/2}^2 < \chi^2 < \chi_{n,1-\alpha/2}^2$	$\chi_{n-1,\alpha/2}^2 < \chi^2 < \chi_{n-1,1-\alpha/2}^2$
CI for $\sigma^2$ :	$\left( \frac{\sum_{k=1}^n (X_k - \mu)^2}{\chi_{n,1-\alpha/2}^2}, \frac{\sum_{k=1}^n (X_k - \mu)^2}{\chi_{n,\alpha/2}^2} \right)$	$\left( \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{\chi_{n-1,1-\alpha/2}^2}, \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{\chi_{n-1,\alpha/2}^2} \right)$

Asymptotic CIs

# Asymptotic CIs

Up to now, we have constructed CIs for our important basic Models - say, Bernoulli and Normal. Unfortunately, it is not an easy job to construct CIs for any Parametric Model.

# Asymptotic CIs

Up to now, we have constructed CIs for our important basic Models - say, Bernoulli and Normal. Unfortunately, it is not an easy job to construct CIs for any Parametric Model. But fortunately, if the Sample Size is large enough, CLT and  $C^\circ$  are helping to construct an *Asymptotic CIs*, covering the unknown Parameter value with high Probability for large Sample Size.



# Asymptotic CIs

Up to now, we have constructed CIs for our important basic Models - say, Bernoulli and Normal. Unfortunately, it is not an easy job to construct CIs for any Parametric Model. But fortunately, if the Sample Size is large enough, CLT and  $C^\circ$  are helping to construct an *Asymptotic CIs*, covering the unknown Parameter value with high Probability for large Sample Size.

Let us formalize the notion.

# Asymptotic CIs

Up to now, we have constructed CIs for our important basic Models - say, Bernoulli and Normal. Unfortunately, it is not an easy job to construct CIs for any Parametric Model. But fortunately, if the Sample Size is large enough, CLT and  $C^o$  are helping to construct an *Asymptotic CIs*, covering the unknown Parameter value with high Probability for large Sample Size.

Let us formalize the notion.

Again we consider a Parametric Model  $\mathcal{F}_\theta$ , assuming that we have an (infinite) Random Sample from one of these Distributions:

$$X_1, X_2, \dots, X_n, \dots \sim \mathcal{F}_\theta.$$

# Asymptotic CIs

Up to now, we have constructed CIs for our important basic Models - say, Bernoulli and Normal. Unfortunately, it is not an easy job to construct CIs for any Parametric Model. But fortunately, if the Sample Size is large enough, CLT and  $C^\circ$  are helping to construct an *Asymptotic CIs*, covering the unknown Parameter value with high Probability for large Sample Size.

Let us formalize the notion.

Again we consider a Parametric Model  $\mathcal{F}_\theta$ , assuming that we have an (infinite) Random Sample from one of these Distributions:

$$X_1, X_2, \dots, X_n, \dots \sim \mathcal{F}_\theta.$$

To construct an Asymptotic CI for  $\theta$ , we take  $\alpha \in (0, 1)$ .

# Asymptotic CIs

**Definition:** Assume, for any  $n$ ,  $L_n = L_n(x_1, \dots, x_n, \alpha)$ ,  $U_n = U_n(x_1, \dots, x_n, \alpha)$  be two functions with  $L_n(x_1, \dots, x_n, \alpha) \leq U_n(x_1, \dots, x_n, \alpha)$  for all  $(x_1, \dots, x_n, \alpha)$ . The sequence of Random Intervals

$$(L_n, U_n) = (L_n(X_1, \dots, X_n, \alpha); U_n(X_1, \dots, X_n, \alpha))$$

is called an **Asymptotic Confidence Interval sequence** (or just an Asymptotic Confidence Interval for  $\theta$  of (Asymptotic) level  $1 - \alpha$ , if for any  $\theta \in \Theta$ ,

$$\liminf_{n \rightarrow +\infty} \mathbb{P}(L_n < \theta < U_n) \geq 1 - \alpha.$$

# Asymptotic CIs

**Definition:** Assume, for any  $n$ ,  $L_n = L_n(x_1, \dots, x_n, \alpha)$ ,  $U_n = U_n(x_1, \dots, x_n, \alpha)$  be two functions with  $L_n(x_1, \dots, x_n, \alpha) \leq U_n(x_1, \dots, x_n, \alpha)$  for all  $(x_1, \dots, x_n, \alpha)$ . The sequence of Random Intervals

$$(L_n, U_n) = (L_n(X_1, \dots, X_n, \alpha); U_n(X_1, \dots, X_n, \alpha))$$

is called an **Asymptotic Confidence Interval sequence** (or just an Asymptotic Confidence Interval for  $\theta$  of (Asymptotic) level  $1 - \alpha$ , if for any  $\theta \in \Theta$ ,

$$\liminf_{n \rightarrow +\infty} \mathbb{P}(L_n < \theta < U_n) \geq 1 - \alpha.$$

Usually, we will have that the limit above exists, so we will use

$$\lim_{n \rightarrow +\infty} \mathbb{P}(L_n < \theta < U_n) \geq 1 - \alpha.$$

## Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want the Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ .

## Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want the Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ . Here we assume that the Variance  $\sigma^2 = \text{Var}(X_k)$  is finite, but unknown.

## Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want the Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ . Here we assume that the Variance  $\sigma^2 = \text{Var}(X_k)$  is finite, but unknown.

So our Problem is to construct an Asymptotic CI of level  $1 - \alpha$  for  $\mu$ .



## Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want the Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ . Here we assume that the Variance  $\sigma^2 = \text{Var}(X_k)$  is finite, but unknown.

So our Problem is to construct an Asymptotic CI of level  $1 - \alpha$  for  $\mu$ .

We consider the following  $t$ -Statistics (or, rather, a sequence of Statistics):

$$t_n = \frac{\bar{X}_n - \mu}{S_n / \sqrt{n}}.$$

where  $(S_n)^2 = \frac{\sum_{k=1}^n (X_k - \bar{X}_n)^2}{n-1}$  is the Sample SD.

# Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want the Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ . Here we assume that the Variance  $\sigma^2 = \text{Var}(X_k)$  is finite, but unknown.

So our Problem is to construct an Asymptotic CI of level  $1 - \alpha$  for  $\mu$ .

We consider the following  $t$ -Statistics (or, rather, a sequence of Statistics):

$$t_n = \frac{\bar{X}_n - \mu}{S_n / \sqrt{n}}.$$

where  $(S_n)^2 = \frac{\sum_{k=1}^n (X_k - \bar{X}_n)^2}{n-1}$  is the Sample SD. Unfortunately, this is not a Pivot anymore, in the general case - the Distribution of  $t_n$  was  $t(n-1)$  **only in the case when**  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ .

# Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want the Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ . Here we assume that the Variance  $\sigma^2 = \text{Var}(X_k)$  is finite, but unknown.

So our Problem is to construct an Asymptotic CI of level  $1 - \alpha$  for  $\mu$ .

We consider the following  $t$ -Statistics (or, rather, a sequence of Statistics):

$$t_n = \frac{\bar{X}_n - \mu}{S_n / \sqrt{n}}.$$

where  $(S_n)^2 = \frac{\sum_{k=1}^n (X_k - \bar{X}_n)^2}{n-1}$  is the Sample SD. Unfortunately, this is not a Pivot anymore, in the general case - the Distribution of  $t_n$  was  $t(n-1)$  **only in the case when**  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ . Fortunately, the Asymptotic Distribution is independent on  $\mu$ , so  $t_n$  is an *Asymptotic Pivot*.

# Asymptotic CI for the Mean of General Distribution

Assume we have an observation from a Random Sample  $X_1, X_2, \dots, X_n, \dots$ . We want the Estimate, using CIs, the Mean  $\mu = \mathbb{E}(X_k)$ . Here we assume that the Variance  $\sigma^2 = \text{Var}(X_k)$  is finite, but unknown.

So our Problem is to construct an Asymptotic CI of level  $1 - \alpha$  for  $\mu$ .

We consider the following  $t$ -Statistics (or, rather, a sequence of Statistics):

$$t_n = \frac{\bar{X}_n - \mu}{S_n / \sqrt{n}}.$$

where  $(S_n)^2 = \frac{\sum_{k=1}^n (X_k - \bar{X}_n)^2}{n-1}$  is the Sample SD. Unfortunately, this is not a Pivot anymore, in the general case - the Distribution of  $t_n$  was  $t(n-1)$  **only in the case when**  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ . Fortunately, the Asymptotic Distribution is independent on  $\mu$ , so  $t_n$  is an *Asymptotic Pivot*. To see this, we can write  $t_n$  as:

## Asymptotic CI for the Mean of General Distribution

$$t_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{S_n}.$$

## Asymptotic CI for the Mean of General Distribution

$$t_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{S_n}.$$

Now, by the CLT, the first factor tends to  $\mathcal{N}(0, 1)$  in Distributions, and the second one, as can be proved, tends to 1 in Probability.

## Asymptotic CI for the Mean of General Distribution

$$t_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{S_n}.$$

Now, by the CLT, the first factor tends to  $\mathcal{N}(0, 1)$  in Distributions, and the second one, as can be proved, tends to 1 in Probability.

From this we can obtain that

$$t_n \xrightarrow{D} \mathcal{N}(0, 1).$$

## Asymptotic CI for the Mean of General Distribution

$$t_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{S_n}.$$

Now, by the CLT, the first factor tends to  $\mathcal{N}(0, 1)$  in Distributions, and the second one, as can be proved, tends to 1 in Probability.

From this we can obtain that

$$t_n \xrightarrow{D} \mathcal{N}(0, 1).$$

So the Asymptotic Distribution of  $t_n$  is independent of  $\mu$ .



## Asymptotic CI for the Mean of General Distribution

$$t_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{S_n}.$$

Now, by the CLT, the first factor tends to  $\mathcal{N}(0, 1)$  in Distributions, and the second one, as can be proved, tends to 1 in Probability.

From this we can obtain that

$$t_n \xrightarrow{D} \mathcal{N}(0, 1).$$

So the Asymptotic Distribution of  $t_n$  is independent of  $\mu$ .

The rest is standard: first we find numbers  $a, b$  such that

$$\mathbb{P}(a < Z < b) = 1 - \alpha, \quad \text{where } Z \sim \mathcal{N}(0, 1).$$

## Asymptotic CI for the Mean of General Distribution

$$t_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{S_n}.$$

Now, by the CLT, the first factor tends to  $\mathcal{N}(0, 1)$  in Distributions, and the second one, as can be proved, tends to 1 in Probability.

From this we can obtain that

$$t_n \xrightarrow{D} \mathcal{N}(0, 1).$$

So the Asymptotic Distribution of  $t_n$  is independent of  $\mu$ .

The rest is standard: first we find numbers  $a, b$  such that

$$\mathbb{P}(a < Z < b) = 1 - \alpha, \quad \text{where } Z \sim \mathcal{N}(0, 1).$$

We can take, as previously,  $-a = b = z_{1-\alpha/2}$ .

## Asymptotic CI for the Mean of General Distribution

$$t_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{S_n}.$$

Now, by the CLT, the first factor tends to  $\mathcal{N}(0, 1)$  in Distributions, and the second one, as can be proved, tends to 1 in Probability.

From this we can obtain that

$$t_n \xrightarrow{D} \mathcal{N}(0, 1).$$

So the Asymptotic Distribution of  $t_n$  is independent of  $\mu$ .

The rest is standard: first we find numbers  $a, b$  such that

$$\mathbb{P}(a < Z < b) = 1 - \alpha, \quad \text{where } Z \sim \mathcal{N}(0, 1).$$

We can take, as previously,  $-a = b = z_{1-\alpha/2}$ . Then

$$\mathbb{P}(-z_{1-\alpha/2} < Z < z_{1-\alpha/2}) = 1 - \alpha.$$

## Asymptotic CI for the Mean of General Distribution

$$t_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \cdot \frac{\sigma}{S_n}.$$

Now, by the CLT, the first factor tends to  $\mathcal{N}(0, 1)$  in Distributions, and the second one, as can be proved, tends to 1 in Probability.

From this we can obtain that

$$t_n \xrightarrow{D} \mathcal{N}(0, 1).$$

So the Asymptotic Distribution of  $t_n$  is independent of  $\mu$ .

The rest is standard: first we find numbers  $a, b$  such that

$$\mathbb{P}(a < Z < b) = 1 - \alpha, \quad \text{where } Z \sim \mathcal{N}(0, 1).$$

We can take, as previously,  $-a = b = z_{1-\alpha/2}$ . Then

$$\mathbb{P}(-z_{1-\alpha/2} < Z < z_{1-\alpha/2}) = 1 - \alpha.$$

Since  $t_n \xrightarrow{D} \mathcal{N}(0, 1)$ , then

$$\mathbb{P}(-z_{1-\alpha/2} < t_n < z_{1-\alpha/2}) \rightarrow \mathbb{P}(-z_{1-\alpha/2} < Z < z_{1-\alpha/2}) = 1 - \alpha.$$

## Asymptotic CI for the Mean of General Distribution

We plug the value of  $t_n$  here and solve for  $\mu$  to obtain

$$\mathbb{P}\left(\bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} < \mu < \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}\right) \rightarrow 1 - \alpha$$

## Asymptotic CI for the Mean of General Distribution

We plug the value of  $t_n$  here and solve for  $\mu$  to obtain

$$\mathbb{P}\left(\bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} < \mu < \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}\right) \rightarrow 1 - \alpha$$

so the Random Interval (or, rather, the sequence of Intervals)

$$\left(\bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}\right)$$

is a  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ .

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:

$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$



## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:

$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

This is because

- ▶ the same form of CI was obtained for the Normal Model, when  $\sigma^2$  was unknown;

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:

$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

This is because

- ▶ the same form of CI was obtained for the Normal Model, when  $\sigma^2$  was unknown;
- ▶ this interval is a little bit larger than the previous one, so it is also an AsympCI for  $\mu$  of level  $1 - \alpha$ ;

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:

$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \quad \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

This is because

- ▶ the same form of CI was obtained for the Normal Model, when  $\sigma^2$  was unknown;
- ▶ this interval is a little bit larger than the previous one, so it is also an AsympCI for  $\mu$  of level  $1 - \alpha$ ;
- ▶ when  $n \geq 30$ , these two almost coincide;

## Asymptotic CI for the Mean of General Distribution

**Note:** We have obtained the following  $(1 - \alpha)$ -level Asymptotic CI for  $\mu$ :

$$\left( \bar{X}_n - z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \bar{X}_n + z_{1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

Usually, people use not this one, but the following one:


$$\left( \bar{X}_n - t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}}; \bar{X}_n + t_{n-1, 1-\alpha/2} \cdot \frac{S_n}{\sqrt{n}} \right)$$

This is because

- ▶ the same form of CI was obtained for the Normal Model, when  $\sigma^2$  was unknown;
- ▶ this interval is a little bit larger than the previous one, so it is also an AsympCI for  $\mu$  of level  $1 - \alpha$ ;
- ▶ when  $n \geq 30$ , these two almost coincide;
- ▶ although in the theory these intervals work for large  $n$ , but, in practice, the latter one works also for small  $n$


# Example

Asymptotic CI for the Mean with  $z$ ,  $n = 50$



# Example

Asymptotic CI for the Mean with  $z$ ,  $n = 20$



# Example

