

AUA CS108, Statistics, Fall 2020

Lecture 35

Michael Poghosyan

16 Nov 2020

Contents

- ▶ Maximum Likelihood Method (MLE)
- ▶ Confidence Intervals

Examples, MLE

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

X	0	1	2
$\mathbb{P}(X = x)$	$\frac{\theta}{10}$	$\frac{\theta}{5}$	$1 - \frac{3\theta}{10}$

where $\theta \in [0, \frac{10}{3}]$.

Examples, MLE

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

$$\begin{array}{c|c|c|c} X & 0 & 1 & 2 \\ \hline \mathbb{P}(X = x) & \frac{\theta}{10} & \frac{\theta}{5} & 1 - \frac{3\theta}{10}, \end{array}$$

where $\theta \in [0, \frac{10}{3}]$. Find the MLE Estimator and MLE Estimate for θ .

Solution: OTB

Examples, MLE

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

$$\begin{array}{c|c|c|c} X & 0 & 1 & 2 \\ \hline \mathbb{P}(X = x) & \frac{\theta}{10} & \frac{\theta}{5} & 1 - \frac{3\theta}{10}, \end{array}$$

where $\theta \in [0, \frac{10}{3}]$. Find the MLE Estimator and MLE Estimate for θ .

Solution: OTB

Example: Find the MLE Estimator for (μ, σ^2) in the $\mathcal{N}(\mu, \sigma^2)$ Model.

Solution: OTB

Some Notes about MLE

Note: The MLE does not always exist;

Some Notes about MLE

Note: The MLE does not always exist;

Note: The MLE is not always unique;

Some Notes about MLE

Note: The MLE does not always exist;

Note: The MLE is not always unique;

Note: Sometimes it is necessary to find it numerically, say, using the NR Method

Some Notes about MLE

Note: The MLE does not always exist;

Note: The MLE is not always unique;

Note: Sometimes it is necessary to find it numerically, say, using the NR Method

Note: It is remarkable, that ML Estimators, in general (if they exist, of course 😊), possess some nice properties. These properties make MLE one of the widely used methods of Estimation.

Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family \mathcal{F}_θ ,

- ▶ The MLE $\hat{\theta}_n^{MLE}$ is Consistent, i.e.,

Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family \mathcal{F}_θ ,

- ▶ The MLE $\hat{\theta}_n^{MLE}$ is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family \mathcal{F}_θ ,

- ▶ The MLE $\hat{\theta}_n^{MLE}$ is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

- ▶ The MLE $\hat{\theta}_n^{MLE}$ is Asymptotically Normal and Efficient:

$$\hat{\theta}_n^{MLE} \overset{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

Properties of the MLE, Cont'd

We can interpret this as:

- ▶ $\hat{\theta}_n^{MLE}$ is approximately (Asymptotically) Normal;

Properties of the MLE, Cont'd

We can interpret this as:

- ▶ $\hat{\theta}_n^{MLE}$ is approximately (Asymptotically) Normal;
- ▶ The Variance of $\hat{\theta}_n^{MLE}$ is approximately $\frac{1}{n \cdot \mathcal{I}(\theta)}$. This is the possible smallest Variance we can have with Unbiased Estimators.

Properties of the MLE, Cont'd

We can interpret this as:

- ▶ $\hat{\theta}_n^{MLE}$ is approximately (Asymptotically) Normal;
- ▶ The Variance of $\hat{\theta}_n^{MLE}$ is approximately $\frac{1}{n \cdot \mathcal{I}(\theta)}$. This is the possible smallest Variance we can have with Unbiased Estimators. So we interpret this as **MLE is Asymptotically Efficient**.

Properties of the MLE, Cont'd

We can interpret this as:

- ▶ $\hat{\theta}_n^{MLE}$ is approximately (Asymptotically) Normal;
- ▶ The Variance of $\hat{\theta}_n^{MLE}$ is approximately $\frac{1}{n \cdot \mathcal{I}(\theta)}$. This is the possible smallest Variance we can have with Unbiased Estimators. So we interpret this as **MLE is Asymptotically Efficient**.

So, MLE is **Consistent** and **Asymptotically Efficient**.

Properties of the MLE, Cont'd

We can interpret this as:

- ▶ $\hat{\theta}_n^{MLE}$ is approximately (Asymptotically) Normal;
- ▶ The Variance of $\hat{\theta}_n^{MLE}$ is approximately $\frac{1}{n \cdot \mathcal{I}(\theta)}$. This is the possible smallest Variance we can have with Unbiased Estimators. So we interpret this as **MLE is Asymptotically Efficient**.

So, MLE is **Consistent** and **Asymptotically Efficient**. And this is why, for large Sample Size n , MLE is the Top 1 Choice, is (almost) unbeatable.

Fisher Information

in the above formulas,

$$\mathcal{I}(\theta) = -\mathbb{E} \left(\frac{\partial^2}{\partial \theta^2} \ln f(X|\theta) \right) = \mathbb{E} \left[\left(\frac{\partial}{\partial \theta} \ln f(X|\theta) \right)^2 \right],$$

is the **Fisher Information** for θ (X is a r.v. from the Distribution \mathcal{F}_θ , and $f(x|\theta)$ is the corresponding PD(M)F).

Properties of the MLE, Cont'd

► Also,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}\right)$$

Properties of the MLE, Cont'd

► Also,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}\right)$$

Note: This is almost the above Property,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right),$$

but, instead of $\mathcal{I}(\theta)$ we have $\mathcal{I}(\hat{\theta}_n^{MLE})$.

Properties of the MLE, Cont'd

► Also,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}\right)$$

Note: This is almost the above Property,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right),$$

but, instead of $\mathcal{I}(\theta)$ we have $\mathcal{I}(\hat{\theta}_n^{MLE})$.

Note: We will use this later, to construct an (approximate) Confidence Interval for θ and for testing Hypotheses about θ .

Properties of the MLE, Cont'd

- ▶ If $\hat{\theta}$ is the MLE for θ , then for any function g , the MLE of $g(\theta)$ is $g(\hat{\theta})$, i.e.,

$$\widehat{g(\theta)}^{MLE} = g\left(\hat{\theta}^{MLE}\right).$$

Properties of the MLE, Cont'd

- ▶ If $\hat{\theta}$ is the MLE for θ , then for any function g , the MLE of $g(\theta)$ is $g(\hat{\theta})$, i.e.,

$$\widehat{g(\theta)}^{MLE} = g\left(\hat{\theta}^{MLE}\right).$$

This is sometimes called the **invariance Principle of MLE**.

Properties of the MLE, Cont'd

- ▶ If $\hat{\theta}$ is the MLE for θ , then for any function g , the MLE of $g(\theta)$ is $g(\hat{\theta})$, i.e.,

$$\widehat{g(\theta)}^{MLE} = g\left(\hat{\theta}^{MLE}\right).$$

This is sometimes called the **invariance Principle of MLE**.

Example Find the MLE for σ in $\mathcal{N}(\mu, \sigma^2)$ Model.

Solution: OTB

Other Methods to construct Point Estimators/Estimates

There are other important methods to construct Estimators: e.g.

- ▶ Bayesian Estimation: Maximum APosteriori (MAP) Estimators
- ▶ Bayesian Estimation: Bayes Estimators;
- ▶ OLS
- ▶ etc

Confidence Intervals

Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume θ is our Parameter to be Estimated, and $\hat{\theta}$ is a good Estimator for θ .

Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume θ is our Parameter to be Estimated, and $\hat{\theta}$ is a good Estimator for θ . Assume $\hat{\theta}$ is a Continuous r.v.

Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume θ is our Parameter to be Estimated, and $\hat{\theta}$ is a good Estimator for θ . Assume $\hat{\theta}$ is a Continuous r.v. If the True value of our Parameter is θ^* , then

$$\mathbb{P}(\hat{\theta} = \theta^*) = 0,$$

i.e., we will (almost) **never** be correct in our guess.

Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume θ is our Parameter to be Estimated, and $\hat{\theta}$ is a good Estimator for θ . Assume $\hat{\theta}$ is a Continuous r.v. If the True value of our Parameter is θ^* , then

$$\mathbb{P}(\hat{\theta} = \theta^*) = 0,$$

i.e., we will (almost) **never** be correct in our guess. Sad news!

Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using $\hat{\theta}$, if $\hat{\theta}$ possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for θ^* .

¹Recall the \widehat{SE} , the Estimated Standard Error reporting story.

Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using $\hat{\theta}$, if $\hat{\theta}$ possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for θ^* .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small¹,

¹Recall the \widehat{SE} , the Estimated Standard Error reporting story.

Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using $\hat{\theta}$, if $\hat{\theta}$ possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for θ^* .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small¹, or, rather, as

$$\theta^* \in (\hat{\theta} - \text{error}, \hat{\theta} + \text{error}).$$

¹Recall the \widehat{SE} , the Estimated Standard Error reporting story.

Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using $\hat{\theta}$, if $\hat{\theta}$ possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for θ^* .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small¹, or, rather, as

$$\theta^* \in (\hat{\theta} - \text{error}, \hat{\theta} + \text{error}).$$

And we can ask questions about how small is the *error* or how much sure are we in our Estimate (for the Unknown Parameter), and how large n needs to be to have a good estimate.

¹Recall the \widehat{SE} , the Estimated Standard Error reporting story.

Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using $\hat{\theta}$, if $\hat{\theta}$ possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for θ^* .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small¹, or, rather, as

$$\theta^* \in (\hat{\theta} - \text{error}, \hat{\theta} + \text{error}).$$

And we can ask questions about how small is the *error* or how much sure are we in our Estimate (for the Unknown Parameter), and how large n needs to be to have a good estimate.

Here we want to develop the theory of Confidence Intervals, which will contain answers to these questions.

¹Recall the \widehat{SE} , the Estimated Standard Error reporting story.

CI Problem Setting

Assume we have a Random Sample from a Parametric Model \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

CI Problem Setting

Assume we have a Random Sample from a Parametric Model \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

CI Problem Setting

Assume we have a Random Sample from a Parametric Model \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

More rigorously, we want to have an interval

- ▶ which contains the True Parameter value with high Probability;

CI Problem Setting

Assume we have a Random Sample from a Parametric Model \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

More rigorously, we want to have an interval

- ▶ which contains the True Parameter value with high Probability;
- ▶ which has the possible smallest length.

CI Problem Setting

Assume we have a Random Sample from a Parametric Model \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

More rigorously, we want to have an interval

- ▶ which contains the True Parameter value with high Probability;
- ▶ which has the possible smallest length.

Let us state this in Mathematical terms. We will consider here only 1D case, i.e., we will assume $\theta \in \Theta \subset \mathbb{R}$.

Random Intervals and CI

We start by **Random Intervals**:

Random Intervals and CI

We start by **Random Intervals**:

Assume X and Y are two r.v. on the same probability space and $X \leq Y$ a.s., that is, $\mathbb{P}(X > Y) = 0$.

Random Intervals and CI

We start by **Random Intervals**:

Assume X and Y are two r.v. on the same probability space and $X \leq Y$ a.s., that is, $\mathbb{P}(X > Y) = 0$. We will call intervals of the form $[X, Y]$ or (X, Y) or $(X, Y]$ or $[X, Y)$ or $(X, +\infty), \dots$, **random intervals**.

Random Intervals and CI

We start by **Random Intervals**:

Assume X and Y are two r.v. on the same probability space and $X \leq Y$ a.s., that is, $\mathbb{P}(X > Y) = 0$. We will call intervals of the form $[X, Y]$ or (X, Y) or $(X, Y]$ or $[X, Y)$ or $(X, +\infty), \dots$, **random intervals**.

Example: Assume $X \sim \text{Pois}(2.3)$. Then

$$[X, X + 2)$$

is a Random Interval.

Random Intervals and CI

We start by **Random Intervals**:

Assume X and Y are two r.v. on the same probability space and $X \leq Y$ a.s., that is, $\mathbb{P}(X > Y) = 0$. We will call intervals of the form $[X, Y]$ or (X, Y) or $(X, Y]$ or $[X, Y)$ or $(X, +\infty), \dots$, **random intervals**.

Example: Assume $X \sim \text{Pois}(2.3)$. Then

$$[X, X + 2)$$

is a Random Interval.

Example: Let X_1, X_2, \dots, X_n are IID r.v.s. Then

$$(\bar{X} - 0.1, \bar{X} + 0.1)$$

is a Random Interval.