

CS 107, Probability, Spring 2019

Lecture 22

Michael Poghosyan

AUA

15 March 2019

- Examples of Important Discrete R.V.s

$$0 = 1$$

$$0 = 1$$

Proof:

$$\int \frac{1}{x} dx =$$

$$0 = 1$$

Proof:

$$\int \frac{1}{x} dx = (IBP)$$

$$0 = 1$$

Proof:

$$\int \frac{1}{x} dx = (IBP) = x \cdot \frac{1}{x} - \int x \cdot \left(\frac{1}{x}\right)' dx =$$
$$1 + \int \frac{1}{x} dx$$

And we cancel $\int \frac{1}{x} dx$ from the LHS and RHS.

Reminder on RVs

We know that we can describe a r.v. through:

Reminder on RVs

We know that we can describe a r.v. through:

- Its CDF (general case, both Discrete, Continuous, and even Singular r.v.s)

Reminder on RVs

We know that we can describe a r.v. through:

- Its CDF (general case, both Discrete, Continuous, and even Singular r.v.s)
- Its PMF (if our r.v. is Discrete) or PDF (if our r.v. is Continuous)

Reminder on RVs

We know that we can describe a r.v. through:

- Its CDF (general case, both Discrete, Continuous, and even Singular r.v.s)
- Its PMF (if our r.v. is Discrete) or PDF (if our r.v. is Continuous)

That is, we can calculate Probabilities about the possible values of a r.v. through the CDF or PMF/PDF.

Probabilities through the CDF

If X is any r.v., then the CDF of X is defined by

$$F(x) =$$

Probabilities through the CDF

If X is any r.v., then the CDF of X is defined by

$$F(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R}.$$

Probabilities through the CDF

If X is any r.v., then the CDF of X is defined by

$$F(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R}.$$

And using the CDF, we can calculate probabilities like

$$\mathbb{P}(X = a) =$$

Probabilities through the CDF

If X is any r.v., then the CDF of X is defined by

$$F(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R}.$$

And using the CDF, we can calculate probabilities like

$$\mathbb{P}(X = a) = F(a) - F(a-),$$

$$\mathbb{P}(a < X \leq b) =$$

Probabilities through the CDF

If X is any r.v., then the CDF of X is defined by

$$F(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R}.$$

And using the CDF, we can calculate probabilities like

$$\mathbb{P}(X = a) = F(a) - F(a-),$$

$$\mathbb{P}(a < X \leq b) = F(b) - F(a).$$

Probabilities through the PM(D)F

Let's consider now two cases - when X is Discrete or Continuous:

Discrete

Continuous

Probabilities through the PM(D)F

Let's consider now two cases - when X is Discrete or Continuous:

Discrete	Continuous
PMF: $\mathbb{P}(X = x_k) = F(x_k) - F(x_{k-})$	PDF: $f(x) = F'(x)$

Probabilities through the PM(D)F

Let's consider now two cases - when X is Discrete or Continuous:

Discrete	Continuous
PMF: $\mathbb{P}(X = x_k) = F(x_k) - F(x_{k-})$	PDF: $f(x) = F'(x)$
$\mathbb{P}(X = x_k) = p_k$	$\mathbb{P}(X = x) = 0, \forall x \in \mathbb{R}$

Probabilities through the PM(D)F

Let's consider now two cases - when X is Discrete or Continuous:

Discrete	Continuous
PMF: $\mathbb{P}(X = x_k) = F(x_k) - F(x_k -)$	PDF: $f(x) = F'(x)$
$\mathbb{P}(X = x_k) = p_k$	$\mathbb{P}(X = x) = 0, \forall x \in \mathbb{R}$
$\sum_k p_k = 1$	$\int_{-\infty}^{+\infty} f(x) dx = 1$

Probabilities through the PM(D)F

Let's consider now two cases - when X is Discrete or Continuous:

Discrete	Continuous
PMF: $\mathbb{P}(X = x_k) = F(x_k) - F(x_k -)$	PDF: $f(x) = F'(x)$
$\mathbb{P}(X = x_k) = p_k$	$\mathbb{P}(X = x) = 0, \forall x \in \mathbb{R}$
$\sum_k p_k = 1$	$\int_{-\infty}^{+\infty} f(x) dx = 1$
$F(x) = \sum_{x_k \leq x} p_k$	$F(x) = \int_{-\infty}^x f(t) dt$

Probabilities through the PM(D)F

Let's consider now two cases - when X is Discrete or Continuous:

Discrete

$$\text{PMF: } \mathbb{P}(X = x_k) = F(x_k) - F(x_{k-1})$$

$$\mathbb{P}(X = x_k) = p_k$$

$$\sum_k p_k = 1$$

$$F(x) = \sum_{x_k \leq x} p_k$$

$$\mathbb{P}(a \leq X \leq b) = \sum_{a \leq x_k \leq b} p_k$$

Continuous

$$\text{PDF: } f(x) = F'(x)$$

$$\mathbb{P}(X = x) = 0, \forall x \in \mathbb{R}$$

$$\int_{-\infty}^{+\infty} f(x) dx = 1$$

$$F(x) = \int_{-\infty}^x f(t) dt$$

$$\mathbb{P}(a \leq X \leq b) = \int_a^b f(x) dx$$

Warnings!

Assume X is a r.v., and a is a number.

Warnings!

Assume X is a r.v., and a is a number. Then X **is not an Event**.

Warnings!

Assume X is a r.v., and a is a number. Then X **is not an Event**. Instead, the sets $\{X = a\}$, $\{X < a\}$, $\{X \leq a\}$, ... are Events. So:

Warnings!

Assume X is a r.v., and a is a number. Then X **is not an Event**. Instead, the sets $\{X = a\}$, $\{X < a\}$, $\{X \leq a\}, \dots$ are Events. So:

- Writing $\mathbb{P}(X)$ is **incorrect**;

Warnings!

Assume X is a r.v., and a is a number. Then X **is not an Event**. Instead, the sets $\{X = a\}$, $\{X < a\}$, $\{X \leq a\}$, ... are Events. So:

- Writing $\mathbb{P}(X)$ is **incorrect**;
- Writing $\mathbb{P}(a)$ is **incorrect**;

Warnings!

Assume X is a r.v., and a is a number. Then X **is not an Event**. Instead, the sets $\{X = a\}$, $\{X < a\}$, $\{X \leq a\}$, ... are Events. So:

- Writing $\mathbb{P}(X)$ is **incorrect**;
- Writing $\mathbb{P}(a)$ is **incorrect**;
- Writing $\mathbb{P}(X = a)$ or $\mathbb{P}(X < a)$ or $\mathbb{P}(X \geq a)$, ... are **correct**.

Warnings!

Assume X is a r.v., and a is a number. Then X **is not an Event**. Instead, the sets $\{X = a\}$, $\{X < a\}$, $\{X \leq a\}$, ... are Events. So:

- Writing $\mathbb{P}(X)$ is **incorrect**;
- Writing $\mathbb{P}(a)$ is **incorrect**;
- Writing $\mathbb{P}(X = a)$ or $\mathbb{P}(X < a)$ or $\mathbb{P}(X \geq a)$, ... are **correct**.

Example: Let X be the number of emails I will get today. Then X is a r.v.

Warnings!

Assume X is a r.v., and a is a number. Then X **is not an Event**. Instead, the sets $\{X = a\}$, $\{X < a\}$, $\{X \leq a\}$, ... are Events. So:

- Writing $\mathbb{P}(X)$ is **incorrect**;
- Writing $\mathbb{P}(a)$ is **incorrect**;
- Writing $\mathbb{P}(X = a)$ or $\mathbb{P}(X < a)$ or $\mathbb{P}(X \geq a)$, ... are **correct**.

Example: Let X be the number of emails I will get today. Then X is a r.v. Is it Discrete or Continuous?

Warnings!

Assume X is a r.v., and a is a number. Then X **is not an Event**. Instead, the sets $\{X = a\}$, $\{X < a\}$, $\{X \leq a\}$, ... are Events. So:

- Writing $\mathbb{P}(X)$ is **incorrect**;
- Writing $\mathbb{P}(a)$ is **incorrect**;
- Writing $\mathbb{P}(X = a)$ or $\mathbb{P}(X < a)$ or $\mathbb{P}(X \geq a)$, ... are **correct**.

Example: Let X be the number of emails I will get today. Then X is a r.v. Is it Discrete or Continuous? Then $\mathbb{P}(X)$ denotes "the Probability of number of emails I will get today". Nonsense, of course 😊

Warnings!

Assume X is a r.v., and a is a number. Then X **is not an Event**. Instead, the sets $\{X = a\}$, $\{X < a\}$, $\{X \leq a\}$, ... are Events. So:

- Writing $\mathbb{P}(X)$ is **incorrect**;
- Writing $\mathbb{P}(a)$ is **incorrect**;
- Writing $\mathbb{P}(X = a)$ or $\mathbb{P}(X < a)$ or $\mathbb{P}(X \geq a)$, ... are **correct**.

Example: Let X be the number of emails I will get today. Then X is a r.v. Is it Discrete or Continuous? Then $\mathbb{P}(X)$ denotes "the Probability of number of emails I will get today". Nonsense, of course 😊 But we can talk about $\mathbb{P}(X = 0)$ or $\mathbb{P}(X > 10)$, and they have clear meanings.

Important Discrete R.V.

Discrete Uniform Distribution

Discrete Uniform Distribution

We will say that the r.v. X has a Discrete Uniform Distribution with (over) the values x_1, x_2, \dots, x_n ($x_i \neq x_j$, $i \neq j$), and we will write $X \sim \text{DiscreteUnif}(x_1, \dots, x_n)$, if

Values of X	x_1	x_2	\dots	x_n
$\mathbb{P}(X = x)$	$\frac{1}{n}$	$\frac{1}{n}$	\dots	$\frac{1}{n}$

Discrete Uniform Distribution

Discrete Uniform Distribution

We will say that the r.v. X has a Discrete Uniform Distribution with (over) the values x_1, x_2, \dots, x_n ($x_i \neq x_j$, $i \neq j$), and we will write $X \sim \text{DiscreteUnif}(x_1, \dots, x_n)$, if

Values of X	x_1	x_2	\dots	x_n
$\mathbb{P}(X = x)$	$\frac{1}{n}$	$\frac{1}{n}$	\dots	$\frac{1}{n}$

that is, $\mathbb{P}(X = x_k) = \frac{1}{n}$, for any $k = 1, 2, \dots, n$.

Discrete Uniform Distribution: Examples

Discrete Uniform Distribution models equiprobable outcomes case. Some examples:

Discrete Uniform Distribution: Examples

Discrete Uniform Distribution models equiprobable outcomes case. Some examples:

- We are tossing a fair coin, and $X = 0$ if Heads appears, and $X = 1$ in the other case.

Discrete Uniform Distribution: Examples

Discrete Uniform Distribution models equiprobable outcomes case. Some examples:

- We are tossing a fair coin, and $X = 0$ if Heads appears, and $X = 1$ in the other case.
- We are rolling a fair die, X is the top face number.

Discrete Uniform Distribution: Examples

Discrete Uniform Distribution models equiprobable outcomes case. Some examples:

- We are tossing a fair coin, and $X = 0$ if Heads appears, and $X = 1$ in the other case.
- We are rolling a fair die, X is the top face number.
- Can you give some more?

Discrete Uniform Distribution: R Examples

R Code

```
#Discrete Uniform on data
data <- c(-1,2,4)
sample(data, size = 10, replace = T)

#Generating a sample of size 100
s <- sample(data, size = 100, replace = T)
#Calculating the number of -1 in the sample s
length(s[s == -1])
```