

# CS 107, Probability, Spring 2019

## Lecture 43

Michael Poghosyan

AUA

- 3 lectures

Ta-da-da-daaaam! Quiz time!

- The Central Limit Theorem
- Expectation and Variance for Important Distributions
- Intro to Markov Chains

# The Central Limit Theorem

CLT gives more info about the Distribution of  $S_n$  and  $\bar{X}_n$ :

## The Central Limit Theorem, CLT

Assume  $X_1, X_2, \dots, X_n$  are IID with finite Expectation  $\mu = \mathbb{E}(X_1)$  and Variance  $\sigma^2 = \text{Var}(X_1)$ . We Standardize  $S_n$  (or  $\bar{X}_n$ ):

$$Z_n = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}} = \frac{S_n - n \cdot \mu}{\sqrt{n} \cdot \sigma}$$

$$\left( Z_n = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}} = \frac{\bar{X}_n - \mu}{\sigma / \sqrt{n}} \right).$$

Then, for any subset  $A \subset \mathbb{R}$ ,

$$\mathbb{P}(Z_n \in A) \rightarrow \int_A \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx.$$

# Example:

**Example:** Assume I have a Piggy Bank.

# Example:

**Example:** Assume I have a Piggy Bank.



# Example:

**Example:** Assume I have a Piggy Bank.  
I am collecting coins worth 500 AMD in my Piggy.

# Example:

**Example:** Assume I have a Piggy Bank.

I am collecting coins worth 500 AMD in my Piggy. The mean weight of a 500 AMD coin is 5 gr with Standard Deviation 0.1 gr.



# Example:

**Example:** Assume I have a Piggy Bank.

I am collecting coins worth 500 AMD in my Piggy. The mean weight of a 500 AMD coin is 5 gr with Standard Deviation 0.1 gr. I have collected 250 coins.

# Example:

**Example:** Assume I have a Piggy Bank.

I am collecting coins worth 500 AMD in my Piggy. The mean weight of a 500 AMD coin is 5 gr with Standard Deviation 0.1 gr. I have collected 250 coins. Assume  $W$  is the weight, in grams, of all that 250 coins.

# Example:

**Example:** Assume I have a Piggy Bank.

I am collecting coins worth 500 AMD in my Piggy. The mean weight of a 500 AMD coin is 5 gr with Standard Deviation 0.1 gr. I have collected 250 coins. Assume  $W$  is the weight, in grams, of all that 250 coins. Clearly, the average weight of all coins is  $250 \cdot 5gr = 1250gr$ , i.e.,  $\mathbb{E}(W) = 1250gr$ .

# Example:

**Example:** Assume I have a Piggy Bank.

I am collecting coins worth 500 AMD in my Piggy. The mean weight of a 500 AMD coin is 5 gr with Standard Deviation 0.1 gr. I have collected 250 coins. Assume  $W$  is the weight, in grams, of all that 250 coins. Clearly, the average weight of all coins is  $250 \cdot 5gr = 1250gr$ , i.e.,  $\mathbb{E}(W) = 1250gr$ .

- Calculate, approximately,

$$\mathbb{P}(W > 1200);$$

# Example:

**Example:** Assume I have a Piggy Bank.

I am collecting coins worth 500 AMD in my Piggy. The mean weight of a 500 AMD coin is 5 gr with Standard Deviation 0.1 gr. I have collected 250 coins. Assume  $W$  is the weight, in grams, of all that 250 coins. Clearly, the average weight of all coins is  $250 \cdot 5gr = 1250gr$ , i.e.,  $\mathbb{E}(W) = 1250gr$ .

- Calculate, approximately,

$$\mathbb{P}(W > 1200);$$

- Calculate, approximately,

$$\mathbb{P}(1100 \leq W < 1300);$$

# Example:

**Example:** Assume I have a Piggy Bank.

I am collecting coins worth 500 AMD in my Piggy. The mean weight of a 500 AMD coin is 5 gr with Standard Deviation 0.1 gr. I have collected 250 coins. Assume  $W$  is the weight, in grams, of all that 250 coins. Clearly, the average weight of all coins is  $250 \cdot 5gr = 1250gr$ , i.e.,  $\mathbb{E}(W) = 1250gr$ .

- Calculate, approximately,

$$\mathbb{P}(W > 1200);$$

- Calculate, approximately,

$$\mathbb{P}(1100 \leq W < 1300);$$

- Guess what I could do with the collected amount 😊

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) =$

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) =$



# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$
- If  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{E}(X) =$

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$
- If  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{E}(X) = np$ ,  $\text{Var}(X) =$

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$
- If  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{E}(X) = np$ ,  $\text{Var}(X) = np(1 - p)$

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$
- If  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{E}(X) = np$ ,  $\text{Var}(X) = np(1 - p)$   
Idea:  $X = X_1 + X_2 + \dots + X_n$ , where  $X_k \sim \text{Bernoulli}(p)$  are IID

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$
- If  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{E}(X) = np$ ,  $\text{Var}(X) = np(1 - p)$   
Idea:  $X = X_1 + X_2 + \dots + X_n$ , where  $X_k \sim \text{Bernoulli}(p)$  are IID
- If  $X \sim \text{Geom}(p)$ , then  $\mathbb{E}(X) =$

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$
- If  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{E}(X) = np$ ,  $\text{Var}(X) = np(1 - p)$   
Idea:  $X = X_1 + X_2 + \dots + X_n$ , where  $X_k \sim \text{Bernoulli}(p)$  are IID
- If  $X \sim \text{Geom}(p)$ , then  $\mathbb{E}(X) = \frac{1}{p}$ ,  $\text{Var}(X) = \frac{1 - p}{p^2}$

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$
- If  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{E}(X) = np$ ,  $\text{Var}(X) = np(1 - p)$   
Idea:  $X = X_1 + X_2 + \dots + X_n$ , where  $X_k \sim \text{Bernoulli}(p)$  are IID
- If  $X \sim \text{Geom}(p)$ , then  $\mathbb{E}(X) = \frac{1}{p}$ ,  $\text{Var}(X) = \frac{1 - p}{p^2}$
- If  $X \sim \text{Poisson}(\lambda)$ , then  $\mathbb{E}(X) =$



# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$
- If  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{E}(X) = np$ ,  $\text{Var}(X) = np(1 - p)$   
Idea:  $X = X_1 + X_2 + \dots + X_n$ , where  $X_k \sim \text{Bernoulli}(p)$  are IID
- If  $X \sim \text{Geom}(p)$ , then  $\mathbb{E}(X) = \frac{1}{p}$ ,  $\text{Var}(X) = \frac{1 - p}{p^2}$
- If  $X \sim \text{Poisson}(\lambda)$ , then  $\mathbb{E}(X) = \lambda$ ,  $\text{Var}(X) =$

# Expectations and Variances for Important RVs

## Discrete Distributions

- If  $X \sim \text{Bernoulli}(p)$ , then  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$
- If  $X \sim \text{Binom}(n, p)$ , then  $\mathbb{E}(X) = np$ ,  $\text{Var}(X) = np(1 - p)$   
Idea:  $X = X_1 + X_2 + \dots + X_n$ , where  $X_k \sim \text{Bernoulli}(p)$  are IID
- If  $X \sim \text{Geom}(p)$ , then  $\mathbb{E}(X) = \frac{1}{p}$ ,  $\text{Var}(X) = \frac{1 - p}{p^2}$
- If  $X \sim \text{Poisson}(\lambda)$ , then  $\mathbb{E}(X) = \lambda$ ,  $\text{Var}(X) = \lambda$

# Expectations and Variances for Important RVs

## Continuous Distributions

- If  $X \sim \text{Unif}[a, b]$ , then  $\mathbb{E}(X) =$

# Expectations and Variances for Important RVs

## Continuous Distributions

- If  $X \sim \text{Unif}[a, b]$ , then  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) =$

# Expectations and Variances for Important RVs

## Continuous Distributions

- If  $X \sim \text{Unif}[a, b]$ , then  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ ;

# Expectations and Variances for Important RVs

## Continuous Distributions

- If  $X \sim \text{Unif}[a, b]$ , then  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ ;
- If  $X \sim \text{Exp}(\lambda)$ , then  $\mathbb{E}(X) =$

# Expectations and Variances for Important RVs

## Continuous Distributions

- If  $X \sim \text{Unif}[a, b]$ , then  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ ;
- If  $X \sim \text{Exp}(\lambda)$ , then  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ ;

# Expectations and Variances for Important RVs

## Continuous Distributions

- If  $X \sim \text{Unif}[a, b]$ , then  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ ;
- If  $X \sim \text{Exp}(\lambda)$ , then  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ ;
- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\mathbb{E}(X) =$



# Expectations and Variances for Important RVs

## Continuous Distributions

- If  $X \sim \text{Unif}[a, b]$ , then  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ ;
- If  $X \sim \text{Exp}(\lambda)$ , then  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ ;
- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\mathbb{E}(X) = \mu$ ,  $\text{Var}(X) =$

# Expectations and Variances for Important RVs

## Continuous Distributions

- If  $X \sim \text{Unif}[a, b]$ , then  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ ;
- If  $X \sim \text{Exp}(\lambda)$ , then  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ ;
- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\mathbb{E}(X) = \mu$ ,  $\text{Var}(X) = \sigma^2$ ;

# Expectations and Variances for Important RVs

## Continuous Distributions

- If  $X \sim \text{Unif}[a, b]$ , then  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ ;
- If  $X \sim \text{Exp}(\lambda)$ , then  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ ;
- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\mathbb{E}(X) = \mu$ ,  $\text{Var}(X) = \sigma^2$ ;
- If  $\mathbf{X} \sim \mathcal{N}(\mu, \Sigma)$ , then  $\mathbb{E}(\mathbf{X}) = \mu$ ,  $\text{Cov}(\mathbf{X}) = \Sigma$

**Note:**  $\text{Cov}(\mathbf{X})$  is the Covariance Matrix of  $\mathbf{X}$ .

# Intro to Markov Chains

# Intro To Markov Chains

Up to this point we have considered sequences of r.v.  $X_k$  that are IID.

# Intro To Markov Chains

Up to this point we have considered sequences of r.v.  $X_k$  that are IID. This kind of sequences appear naturally in many phenomena, and the IID structure simplifies calculations, and gives us the celebrated LLN and CLT.

# Intro To Markov Chains

Up to this point we have considered sequences of r.v.  $X_k$  that are IID. This kind of sequences appear naturally in many phenomena, and the IID structure simplifies calculations, and gives us the celebrated LLN and CLT.

But in many other real-life phenomena, we cannot model things as Independent r.v.s. Say, in Stock Markets, if  $S_n$  is the daily closing price of some stock, then  $S_0, S_1, S_2, \dots$  are not Independent.

# Intro To Markov Chains

Up to this point we have considered sequences of r.v.  $X_k$  that are IID. This kind of sequences appear naturally in many phenomena, and the IID structure simplifies calculations, and gives us the celebrated LLN and CLT.

But in many other real-life phenomena, we cannot model things as Independent r.v.s. Say, in Stock Markets, if  $S_n$  is the daily closing price of some stock, then  $S_0, S_1, S_2, \dots$  are not Independent. Or, if  $W_1, W_2, W_3, \dots$  are daily weather condition r.v.s (say,  $W_i = 1$ , if the  $i$ -th day was Sunny,  $W_i = 0$ , if Rainy), then  $W_k$ -s are not Independent.



# Intro To Markov Chains

Up to this point we have considered sequences of r.v.  $X_k$  that are IID. This kind of sequences appear naturally in many phenomena, and the IID structure simplifies calculations, and gives us the celebrated LLN and CLT.

But in many other real-life phenomena, we cannot model things as Independent r.v.s. Say, in Stock Markets, if  $S_n$  is the daily closing price of some stock, then  $S_0, S_1, S_2, \dots$  are not Independent. Or, if  $W_1, W_2, W_3, \dots$  are daily weather condition r.v.s (say,  $W_i = 1$ , if the  $i$ -th day was Sunny,  $W_i = 0$ , if Rainy), then  $W_k$ -s are not Independent. So we want to be able to model Dependent Data sequences, and one of the simplest and very important Models is the Markov Chain Model.

# Intro To Markov Chains

Up to this point we have considered sequences of r.v.  $X_k$  that are IID. This kind of sequences appear naturally in many phenomena, and the IID structure simplifies calculations, and gives us the celebrated LLN and CLT.

But in many other real-life phenomena, we cannot model things as Independent r.v.s. Say, in Stock Markets, if  $S_n$  is the daily closing price of some stock, then  $S_0, S_1, S_2, \dots$  are not Independent. Or, if  $W_1, W_2, W_3, \dots$  are daily weather condition r.v.s (say,  $W_i = 1$ , if the  $i$ -th day was Sunny,  $W_i = 0$ , if Rainy), then  $W_k$ -s are not Independent. So we want to be able to model Dependent Data sequences, and one of the simplest and very important Models is the Markov Chain Model.

Recall also that we have considered a Markov Chain - when talking about the Language Modeling!

# Intro To Markov Chains

# Intro To Markov Chains

Here we will consider Stochastic Processes  $X_t : t \in T$ , where  $X_t$  is a r.v. for each  $t$ .

# Intro To Markov Chains

Here we will consider Stochastic Processes  $X_t : t \in T$ , where  $X_t$  is a r.v. for each  $t$ . Usually, we refer to  $t$  as time, and distinguish between

# Intro To Markov Chains

Here we will consider Stochastic Processes  $X_t : t \in T$ , where  $X_t$  is a r.v. for each  $t$ . Usually, we refer to  $t$  as time, and distinguish between

- **Discrete Time Stochastic Process:** here  $T = \{0, 1, 2, \dots\}$ , so  $t$  can take values  $0, 1, 2, \dots$ , and we will have just a sequence of r.v.s

$$X_0, X_1, X_2, \dots$$

# Intro To Markov Chains

Here we will consider Stochastic Processes  $X_t : t \in T$ , where  $X_t$  is a r.v. for each  $t$ . Usually, we refer to  $t$  as time, and distinguish between

- **Discrete Time Stochastic Process:** here  $T = \{0, 1, 2, \dots\}$ , so  $t$  can take values  $0, 1, 2, \dots$ , and we will have just a sequence of r.v.s

$$X_0, X_1, X_2, \dots$$

- **Continuous Time Stochastic Process:** here  $T = [0, +\infty)$  or  $T = [0, A]$ , so  $t$  can take any real values from the specified interval. Say, we can have  $X_1, X_{1.3}, X_{\sqrt{2}}$  etc.

# Intro To Markov Chains

Here we will consider Stochastic Processes  $X_t : t \in T$ , where  $X_t$  is a r.v. for each  $t$ . Usually, we refer to  $t$  as time, and distinguish between

- **Discrete Time Stochastic Process:** here  $T = \{0, 1, 2, \dots\}$ , so  $t$  can take values 0, 1, 2, ..., and we will have just a sequence of r.v.s

$$X_0, X_1, X_2, \dots$$

- **Continuous Time Stochastic Process:** here  $T = [0, +\infty)$  or  $T = [0, A]$ , so  $t$  can take any real values from the specified interval. Say, we can have  $X_1, X_{1.3}, X_{\sqrt{2}}$  etc.

We will consider only Discrete Time Markov Chains, which is an example of Discrete Time Stochastic Processes.



# Definition of the Markov Chain: State Space

First, we fix a set  $S =$  the State Space.

# Definition of the Markov Chain: State Space

First, we fix a set  $S =$  the State Space.  $S$  is the set of possible values of  $X_k$ .

# Definition of the Markov Chain: State Space

First, we fix a set  $S =$  the State Space.  $S$  is the set of possible values of  $X_k$ . We will assume here again that  $S$  is Finite: in this case we talk about **Finite State Discrete Time Markov Chains**.

# Definition of the Markov Chain: State Space

First, we fix a set  $S =$  the State Space.  $S$  is the set of possible values of  $X_k$ . We will assume here again that  $S$  is Finite: in this case we talk about **Finite State Discrete Time Markov Chains**. For simplicity, we can take  $S = \{1, 2, 3, \dots, N\}$  (we just enumerate the States, we can start also from 0).

# Definition of the Markov Chain: State Space

First, we fix a set  $S =$  the State Space.  $S$  is the set of possible values of  $X_k$ . We will assume here again that  $S$  is Finite: in this case we talk about **Finite State Discrete Time Markov Chains**. For simplicity, we can take  $S = \{1, 2, 3, \dots, N\}$  (we just enumerate the States, we can start also from 0). Assume now we are modeling a (Probabilistic, Stochastic) System, which can be in any of states from  $S$ .

# Definition of the Markov Chain: State Space

First, we fix a set  $S =$  the State Space.  $S$  is the set of possible values of  $X_k$ . We will assume here again that  $S$  is Finite: in this case we talk about **Finite State Discrete Time Markov Chains**. For simplicity, we can take  $S = \{1, 2, 3, \dots, N\}$  (we just enumerate the States, we can start also from 0).

Assume now we are modeling a (Probabilistic, Stochastic) System, which can be in any of states from  $S$ . We denote the state of our System at time  $t$  by  $X_t$ ,  $t = 0, 1, 2, \dots$ . So  $X_t$  can take values from  $S$ , and  $X_t$  is a r.v., since we do not know in which State will be our  $X_t$ .

# Definition of the Markov Chain: State Space

First, we fix a set  $S =$  the State Space.  $S$  is the set of possible values of  $X_k$ . We will assume here again that  $S$  is Finite: in this case we talk about **Finite State Discrete Time Markov Chains**. For simplicity, we can take  $S = \{1, 2, 3, \dots, N\}$  (we just enumerate the States, we can start also from 0).

Assume now we are modeling a (Probabilistic, Stochastic) System, which can be in any of states from  $S$ . We denote the state of our System at time  $t$  by  $X_t$ ,  $t = 0, 1, 2, \dots$ . So  $X_t$  can take values from  $S$ , and  $X_t$  is a r.v., since we do not know in which State will be our  $X_t$ .

**Example:** Say, we are modeling the Weather. The States are *Rainy, Sunny*, or, say, 0, 1.  $W_t$  is the weather at the  $t$ -th day, started from today ( $t = 0$ ). So  $W_0$  is today's weather,  $W_1$  is the tomorrow's weather etc.

# Definition of the Markov Chain

Now, we give the Definition:



# Definition of the Markov Chain

Now, we give the Definition:

## Markov Chain

Assume  $X_0, X_1, X_2, \dots$  is a sequence of r.v.s (Discrete Stochastic Process), which take values from  $S = \{1, 2, \dots, N\}$ . We say that  $X_n, n = 0, 1, 2, \dots$  is a (Finite State Discrete Time) **Markov Chain**, if

$$\mathbb{P}(X_{t+1} = j | X_t = i, X_{t-1} = k, \dots, X_0 = m) = \mathbb{P}(X_{t+1} = j | X_t = i)$$

for any time  $t$ , for any state  $j, i, k, \dots, m$ .

# Definition of the Markov Chain

Now, we give the Definition:

## Markov Chain

Assume  $X_0, X_1, X_2, \dots$  is a sequence of r.v.s (Discrete Stochastic Process), which take values from  $S = \{1, 2, \dots, N\}$ . We say that  $X_n, n = 0, 1, 2, \dots$  is a (Finite State Discrete Time) **Markov Chain**, if

$$\mathbb{P}(X_{t+1} = j | X_t = i, X_{t-1} = k, \dots, X_0 = m) = \mathbb{P}(X_{t+1} = j | X_t = i)$$

for any time  $t$ , for any state  $j, i, k, \dots, m$ .

**Interpretation:** Given today's State, tomorrow's State is independent of the past States.

# Definition of the Markov Chain

Now, we give the Definition:

## Markov Chain

Assume  $X_0, X_1, X_2, \dots$  is a sequence of r.v.s (Discrete Stochastic Process), which take values from  $S = \{1, 2, \dots, N\}$ . We say that  $X_n, n = 0, 1, 2, \dots$  is a (Finite State Discrete Time) **Markov Chain**, if

$$\mathbb{P}(X_{t+1} = j | X_t = i, X_{t-1} = k, \dots, X_0 = m) = \mathbb{P}(X_{t+1} = j | X_t = i)$$

for any time  $t$ , for any state  $j, i, k, \dots, m$ .

**Interpretation:** Given today's State, tomorrow's State is independent of the past States. Or, in other words, today's information is enough to completely determine the probabilities of Tomorrow's states.