

CS 107, Probability, Spring 2019

Lecture 07

Michael Poghosyan

AUA

1 February 2019

- Classical Probability Models: Finite Sample Spaces with Equally Likely Outcomes = Combinatorial Problems, Cont'd

Strange Mailman Problem

Mailman needs to deliver 100 letters to given addresses. But he is putting letters at random to the mailboxes. What is the probability that at least one person will receive the correct letter?

Strange Mailman Problem

Mailman needs to deliver 100 letters to given addresses. But he is putting letters at random to the mailboxes. What is the probability that at least one person will receive the correct letter?

Answer: the Probability is about 63%.

Example:

- **Problem:** We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?
-

Example:

- **Problem:** We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?
-
- Clear assumption: Choosing each ball is equiprobable

Example:

- **Problem:** We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?
-
- Clear assumption: Choosing each ball is equiprobable
 - **Case 1:** We take balls without a replacement:

Example:

- **Problem:** We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?
-

- Clear assumption: Choosing each ball is equiprobable
- **Case 1:** We take balls without a replacement:
- **Case 2:** We take balls with a replacement:

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Our Experiment is taking 12 balls at random.

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Our Experiment is taking 12 balls at random. Let us fix, as an Outcome of our Experiment, the number of white and black balls in the chosen set of 12 balls.

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Our Experiment is taking 12 balls at random. Let us fix, as an Outcome of our Experiment, the number of white and black balls in the chosen set of 12 balls. Say, if we will get 3 whites and 9 blacks, our Outcome will be $(3, 9)$, or if we will have 5 whites and 7 blacks, then our Outcome will be $(5, 7)$.

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

So our Sample Space is:

$$\Omega = \{(n, 12 - n) : n = 0, 1, \dots, 12\}, \quad \#\Omega =$$

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

So our Sample Space is:

$$\Omega = \{(n, 12 - n) : n = 0, 1, \dots, 12\}, \quad \#\Omega = 13$$

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

So our Sample Space is:

$$\Omega = \{(n, 12 - n) : n = 0, 1, \dots, 12\}, \quad \#\Omega = 13$$

The Event of our interest is:

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

So our Sample Space is:

$$\Omega = \{(n, 12 - n) : n = 0, 1, \dots, 12\}, \quad \#\Omega = 13$$

The Event of our interest is:

$$A = \{(4, 8)\}, \quad \#A = 1.$$

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

So our Sample Space is:

$$\Omega = \{(n, 12 - n) : n = 0, 1, \dots, 12\}, \quad \#\Omega = 13$$

The Event of our interest is:

$$A = \{(4, 8)\}, \quad \#A = 1.$$

So the Probability of A is:

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

So our Sample Space is:

$$\Omega = \{(n, 12 - n) : n = 0, 1, \dots, 12\}, \quad \#\Omega = 13$$

The Event of our interest is:

$$A = \{(4, 8)\}, \quad \#A = 1.$$

So the Probability of A is:

$$\mathbb{P}(A) = \frac{1}{13}.$$

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

So our Sample Space is:

$$\Omega = \{(n, 12 - n) : n = 0, 1, \dots, 12\}, \quad \#\Omega = 13$$

The Event of our interest is:

$$A = \{(4, 8)\}, \quad \#A = 1.$$

So the Probability of A is:

$$\mathbb{P}(A) = \frac{1}{13}.$$

Of course, not! The solution is incorrect :P

Without Replacements Model, Solution 1:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

So our Sample Space is:

$$\Omega = \{(n, 12 - n) : n = 0, 1, \dots, 12\}, \quad \#\Omega = 13$$

The Event of our interest is:

$$A = \{(4, 8)\}, \quad \#A = 1.$$

So the Probability of A is:

$$\mathbb{P}(A) = \frac{1}{13}.$$

Of course, not! The solution is incorrect :P Why?

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Our Experiment is taking 12 balls at random.

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Our Experiment is taking 12 balls at random. Let us record, as an Outcome of our Experiment, the color of each of 12 balls.

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Our Experiment is taking 12 balls at random. Let us record, as an Outcome of our Experiment, the color of each of 12 balls. Say, one of the outcomes can be

$$(w, b, w, w, b, w, w, w, w, w, b, b).$$

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Our Experiment is taking 12 balls at random. Let us record, as an Outcome of our Experiment, the color of each of 12 balls. Say, one of the outcomes can be

$$(w, b, w, w, b, w, w, w, w, w, b, b).$$

In this case, the Sample Space is:

$$\Omega =$$

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Our Experiment is taking 12 balls at random. Let us record, as an Outcome of our Experiment, the color of each of 12 balls. Say, one of the outcomes can be

$$(w, b, w, w, b, w, w, w, w, w, b, b).$$

In this case, the Sample Space is:

$$\Omega = \{(x_1, x_2, \dots, x_{12}) : x_k \in \{w, b\}\}, \quad \text{and } \#\Omega =$$

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Our Experiment is taking 12 balls at random. Let us record, as an Outcome of our Experiment, the color of each of 12 balls. Say, one of the outcomes can be

$$(w, b, w, w, b, w, w, w, w, w, b, b).$$

In this case, the Sample Space is:

$$\Omega = \{(x_1, x_2, \dots, x_{12}) : x_k \in \{w, b\}\}, \quad \text{and } \#\Omega = 2^{12}.$$

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

$$\Omega = \{(x_1, x_2, \dots, x_{12}) : x_k \in \{w, b\}\}, \quad \text{and } \#\Omega = 2^{12}.$$

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

$$\Omega = \{(x_1, x_2, \dots, x_{12}) : x_k \in \{w, b\}\}, \quad \text{and } \#\Omega = 2^{12}.$$

Now, the Event in which we are interested is

$$A = \{(x_1, x_2, \dots, x_{12}) : \text{exactly 4 of } x_k\text{-s are w, and the rest are b}\}$$

And $\#A =$

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

$$\Omega = \{(x_1, x_2, \dots, x_{12}) : x_k \in \{w, b\}\}, \quad \text{and } \#\Omega = 2^{12}.$$

Now, the Event in which we are interested is

$$A = \{(x_1, x_2, \dots, x_{12}) : \text{exactly 4 of } x_k\text{-s are w, and the rest are b}\}$$

$$\text{And } \#A = \binom{12}{4}. \text{ So}$$

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

$$\Omega = \{(x_1, x_2, \dots, x_{12}) : x_k \in \{w, b\}\}, \quad \text{and } \#\Omega = 2^{12}.$$

Now, the Event in which we are interested is

$$A = \{(x_1, x_2, \dots, x_{12}) : \text{exactly 4 of } x_k\text{-s are w, and the rest are b}\}$$

$$\text{And } \#A = \binom{12}{4}. \text{ So}$$

$$\mathbb{P}(A) = \binom{12}{4} \cdot \frac{1}{2^{12}}.$$

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

$$\Omega = \{(x_1, x_2, \dots, x_{12}) : x_k \in \{w, b\}\}, \quad \text{and } \#\Omega = 2^{12}.$$

Now, the Event in which we are interested is

$$A = \{(x_1, x_2, \dots, x_{12}) : \text{exactly 4 of } x_k\text{-s are w, and the rest are b}\}$$

$$\text{And } \#A = \binom{12}{4}. \text{ So}$$

$$\mathbb{P}(A) = \binom{12}{4} \cdot \frac{1}{2^{12}}.$$

Right?

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

$$\Omega = \{(x_1, x_2, \dots, x_{12}) : x_k \in \{w, b\}\}, \quad \text{and } \#\Omega = 2^{12}.$$

Now, the Event in which we are interested is

$$A = \{(x_1, x_2, \dots, x_{12}) : \text{exactly 4 of } x_k\text{-s are } w, \text{ and the rest are } b\}$$

And $\#A = \binom{12}{4}$. So

$$\mathbb{P}(A) = \binom{12}{4} \cdot \frac{1}{2^{12}}.$$

Right? Not, of course!

Without Replacements Model, Solution 2:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

$$\Omega = \{(x_1, x_2, \dots, x_{12}) : x_k \in \{w, b\}\}, \quad \text{and } \#\Omega = 2^{12}.$$

Now, the Event in which we are interested is

$$A = \{(x_1, x_2, \dots, x_{12}) : \text{exactly 4 of } x_k\text{-s are w, and the rest are b}\}$$

And $\#A = \binom{12}{4}$. So

$$\mathbb{P}(A) = \binom{12}{4} \cdot \frac{1}{2^{12}}.$$

Right? Not, of course! Why?

Without Replacements Model, Solution 3:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Without Replacements Model, Solution 3:

Problem: We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

Now, the correct solution:

An Idea for Solving Probability Problems

- Assume we want to calculate the Probability of an Event A .

An Idea for Solving Probability Problems

- Assume we want to calculate the Probability of an Event A .
- Sometimes, when calculation of $\mathbb{P}(A)$ is not so easy, one can try calculating the probability of an Opposite Event, \overline{A} , and use

An Idea for Solving Probability Problems

- Assume we want to calculate the Probability of an Event A .
- Sometimes, when calculation of $\mathbb{P}(A)$ is not so easy, one can try calculating the probability of an Opposite Event, \overline{A} , and use

$$\mathbb{P}(A) = 1 - \mathbb{P}(\overline{A}).$$

An Idea for Solving Probability Problems

- Assume we want to calculate the Probability of an Event A .
- Sometimes, when calculation of $\mathbb{P}(A)$ is not so easy, one can try calculating the probability of an Opposite Event, \overline{A} , and use

$$\mathbb{P}(A) = 1 - \mathbb{P}(\overline{A}).$$

- This trick helps in different situations

Example: Birthday Problem

Problem: We have 36 participants in our group of Probability class, including the instructor. What is the probability that at least two participants share the same birthday?

Example: Birthday Problem

Problem: We have 36 participants in our group of Probability class, including the instructor. What is the probability that at least two participants share the same birthday?

Assumptions: There are 365 days in a year,

Example: Birthday Problem

Problem: We have 36 participants in our group of Probability class, including the instructor. What is the probability that at least two participants share the same birthday?

Assumptions: There are 365 days in a year, and the probability of being born on each day is the same.