

# AUA CS 108, Statistics, Fall 2019

## Lecture 40

Michael Poghosyan

YSU, AUA

[michael@ysu.am](mailto:michael@ysu.am), [mpoghosyan@aua.am](mailto:mpoghosyan@aua.am)

25 Nov 2019

# Contents

- ▶ Two-Sample Tests

## Last Lecture ReCap

- ▶ Describe the Asymptotic Test by MLE.

## Wald Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta;$

## Wald Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$ ; We take any Estimator  $\hat{\theta}$  which is Asymptotically Normal, when  $\theta = \theta_0$ :

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

## Wald Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$ ; We take any Estimator  $\hat{\theta}$  which is Asymptotically Normal, when  $\theta = \theta_0$ :

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

**Null Hypothesis:**  $\mathcal{H}_0 : \theta = \theta_0$

## Wald Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$ ; We take any Estimator  $\hat{\theta}$  which is Asymptotically Normal, when  $\theta = \theta_0$ :

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

**Null Hypothesis:**  $\mathcal{H}_0 : \theta = \theta_0$

**Asymptotic Significance Level:**  $\alpha \in (0, 1)$ ;

## Wald Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$ ; We take any Estimator  $\hat{\theta}$  which is Asymptotically Normal, when  $\theta = \theta_0$ :

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

**Null Hypothesis:**  $\mathcal{H}_0 : \theta = \theta_0$

**Asymptotic Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})}$  or  $W = \frac{\hat{\theta} - \theta_0}{SE(\theta_0)}$



## Wald Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$ ; We take any Estimator  $\hat{\theta}$  which is Asymptotically Normal, when  $\theta = \theta_0$ :

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

**Null Hypothesis:**  $\mathcal{H}_0 : \theta = \theta_0$

**Asymptotic Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})}$  or  $W = \frac{\hat{\theta} - \theta_0}{SE(\theta_0)}$

**Asymptotic Distrib of the TS Under  $\mathcal{H}_0$ :**  $W \xrightarrow{D}$

## Wald Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$ ; We take any Estimator  $\hat{\theta}$  which is Asymptotically Normal, when  $\theta = \theta_0$ :

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

**Null Hypothesis:**  $\mathcal{H}_0 : \theta = \theta_0$

**Asymptotic Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})}$  or  $W = \frac{\hat{\theta} - \theta_0}{SE(\theta_0)}$

**Asymptotic Distrib of the TS Under  $\mathcal{H}_0$ :**  $W \xrightarrow{D} \mathcal{N}(0, 1)$ ;

## Wald Test

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$ ; We take any Estimator  $\hat{\theta}$  which is Asymptotically Normal, when  $\theta = \theta_0$ :

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

**Null Hypothesis:**  $\mathcal{H}_0 : \theta = \theta_0$

**Asymptotic Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})}$  or  $W = \frac{\hat{\theta} - \theta_0}{SE(\theta_0)}$

**Asymptotic Distrib of the TS Under  $\mathcal{H}_0$ :**  $W \xrightarrow{D} \mathcal{N}(0, 1)$ ;

$\mathcal{H}_1$ is	RR is
$\theta \neq \theta_0$	$ W  > z_{1-\frac{\alpha}{2}}$
$\theta > \theta_0$	$W > z_{1-\alpha}$
$\theta < \theta_0$	$W < z_\alpha$

## Note

**Note:** In all above Asymptotic Tests, one can replace the quantiles  $z_p$  of the Standard Normal by the quantiles  $t_{n-1,p}$  of  $t(n-1)$ , since, for large  $n$ ,

$$t_{n-1,p} \approx z_p.$$

## Note

**Note:** In all above Asymptotic Tests, one can replace the quantiles  $z_p$  of the Standard Normal by the quantiles  $t_{n-1,p}$  of  $t(n-1)$ , since, for large  $n$ ,

$$t_{n-1,p} \approx z_p.$$

```
a <- 0.05  
qnorm(1-a/2)
```

```
## [1] 1.959964
```

```
n <- seq(50,250,50)  
qt(1-a/2, df = n)
```

```
## [1] 2.008559 1.983972 1.975905 1.971896 1.969498
```

## Note

**Note:** In all above Asymptotic Tests, one can replace the quantiles  $z_p$  of the Standard Normal by the quantiles  $t_{n-1,p}$  of  $t(n-1)$ , since, for large  $n$ ,

$$t_{n-1,p} \approx z_p.$$

```
a <- 0.05  
qnorm(1-a/2)
```

```
## [1] 1.959964
```

```
n <- seq(50,250,50)  
qt(1-a/2, df = n)
```

```
## [1] 2.008559 1.983972 1.975905 1.971896 1.969498
```

Another point is that, since  $|t_{n-1,p}| > |z_p|$ ,  $p \neq 0.5$ , it is safer to have a little bit smaller Rejection Region:

## Note

**Note:** In all above Asymptotic Tests, one can replace the quantiles  $z_p$  of the Standard Normal by the quantiles  $t_{n-1,p}$  of  $t(n-1)$ , since, for large  $n$ ,

$$t_{n-1,p} \approx z_p.$$

```
a <- 0.05  
qnorm(1-a/2)
```

```
## [1] 1.959964
```

```
n <- seq(50,250,50)  
qt(1-a/2, df = n)
```

```
## [1] 2.008559 1.983972 1.975905 1.971896 1.969498
```

Another point is that, since  $|t_{n-1,p}| > |z_p|$ ,  $p \neq 0.5$ , it is safer to have a little bit smaller Rejection Region: say, for the Two-Sided Tests, if  $|W| > t_{n-1,1-\alpha/2}$ , then, for sure, also  $|W| > z_{1-\alpha/2}$ .

# Two Sample Tests



## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2),$

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2), Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2),$

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  
 $\sigma_X, \sigma_Y$  are known,

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  
 $\sigma_X, \sigma_Y$  **are known**, and  $X_k$ -s and  $Y_j$ -s are all Independent.

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are known**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are known**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are known**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are known**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z =$



## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are known**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sigma_{\bar{X} - \bar{Y}}}$

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are known**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:** 
$$Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}$$

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are known**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:** 
$$Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}$$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim$

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are known**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:** 
$$Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}$$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim \mathcal{N}(0, 1)$ ;

## Z-Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  are known, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:** 
$$Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}$$

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $Z \sim \mathcal{N}(0, 1)$ ;

$\mathcal{H}_1$ is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ Z  > z_{1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$Z > z_{1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$Z < z_\alpha$

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2),$

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2), Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2),$

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  
 $\sigma_X, \sigma_Y$  are unknown,



## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  
 $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent.

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X = \sigma_Y$

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X = \sigma_Y$  (can be Tested separately, by  $F$ -Test)

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X = \sigma_Y$  (can be Tested separately, by  $F$ -Test)

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X = \sigma_Y$  (can be Tested separately, by  $F$ -Test)

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X = \sigma_Y$  (can be Tested separately, by  $F$ -Test)

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{(\bar{X} - \bar{Y}) - \mu_0}{S_p \cdot \sqrt{\frac{1}{n} + \frac{1}{m}}}$ , where  $S_p$  is the **Pooled**

**Sample Deviation:**

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X = \sigma_Y$  (can be Tested separately, by  $F$ -Test)

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**  $t = \frac{(\bar{X} - \bar{Y}) - \mu_0}{S_p \cdot \sqrt{\frac{1}{n} + \frac{1}{m}}}$ , where  $S_p$  is the **Pooled**

**Sample Deviation:**

$$S_p^2 = \frac{(n-1)S_X^2 + (m-1)S_Y^2}{n+m-2} = \frac{\sum_{k=1}^n (X_k - \bar{X})^2 + \sum_{k=1}^m (Y_k - \bar{Y})^2}{n+m-2}.$$



## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim$

## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n + m - 2)$ ;

## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n + m - 2)$ ;

**Rejection Region:**

$\mathcal{H}_1$ is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t  > t_{n+m-2, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{n+m-2, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{n+m-2, \alpha}$

## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**  $t \sim t(n + m - 2)$ ;

**Rejection Region:**

$\mathcal{H}_1$ is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t  > t_{n+m-2, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{n+m-2, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{n+m-2, \alpha}$

**Note:** This Test is called the **Pooled  $t$ -Test**

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2),$

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2), Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2),$

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  
 $\sigma_X, \sigma_Y$  are unknown,

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  
 $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent.



## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X \neq \sigma_Y$  (can be Tested separately, by  $F$ -Test)

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X \neq \sigma_Y$  (can be Tested separately, by  $F$ -Test)

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X \neq \sigma_Y$  (can be Tested separately, by  $F$ -Test)

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  are unknown, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X \neq \sigma_Y$  (can be Tested separately, by  $F$ -Test)

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**

$$t = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{S_X^2}{n} + \frac{S_Y^2}{m}}},$$

## $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  are unknown, and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Additional Assumption:**  $\sigma_X \neq \sigma_Y$  (can be Tested separately, by  $F$ -Test)

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**

$$t = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{S_X^2}{n} + \frac{S_Y^2}{m}}},$$

where  $S_X$  and  $S_Y$  are the Sample SDs for  $X$  and  $Y$ , respectively.

## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :** Approximately,

$$t \approx$$

## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :** Approximately,  
 $t \approx t(\nu)$ , where  $\nu$  is



## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :** Approximately,  
 $t \approx t(\nu)$ , where  $\nu$  is some scary thing...

## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :** Approximately,  
 $t \approx t(\nu)$ , where  $\nu$  is some scary thing... given by

$$\nu = \left\lfloor \frac{\left( \frac{S_X^2}{n} + \frac{S_Y^2}{m} \right)^2}{\frac{(S_X^2/n)^2}{n-1} + \frac{(S_Y^2/m)^2}{m-1}} \right\rfloor$$

## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :** Approximately,  
 $t \approx t(\nu)$ , where  $\nu$  is some scary thing... given by

$$\nu = \left\lfloor \frac{\left( \frac{S_X^2}{n} + \frac{S_Y^2}{m} \right)^2}{\frac{(S_X^2/n)^2}{n-1} + \frac{(S_Y^2/m)^2}{m-1}} \right\rfloor$$

**Rejection Region:**

$\mathcal{H}_1$ is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t  > t_{\nu, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{\nu, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{\nu, \alpha}$

## $t$ -Test for the Difference of two Normals Means, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :** Approximately,  
 $t \approx t(\nu)$ , where  $\nu$  is some scary thing... given by

$$\nu = \left\lfloor \frac{\left( \frac{S_X^2}{n} + \frac{S_Y^2}{m} \right)^2}{\frac{(S_X^2/n)^2}{n-1} + \frac{(S_Y^2/m)^2}{m-1}} \right\rfloor$$

**Rejection Region:**

$\mathcal{H}_1$ is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t  > t_{\nu, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{\nu, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{\nu, \alpha}$

**Note:** The formula above for the DF  $\nu$  is called **Welch – Satterthwaite Equation**, and the Tests is called the **Welch Test**.

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2),$

---

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2), Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2),$

---

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  
 $\sigma_X, \sigma_Y$  are unknown.

---

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  
 $\sigma_X, \sigma_Y$  **are unknown**. The Parameter of interest is  $\mu_X - \mu_Y$ ;

---



## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  
 $\sigma_X, \sigma_Y$  **are unknown**. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Note:** Here we have the same number of  $X_k$  and  $Y_k$ ;

---

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Note:** Here we have the same number of  $X_k$  and  $Y_k$ ; also, importantly,  $X_k$  and  $Y_k$  can be dependent!

---

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Note:** Here we have the same number of  $X_k$  and  $Y_k$ ; also, importantly,  $X_k$  and  $Y_k$  can be dependent!

**Notation:**  $D_k = X_k - Y_k$ ;

---

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Note:** Here we have the same number of  $X_k$  and  $Y_k$ ; also, importantly,  $X_k$  and  $Y_k$  can be dependent!

**Notation:**  $D_k = X_k - Y_k$ ; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

---

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Note:** Here we have the same number of  $X_k$  and  $Y_k$ ; also, importantly,  $X_k$  and  $Y_k$  can be dependent!

**Notation:**  $D_k = X_k - Y_k$ ; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

The Variance of  $D_k$ , although the same,  $\sigma_D^2 = \text{Var}(X_k - Y_k)$ , cannot be calculated, since  $X_k$  and  $Y_k$  can be dependent. But that's OK, we do not need it.

---

<sup>1</sup>The Test will work also in the case when the Differences are nor Normally Distributed, but the Sample Size  $n$  is large. We jut need to use the CLT.

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Note:** Here we have the same number of  $X_k$  and  $Y_k$ ; also, importantly,  $X_k$  and  $Y_k$  can be dependent!

**Notation:**  $D_k = X_k - Y_k$ ; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

The Variance of  $D_k$ , although the same,  $\sigma_D^2 = \text{Var}(X_k - Y_k)$ , cannot be calculated, since  $X_k$  and  $Y_k$  can be dependent. But that's OK, we do not need it.

**Additional Assumption:** We will assume that the Differences  $D_k$  are Normally Distributed<sup>1</sup>.

---

<sup>1</sup>The Test will work also in the case when the Differences are nor Normally Distributed, but the Sample Size  $n$  is large. We jut need to use the CLT.

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Note:** Here we have the same number of  $X_k$  and  $Y_k$ ; also, importantly,  $X_k$  and  $Y_k$  can be dependent!

**Notation:**  $D_k = X_k - Y_k$ ; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

The Variance of  $D_k$ , although the same,  $\sigma_D^2 = \text{Var}(X_k - Y_k)$ , cannot be calculated, since  $X_k$  and  $Y_k$  can be dependent. But that's OK, we do not need it.

**Additional Assumption:** We will assume that the Differences  $D_k$  are Normally Distributed<sup>1</sup>.

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

---

<sup>1</sup>The Test will work also in the case when the Differences are nor Normally Distributed, but the Sample Size  $n$  is large. We jut need to use the CLT.

## Paired $t$ -Test for the Difference of two Normals Means

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$ ,  $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$ ,  $\sigma_X, \sigma_Y$  **are unknown**. The Parameter of interest is  $\mu_X - \mu_Y$ ;

**Note:** Here we have the same number of  $X_k$  and  $Y_k$ ; also, importantly,  $X_k$  and  $Y_k$  can be dependent!

**Notation:**  $D_k = X_k - Y_k$ ; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

The Variance of  $D_k$ , although the same,  $\sigma_D^2 = \text{Var}(X_k - Y_k)$ , cannot be calculated, since  $X_k$  and  $Y_k$  can be dependent. But that's OK, we do not need it.

**Additional Assumption:** We will assume that the Differences  $D_k$  are Normally Distributed<sup>1</sup>.

**Null Hypothesis:**  $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

**Asymptotic Significance Level:**  $\alpha \in (0, 1)$ ;

<sup>1</sup>The Test will work also in the case when the Differences are nor Normally Distributed, but the Sample Size  $n$  is large. We jut need to use the CLT.



## Paired $t$ -Test for the Difference of two Normals Means, Cont'd

**Test Statistics:**  $t = \frac{\bar{D} - \mu_0}{S_D/\sqrt{n}}$ , where  $S_D$  is the Sample Deviation of  $D$ .

---

<sup>2</sup>Or, Asymptotically,  $t \approx t(n-1)$  or  $t \approx \mathcal{N}(0,1)$ , if  $D_k$ -s are not Normal, but  $n$  is large.

## Paired $t$ -Test for the Difference of two Normals Means, Cont'd

**Test Statistics:**  $t = \frac{\bar{D} - \mu_0}{S_D/\sqrt{n}}$ , where  $S_D$  is the Sample Deviation of  $D$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**<sup>2</sup>  $t \sim t(n-1)$ ;

---

<sup>2</sup>Or, Asymptotically,  $t \approx t(n-1)$  or  $t \approx \mathcal{N}(0,1)$ , if  $D_k$ -s are not Normal, but  $n$  is large.

## Paired $t$ -Test for the Difference of two Normals Means, Cont'd

**Test Statistics:**  $t = \frac{\bar{D} - \mu_0}{S_D/\sqrt{n}}$ , where  $S_D$  is the Sample Deviation of  $D$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**<sup>2</sup>  $t \sim t(n-1)$ ;

**Rejection Region:**

$\mathcal{H}_1$ is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t  > t_{n-1, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{n-1, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{n-1, \alpha}$

---

<sup>2</sup>Or, Asymptotically,  $t \approx t(n-1)$  or  $t \approx \mathcal{N}(0, 1)$ , if  $D_k$ -s are not Normal, but  $n$  is large.

## Paired $t$ -Test for the Difference of two Normals Means, Cont'd

**Test Statistics:**  $t = \frac{\bar{D} - \mu_0}{S_D/\sqrt{n}}$ , where  $S_D$  is the Sample Deviation of  $D$ .

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :**<sup>2</sup>  $t \sim t(n-1)$ ;

**Rejection Region:**

$\mathcal{H}_1$ is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t  > t_{n-1, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{n-1, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{n-1, \alpha}$

**Note:** This Test is called the **Paired  $t$ -Test**

---

<sup>2</sup>Or, Asymptotically,  $t \approx t(n-1)$  or  $t \approx \mathcal{N}(0, 1)$ , if  $D_k$ -s are not Normal, but  $n$  is large.

## Two Sample test for Proportions

**Model:**  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p_X),$

## Two Sample test for Proportions

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X),$   
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y),$

## Two Sample test for Proportions

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X)$ ,  
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y)$ , and  $X_k$ -s and  $Y_j$ -s are all Independent.

## Two Sample test for Proportions

**Model:**  $X_1, X_2, \dots, X_n \overset{IID}{\sim} \text{Bernoulli}(p_X)$ ,  
 $Y_1, Y_2, \dots, Y_m \overset{IID}{\sim} \text{Bernoulli}(p_Y)$ , and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $p_X - p_Y$ ;



## Two Sample test for Proportions

**Model:**  $X_1, X_2, \dots, X_n \overset{IID}{\sim} \text{Bernoulli}(p_X)$ ,  
 $Y_1, Y_2, \dots, Y_m \overset{IID}{\sim} \text{Bernoulli}(p_Y)$ , and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $p_X - p_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : p_X - p_Y = p_0$

## Two Sample test for Proportions

**Model:**  $X_1, X_2, \dots, X_n \overset{IID}{\sim} \text{Bernoulli}(p_X)$ ,  
 $Y_1, Y_2, \dots, Y_m \overset{IID}{\sim} \text{Bernoulli}(p_Y)$ , and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $p_X - p_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : p_X - p_Y = p_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

## Two Sample test for Proportions

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X)$ ,  
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y)$ , and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $p_X - p_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : p_X - p_Y = p_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**

$$Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\hat{p}(1 - \hat{p}) \left( \frac{1}{n} + \frac{1}{m} \right)}} \quad \text{or} \quad Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\frac{\bar{X}(1 - \bar{X})}{n} + \frac{\bar{Y}(1 - \bar{Y})}{m}}}$$

## Two Sample test for Proportions

**Model:**  $X_1, X_2, \dots, X_n \overset{IID}{\sim} \text{Bernoulli}(p_X)$ ,  
 $Y_1, Y_2, \dots, Y_m \overset{IID}{\sim} \text{Bernoulli}(p_Y)$ , and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $p_X - p_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : p_X - p_Y = p_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**

$$Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\hat{p}(1 - \hat{p}) \left( \frac{1}{n} + \frac{1}{m} \right)}} \quad \text{or} \quad Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\frac{\bar{X}(1 - \bar{X})}{n} + \frac{\bar{Y}(1 - \bar{Y})}{m}}}$$

where  $\hat{p}$  is the **Pooled Sample Proportion**:

$$\hat{p} = \frac{n}{n + m} \cdot \bar{X} + \frac{m}{n + m} \cdot \bar{Y}$$

## Two Sample test for Proportions

**Model:**  $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X)$ ,  
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y)$ , and  $X_k$ -s and  $Y_j$ -s are all Independent. The Parameter of interest is  $p_X - p_Y$ ;

**Null Hypothesis:**  $\mathcal{H}_0 : p_X - p_Y = p_0$

**Significance Level:**  $\alpha \in (0, 1)$ ;

**Test Statistics:**

$$Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\hat{p}(1 - \hat{p}) \left( \frac{1}{n} + \frac{1}{m} \right)}} \quad \text{or} \quad Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\frac{\bar{X}(1 - \bar{X})}{n} + \frac{\bar{Y}(1 - \bar{Y})}{m}}}$$

where  $\hat{p}$  is the **Pooled Sample Proportion**:

$$\hat{p} = \frac{n}{n+m} \cdot \bar{X} + \frac{m}{n+m} \cdot \bar{Y} = \frac{X_1 + \dots + X_n + Y_1 + \dots + Y_m}{n+m}.$$

## Two Sample test for Proportions, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :** Approximately,  
 $Z \approx$

## Two Sample test for Proportions, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :** Approximately,  
 $Z \approx \mathcal{N}(0, 1)$

## Two Sample test for Proportions, Cont'd

**Distrib of the Test-Statistics Under  $\mathcal{H}_0$ :** Approximately,  
 $Z \approx \mathcal{N}(0, 1)$

**Rejection Region:**

$\mathcal{H}_1$ is	RR is
$p_X - p_Y \neq p_0$	$ Z  > z_{1-\frac{\alpha}{2}}$
$p_X - p_Y > p_0$	$Z > z_{1-\alpha}$
$p_X - p_Y < p_0$	$Z < z_\alpha$