

CS 107, Probability, Spring 2019

Lecture 20

Michael Poghosyan

AUA

11 March 2019

Content

- CDF and its Properties
- Discrete r.v.

Welcome back from the Spring Break 😊

Properties of the CDF, reminder

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Properties of the CDF, reminder

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

Properties of the CDF, reminder

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;

Properties of the CDF, reminder

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;
- $F(-\infty) = 0$ and $F(+\infty) = 1$;

Properties of the CDF, reminder

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;
- $F(-\infty) = 0$ and $F(+\infty) = 1$;
- F is an increasing function, i.e., if $x_1 \leq x_2$, then $F(x_1) \leq F(x_2)$;

Properties of the CDF, reminder

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;
- $F(-\infty) = 0$ and $F(+\infty) = 1$;
- F is an increasing function, i.e., if $x_1 \leq x_2$, then $F(x_1) \leq F(x_2)$;
- F is right-continuous at every point, i.e. $F(x_0+) = F(x_0)$ at any $x_0 \in \mathbb{R}$

Properties of the CDF, reminder

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;
- $F(-\infty) = 0$ and $F(+\infty) = 1$;
- F is an increasing function, i.e., if $x_1 \leq x_2$, then $F(x_1) \leq F(x_2)$;
- F is right-continuous at every point, i.e. $F(x_0+) = F(x_0)$ at any $x_0 \in \mathbb{R}$

These four properties characterize completely CDFs!

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF.

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF. Assume X is a r.v., and $F(x)$ is its CDF. Then:

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF. Assume X is a r.v., and $F(x)$ is its CDF. Then:

Probabilities through CDF

- $\mathbb{P}(X = a) = F(a) - F(a-);$
- $\mathbb{P}(a < X \leq b) = F(b) - F(a);$
- $\mathbb{P}(a \leq X \leq b) = F(b) - F(a-);$
- $\mathbb{P}(a \leq X < b) = F(b-) - F(a-);$
- $\mathbb{P}(a < X < b) = F(b-) - F(a);$

Here we can have also $a = -\infty$ or/and $b = +\infty$

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 2)$, $\mathbb{P}(2 < X \leq 3)$, $\mathbb{P}(X \leq 5)$, $\mathbb{P}(X > 2)$:

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 2)$, $\mathbb{P}(2 < X \leq 3)$, $\mathbb{P}(X \leq 5)$, $\mathbb{P}(X > 2)$:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Prblm: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$, $\mathbb{P}(X < -3)$:

Reading Probabilities from the graph of CDF

Prblm: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$, $\mathbb{P}(X < -3)$:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Prblm: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$, $\mathbb{P}(X < -3)$:

Reading Probabilities from the graph of CDF

Prblm: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$, $\mathbb{P}(X < -3)$:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Problem: For the r.v. X given through its CDF below, which values are more probable:

Reading Probabilities from the graph of CDF

Problem: For the r.v. X given through its CDF below, which values are more probable:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$:

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Problem: Below you can find graphs of 2 CDFs: Red is for r.v. X , and Black is for Y . Which one is larger: $\mathbb{P}(2 < X < 4)$ or $\mathbb{P}(2 < Y < 4)$?

Reading Probabilities from the graph of CDF

Problem: Below you can find graphs of 2 CDFs: Red is for r.v. X , and Black is for Y . Which one is larger: $\mathbb{P}(2 < X < 4)$ or $\mathbb{P}(2 < Y < 4)$?




Figure: CDFs of r.v. X and Y

Discrete R.V.s

Discrete Random Variables

In our course, we will consider 2 types of r.v.: **Discrete** and **Continuous**.

¹Right Hand Side

Discrete Random Variables

In our course, we will consider 2 types of r.v.: **Discrete** and **Continuous**. Assume X is a r.v. defined on Ω , i.e., $X: \Omega \rightarrow \mathbb{R}$.

¹Right Hand Side

Discrete Random Variables

In our course, we will consider 2 types of r.v.: **Discrete** and **Continuous**. Assume X is a r.v. defined on Ω , i.e., $X: \Omega \rightarrow \mathbb{R}$.

Discrete Random Variable

We say that the r.v. X is **Discrete**, if the Range of X ,

$Range(X) = \{X(\omega) : \omega \in \Omega\}$ = The set of all possible values of X is finite or countably infinite.

¹Right Hand Side

Discrete Random Variables

In our course, we will consider 2 types of r.v.: **Discrete** and **Continuous**. Assume X is a r.v. defined on Ω , i.e., $X: \Omega \rightarrow \mathbb{R}$.

Discrete Random Variable

We say that the r.v. X is **Discrete**, if the Range of X ,

$Range(X) = \{X(\omega) : \omega \in \Omega\}$ = The set of all possible values of X is finite or countably infinite.

So if X is Discrete, then the Range of X can be written as

$$Range(X) = \{x_1, x_2, x_3, \dots\},$$

where the set on the RHS¹ can be also finite.

¹Right Hand Side

Discrete Random Variables

Examples:

- Let X be the number of Heads when tossing 5 coins.
What is the range of X ? Is X discrete?

Discrete Random Variables

Examples:

- Let X be the number of Heads when tossing 5 coins.
What is the range of X ? Is X discrete?
- Let Y be the number of children in the randomly chosen Armenian family,

Discrete Random Variables

Examples:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Let Y be the number of children in the randomly chosen Armenian family, so if Ω is the set of all Armenian families, $\omega \in \Omega$ is a family, and $Y(\omega)$ is the number of children in the family ω .

Discrete Random Variables

Examples:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Let Y be the number of children in the randomly chosen Armenian family, so if Ω is the set of all Armenian families, $\omega \in \Omega$ is a family, and $Y(\omega)$ is the number of children in the family ω .
- Let Z be the number of car accidents today in Yerevan;

Discrete Random Variables

Examples:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Let Y be the number of children in the randomly chosen Armenian family, so if Ω is the set of all Armenian families, $\omega \in \Omega$ is a family, and $Y(\omega)$ is the number of children in the family ω .
- Let Z be the number of car accidents today in Yerevan;
- Let U be the number of page clicks/search queries in Google;

Discrete Random Variables

Examples:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Let Y be the number of children in the randomly chosen Armenian family, so if Ω is the set of all Armenian families, $\omega \in \Omega$ is a family, and $Y(\omega)$ is the number of children in the family ω .
- Let Z be the number of car accidents today in Yerevan;
- Let U be the number of page clicks/search queries in Google; let V be the number of grammatical errors in my lecture slides, ...

Discrete Random Variables

Examples:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Let Y be the number of children in the randomly chosen Armenian family, so if Ω is the set of all Armenian families, $\omega \in \Omega$ is a family, and $Y(\omega)$ is the number of children in the family ω .
- Let Z be the number of car accidents today in Yerevan;
- Let U be the number of page clicks/search queries in Google; let V be the number of grammatical errors in my lecture slides, ...
- Can you give some more?

Discrete Random Variables

Examples:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Let Y be the number of children in the randomly chosen Armenian family, so if Ω is the set of all Armenian families, $\omega \in \Omega$ is a family, and $Y(\omega)$ is the number of children in the family ω .
- Let Z be the number of car accidents today in Yerevan;
- Let U be the number of page clicks/search queries in Google; let V be the number of grammatical errors in my lecture slides, ...
- Can you give some more?
- If Ω is Discrete, then X will be Discrete;

Discrete Random Variables

Examples:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Let Y be the number of children in the randomly chosen Armenian family, so if Ω is the set of all Armenian families, $\omega \in \Omega$ is a family, and $Y(\omega)$ is the number of children in the family ω .
- Let Z be the number of car accidents today in Yerevan;
- Let U be the number of page clicks/search queries in Google; let V be the number of grammatical errors in my lecture slides, ...
- Can you give some more?
- If Ω is Discrete, then X will be Discrete;
- If Ω is not Discrete, then X CAN BE Discrete: Example: for the Darts game, let X show the points you will get

Characterization of Discrete R.V.s

Now, assume X is a discrete r.v. with the Range (finite or countably infinite)

$$\text{Range}(X) = \{x_1, x_2, x_3, \dots\}.$$

Characterization of Discrete R.V.s

Now, assume X is a discrete r.v. with the Range (finite or countably infinite)

$$\text{Range}(X) = \{x_1, x_2, x_3, \dots\}.$$

Then the function $p(x) = \mathbb{P}(X = x)$, $x \in \mathbb{R}$ is called the Probability Mass Function (PMF) of X .

Characterization of Discrete R.V.s

Now, assume X is a discrete r.v. with the Range (finite or countably infinite)

$$\text{Range}(X) = \{x_1, x_2, x_3, \dots\}.$$

Then the function $p(x) = \mathbb{P}(X = x)$, $x \in \mathbb{R}$ is called the Probability Mass Function (PMF) of X .

In fact, PMF is non-zero only at points $x = x_k$, and we denote $p_k = \mathbb{P}(X = x_k)$, $k = 1, 2, 3, \dots$

Characterization of Discrete R.V.s

Now, assume X is a discrete r.v. with the Range (finite or countably infinite)

$$\text{Range}(X) = \{x_1, x_2, x_3, \dots\}.$$

Then the function $p(x) = \mathbb{P}(X = x)$, $x \in \mathbb{R}$ is called the Probability Mass Function (PMF) of X .

In fact, PMF is non-zero only at points $x = x_k$, and we denote $p_k = \mathbb{P}(X = x_k)$, $k = 1, 2, 3, \dots$. And in this case we write PMF in the table form:

Values of X	x_1	x_2	x_3	\dots
$\mathbb{P}(X = x)$	p_1	p_2	p_3	\dots

Characterization of Discrete R.V.s

Now, assume X is a discrete r.v. with the Range (finite or countably infinite)

$$\text{Range}(X) = \{x_1, x_2, x_3, \dots\}.$$

Then the function $p(x) = \mathbb{P}(X = x)$, $x \in \mathbb{R}$ is called the Probability Mass Function (PMF) of X .

In fact, PMF is non-zero only at points $x = x_k$, and we denote $p_k = \mathbb{P}(X = x_k)$, $k = 1, 2, 3, \dots$. And in this case we write PMF in the table form:

Values of X	x_1	x_2	x_3	\dots
$\mathbb{P}(X = x)$	p_1	p_2	p_3	\dots

We will usually write PMF in the table form.

Characterization of PMF

Clearly, if X is a discrete r.v. with the PMF (in the table form):

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

then

Characterization of PMF

Clearly, if X is a discrete r.v. with the PMF (in the table form):

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

then

- $p_k \geq 0, k = 1, 2, 3, \dots$

Characterization of PMF

Clearly, if X is a discrete r.v. with the PMF (in the table form):

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

then

- $p_k \geq 0, k = 1, 2, 3, \dots$
- $\sum_k p_k =$

Characterization of PMF

Clearly, if X is a discrete r.v. with the PMF (in the table form):

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

then

- $p_k \geq 0, k = 1, 2, 3, \dots$
- $\sum_k p_k = 1$

Characterization of PMF

Clearly, if X is a discrete r.v. with the PMF (in the table form):

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

then

- $p_k \geq 0, k = 1, 2, 3, \dots$
- $\sum_k p_k = 1$

Question: What is the difference between Discrete Probability Models and Discrete r.v.s?

PMF and Calculation of Probabilities

Recall that, given a r.v. X , our aim was to be able to calculate the Probabilities of the type

$$\mathbb{P}(X \in A), \quad A \subset \mathbb{R}.$$

PMF and Calculation of Probabilities

Recall that, given a r.v. X , our aim was to be able to calculate the Probabilities of the type

$$\mathbb{P}(X \in A), \quad A \subset \mathbb{R}.$$

Now, in the case we have the PMF of X ,

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

we can calculate

$$\mathbb{P}(X \in A) =$$

PMF and Calculation of Probabilities

Recall that, given a r.v. X , our aim was to be able to calculate the Probabilities of the type

$$\mathbb{P}(X \in A), \quad A \subset \mathbb{R}.$$

Now, in the case we have the PMF of X ,

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

we can calculate

$$\mathbb{P}(X \in A) = \sum_{x_k \in A} p_k.$$

PMF and Calculation of Probabilities

Recall that, given a r.v. X , our aim was to be able to calculate the Probabilities of the type

$$\mathbb{P}(X \in A), \quad A \subset \mathbb{R}.$$

Now, in the case we have the PMF of X ,

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

we can calculate

$$\mathbb{P}(X \in A) = \sum_{x_k \in A} p_k.$$

For example,

$$\mathbb{P}(a \leq X \leq b) = \sum_{a \leq x_k \leq b} p_k.$$

PMF and CDF for a Discrete r.v.

In particular, if the Discrete r.v. X is given by its PMF

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

and if $F(x)$ is the CDF of X , then:

$$F(x) =$$

PMF and CDF for a Discrete r.v.

In particular, if the Discrete r.v. X is given by its PMF

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

and if $F(x)$ is the CDF of X , then:

$$F(x) = \mathbb{P}(X \leq x) =$$

PMF and CDF for a Discrete r.v.

In particular, if the Discrete r.v. X is given by its PMF

Values of X	x_1	x_2	x_3	...
$\mathbb{P}(X = x)$	p_1	p_2	p_3	...

and if $F(x)$ is the CDF of X , then:

$$F(x) = \mathbb{P}(X \leq x) = \sum_{x_k \leq x} p_k.$$

Example:

Let X is the number of heads shown when tossing a fair coin 4 times. Then X is a r.v. with the PMF:

Values of X					
$\mathbb{P}(X = x)$					

Example:

Let X is the number of heads shown when tossing a fair coin 4 times. Then X is a r.v. with the PMF:

Values of X					
$\mathbb{P}(X = x)$					

Now,

- Calculate $\mathbb{P}(X = 2)$;

Example:

Let X is the number of heads shown when tossing a fair coin 4 times. Then X is a r.v. with the PMF:

Values of X					
$\mathbb{P}(X = x)$					

Now,

- Calculate $\mathbb{P}(X = 2)$;
- Calculate $\mathbb{P}(X \leq 2.5)$;

Example:

Let X is the number of heads shown when tossing a fair coin 4 times. Then X is a r.v. with the PMF:

Values of X					
$\mathbb{P}(X = x)$					

Now,

- Calculate $\mathbb{P}(X = 2)$;
- Calculate $\mathbb{P}(X \leq 2.5)$;
- Graph the PMF of X ;

Example:

Let X is the number of heads shown when tossing a fair coin 4 times. Then X is a r.v. with the PMF:

Values of X					
$\mathbb{P}(X = x)$					

Now,

- Calculate $\mathbb{P}(X = 2)$;
- Calculate $\mathbb{P}(X \leq 2.5)$;
- Graph the PMF of X ;
- Graph the CDF of X .