

# AUA CS 108, Statistics, Fall 2019

## Lecture 25

Michael Poghosyan

YSU, AUA

[michael@ysu.am](mailto:michael@ysu.am), [mpoghosyan@aua.am](mailto:mpoghosyan@aua.am)

21 Oct 2019

# Contents

- ▶ Fisher Information
- ▶ Cramer-Rao Lower Bound

## Last Lecture ReCap

- ▶ Give the definition of a Consistent Estimator

# Last Lecture ReCap

- ▶ Give the definition of a Consistent Estimator
- ▶ In which case and in which way we can use Unbiased Estimator?

## Last Lecture ReCap

- ▶ Give the definition of a Consistent Estimator
- ▶ In which case and in which way we can use Unbiased Estimator?
- ▶ In which case and in which way we can use Consistent Estimator?

# Fisher Information, Refresher

Assume we have a parametric family of distributions  $\mathcal{F}_\theta$ ,  $\theta \in \Theta$ , and  $f(x|\theta)$  is the PD(M)F of  $\mathcal{F}_\theta$ .

# Fisher Information, Refresher

Assume we have a parametric family of distributions  $\mathcal{F}_\theta$ ,  $\theta \in \Theta$ , and  $f(x|\theta)$  is the PD(M)F of  $\mathcal{F}_\theta$ .

**Definition:** The following quantity is called **the Fisher Information** of the parametric family  $\mathcal{F}_\theta$ :

$$I(\theta) = -\mathbb{E} \left( \frac{\partial^2}{\partial \theta^2} \ln f(X|\theta) \right) = \mathbb{E} \left[ \left( \frac{\partial}{\partial \theta} \ln f(X|\theta) \right)^2 \right],$$

where  $X \sim \mathcal{F}_\theta$ .

## Example

**Example:** Calculate the Fisher Information for the  $Bernoulli(p)$  family

**Solution:** OTB



## Example

**Example:** Calculate the Fisher Information for the  $Bernoulli(p)$  family

**Solution:** OTB

**Example:** Calculate the Fisher Information for the  $\mathcal{N}(\mu, \sigma^2)$  family (separately for the Parameter  $\mu$  and  $\sigma^2$ )

**Solution:** OTB

## Cramer-Rao Inequality, C-R Lower Bound

This is a remarkable inequality, giving a lower bound for the Variance of an Unbiased Estimator.

## Cramer-Rao Inequality, C-R Lower Bound

This is a remarkable inequality, giving a lower bound for the Variance of an Unbiased Estimator. Under some regularity conditions on the family of Distributions  $\mathcal{F}_\theta$ , the following holds:

## Cramer-Rao Inequality, C-R Lower Bound

This is a remarkable inequality, giving a lower bound for the Variance of an Unbiased Estimator. Under some regularity conditions on the family of Distributions  $\mathcal{F}_\theta$ , the following holds:

**Theorem (Cramer-Rao):** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$$

and the Fisher Information for the family  $\mathcal{F}_\theta$  is  $I(\theta)$ . Assume also that  $\hat{\theta}$  is an unbiased estimator for  $\theta$  obtained from our Random Sample. Then, under the above mentioned regularity conditions,

$$\text{Var}(\hat{\theta}) \geq \frac{1}{n \cdot I(\theta)}.$$

## Cramer-Rao Inequality, C-R Lower Bound

This is a remarkable inequality, giving a lower bound for the Variance of an Unbiased Estimator. Under some regularity conditions on the family of Distributions  $\mathcal{F}_\theta$ , the following holds:

**Theorem (Cramer-Rao):** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$$

and the Fisher Information for the family  $\mathcal{F}_\theta$  is  $I(\theta)$ . Assume also that  $\hat{\theta}$  is an unbiased estimator for  $\theta$  obtained from our Random Sample. Then, under the above mentioned regularity conditions,

$$\text{Var}(\hat{\theta}) \geq \frac{1}{n \cdot I(\theta)}.$$

**Note:** There is a version of C-R Inequality for the Biased case.

## Consequences of the C-R LB

Recall that for an Unbiased Estimator  $\hat{\theta}$ ,

$$MSE(\hat{\theta}, \theta) = Var(\hat{\theta}).$$

## Consequences of the C-R LB

Recall that for an Unbiased Estimator  $\hat{\theta}$ ,

$$MSE(\hat{\theta}, \theta) = Var(\hat{\theta}).$$

So the C-R LB gives us

$$MSE(\hat{\theta}, \theta) \geq \frac{1}{n \cdot I(\theta)}.$$

## Consequences of the C-R LB

Recall that for an Unbiased Estimator  $\hat{\theta}$ ,

$$MSE(\hat{\theta}, \theta) = Var(\hat{\theta}).$$

So the C-R LB gives us

$$MSE(\hat{\theta}, \theta) \geq \frac{1}{n \cdot I(\theta)}.$$

And this is a fundamental restriction on the MSE: *you cannot do better when estimating  $\theta$  than the Estimator with*

$$MSE(\hat{\theta}, \theta) = \frac{1}{n \cdot I(\theta)},$$

you cannot obtain smaller MSE using Unbiased Estimators!



## Consequences of the C-R LB

Recall that for an Unbiased Estimator  $\hat{\theta}$ ,

$$MSE(\hat{\theta}, \theta) = Var(\hat{\theta}).$$

So the C-R LB gives us

$$MSE(\hat{\theta}, \theta) \geq \frac{1}{n \cdot I(\theta)}.$$

And this is a fundamental restriction on the MSE: *you cannot do better when estimating  $\theta$  than the Estimator with*

$$MSE(\hat{\theta}, \theta) = \frac{1}{n \cdot I(\theta)},$$

you cannot obtain smaller MSE using Unbiased Estimators!

And if there exists an Unbiased Estimator  $\hat{\theta}$  with

$$MSE(\hat{\theta}, \theta) = \frac{1}{n \cdot I(\theta)},$$

we call  $\hat{\theta}$  an **Efficient Estimator** for  $\theta$ , and that Estimator is a MVUE for  $\theta$ .

## Notes

**Note:** Not always there exists an Unbiased Estimator with

$$MSE(\hat{\theta}, \theta) = Var(\hat{\theta}) = \frac{1}{n \cdot I(\theta)}.$$

## Notes

**Note:** Not always there exists an Unbiased Estimator with

$$MSE(\hat{\theta}, \theta) = Var(\hat{\theta}) = \frac{1}{n \cdot I(\theta)}.$$

But even in that cases we can have MVUE (and, in that case, the minimum of the Variance will be  $> \frac{1}{n \cdot I(\theta)}$ ).

# Notes

**Note:** Not always there exists an Unbiased Estimator with

$$MSE(\hat{\theta}, \theta) = Var(\hat{\theta}) = \frac{1}{n \cdot I(\theta)}.$$

But even in that cases we can have MVUE (and, in that case, the minimum of the Variance will be  $> \frac{1}{n \cdot I(\theta)}$ ).

**Note:** Sometimes, in different Textbooks, an Unbiased Estimator with Minimum Variance (not necessarily with  $Var(\hat{\theta}) = \frac{1}{n \cdot I(\theta)}$ ) is called an **Efficient Estimator** for  $\theta$ .

## Example

**Example:** Show that in the Bernoulli Model, with a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p), \quad p \in [0, 1],$$

the Estimator

$$\hat{p} = \bar{X}$$

is the MVUE of  $p$ .

## Example

**Example:** Show that in the Bernoulli Model, with a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p), \quad p \in [0, 1],$$

the Estimator

$$\hat{p} = \bar{X}$$

is the MVUE of  $p$ .

**Example:** Show that in the Poisson Model, with a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda), \quad \lambda > 0,$$

the Estimator

$$\hat{\lambda} = \bar{X}$$

is the MVUE of  $\lambda$ .