CS 107, Probability, Spring 2019 Lecture 37

Michael Poghosyan

AUA

19 April 2019

Content

• Independent Random Variables

Last time we have talked about the following Proposition:

Independence of R.V.s

Last time we have talked about the following Proposition:

Independence of R.V.s

The followings are equivalent:

• X and Y are Independent, $X \perp \!\!\! \perp Y$, i.e., $\mathbb{P}(X \in A, Y \in B) = \mathbb{P}(X \in A) \cdot \mathbb{P}(Y \in B)$ for any $A, B \subset \mathbb{R}$;

Last time we have talked about the following Proposition:

Independence of R.V.s

- X and Y are Independent, $X \perp \!\!\! \perp Y$, i.e., $\mathbb{P}(X \in A, Y \in B) = \mathbb{P}(X \in A) \cdot \mathbb{P}(Y \in B)$ for any $A, B \subset \mathbb{R}$;
- $F_{X,Y}(x,y) = F_X(x) \cdot F_Y(y)$ for any $x,y \in \mathbb{R}$, where $F_{X,Y}$, F_X , F_Y are the Joint CDF of X, Y and the Marginal CDFs of X and Y, respectively;

Last time we have talked about the following Proposition:

Independence of R.V.s

- X and Y are Independent, $X \perp \!\!\! \perp Y$, i.e., $\mathbb{P}(X \in A, Y \in B) = \mathbb{P}(X \in A) \cdot \mathbb{P}(Y \in B)$ for any $A, B \subset \mathbb{R}$;
- $F_{X,Y}(x,y) = F_X(x) \cdot F_Y(y)$ for any $x,y \in \mathbb{R}$, where $F_{X,Y}$, F_X , F_Y are the Joint CDF of X, Y and the Marginal CDFs of X and Y, respectively;
- If X and Y are Discrete, then $\mathbb{P}(X = x_i, Y = y_j) = \mathbb{P}(X = x_i) \cdot \mathbb{P}(Y = y_j)$, for all i, j

Last time we have talked about the following Proposition:

Independence of R.V.s

- X and Y are Independent, $X \perp \!\!\! \perp Y$, i.e., $\mathbb{P}(X \in A, Y \in B) = \mathbb{P}(X \in A) \cdot \mathbb{P}(Y \in B)$ for any $A, B \subset \mathbb{R}$;
- $F_{X,Y}(x,y) = F_X(x) \cdot F_Y(y)$ for any $x,y \in \mathbb{R}$, where $F_{X,Y}$, F_X , F_Y are the Joint CDF of X, Y and the Marginal CDFs of X and Y, respectively;
- If X and Y are Discrete, then $\mathbb{P}(X = x_i, Y = y_j) = \mathbb{P}(X = x_i) \cdot \mathbb{P}(Y = y_j)$, for all i, j
- If X and Y are Jointly Continuous, then $f_{X,Y}(x,y) = f_X(x) \cdot f_Y(y)$ for any f_X , where f_X , where f_X , where f_X , f_Y are the Joint PDF of X, Y and the Marginal PDFs of X.

Type 1 Examples: Given the Joint Distribution, we study the Independence.

Type 1 Examples: Given the Joint Distribution, we study the Independence.

• Assume $(X, Y) \sim \textit{Unif}([0, 1] \times [2, 4])$. Are X and Y Independent?

Type 1 Examples: Given the Joint Distribution, we study the Independence.

- Assume $(X, Y) \sim \textit{Unif}([0, 1] \times [2, 4])$. Are X and Y Independent?
- Assume X ~ Unif[0,1] and Y ~ Unif[2,4]. Are X and Y Independent?

Type 1 Examples: Given the Joint Distribution, we study the Independence.

- Assume $(X, Y) \sim \textit{Unif}([0, 1] \times [2, 4])$. Are X and Y Independent?
- Assume X ~ Unif[0,1] and Y ~ Unif[2,4]. Are X and Y Independent?
- Assume $(X, Y) \sim \textit{Unif}(T)$, where T is the triangle with the vertices at (0,0), (0,1) and (1,0). Are X and Y Independent?

Type 1 Examples: Given the Joint Distribution, we study the Independence.

- Assume $(X, Y) \sim \textit{Unif}([0, 1] \times [2, 4])$. Are X and Y Independent?
- Assume X ~ Unif[0,1] and Y ~ Unif[2,4]. Are X and Y Independent?
- Assume $(X, Y) \sim \textit{Unif}(T)$, where T is the triangle with the vertices at (0,0), (0,1) and (1,0). Are X and Y Independent?

Type 2 Examples: Given Individual Distributions of X and Y, and Independence, we form the Joint Distribution and calculate Probabilities.

Type 1 Examples: Given the Joint Distribution, we study the Independence.

- Assume $(X, Y) \sim \textit{Unif}([0, 1] \times [2, 4])$. Are X and Y Independent?
- Assume X ~ Unif[0,1] and Y ~ Unif[2,4]. Are X and Y Independent?
- Assume $(X, Y) \sim \textit{Unif}(T)$, where T is the triangle with the vertices at (0,0), (0,1) and (1,0). Are X and Y Independent?

Type 2 Examples: Given Individual Distributions of X and Y, and Independence, we form the Joint Distribution and calculate Probabilities.

• Assume $X \sim \textit{Unif}[0,3]$, $Y \sim \textit{Exp}(2)$ and $X \perp \!\!\! \perp Y$. Find $\mathbb{P}(X^2 + Y^2 \leq 1)$.



Facts About Multivariate Uniform Distribution:

Fact 1: Assume $X_1, ..., X_n$ are Independent and $X_k \sim Unif(A_k)$, k = 1, ..., n.

Facts About Multivariate Uniform Distribution:

Fact 1: Assume $X_1, ..., X_n$ are Independent and $X_k \sim Unif(A_k)$, k = 1, ..., n. Then

$$(X_1,...,X_n) \sim Unif(A_1 \times A_2 \times ... \times A_n).$$

Facts About Multivariate Uniform Distribution:

Fact 1: Assume $X_1, ..., X_n$ are Independent and $X_k \sim \textit{Unif}(A_k)$, k = 1, ..., n. Then

$$(X_1,...,X_n) \sim Unif(A_1 \times A_2 \times ... \times A_n).$$

Fact 2: Inversely, if the r.vector $(X_1, ..., X_n)$ has a Multivariate Uniform Distribution on a set, which is a Cartesian Product of n sets $A_1, ..., A_n \subset \mathbb{R}$, i.e., if

$$(X_1,...,X_n) \sim Unif(A_1 \times A_2 \times ... \times A_n),$$

then $X_1,...,X_n$ are Independent and $X_k \sim \textit{Unif}(A_k)$, k = 1,...,n.

Facts About Multivariate Normal Distribution:

Fact 1: Assume $X_1, ..., X_n$ are Independent and $X_k \sim \mathcal{N}(\mu_k, \sigma_k^2)$, k = 1, ..., n.

Facts About Multivariate Normal Distribution:

Fact 1: Assume $X_1, ..., X_n$ are Independent and $X_k \sim \mathcal{N}(\mu_k, \sigma_k^2)$, k = 1, ..., n. Then

$$(X_1,...,X_n) \sim \mathcal{N}(\mu,\Sigma),$$

where

$$\mu = \begin{bmatrix} \mu_1 \\ \vdots \\ \mu_n \end{bmatrix}, \quad \text{and} \quad \Sigma = \begin{bmatrix} \sigma_1^2 & 0 & \cdots & 0 \\ 0 & \sigma_2^2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sigma_n^2 \end{bmatrix}.$$

Facts About Multivariate Normal Distribution:

Fact 1: Assume $X_1, ..., X_n$ are Independent and $X_k \sim \mathcal{N}(\mu_k, \sigma_k^2)$, k = 1, ..., n. Then

$$(X_1,...,X_n) \sim \mathcal{N}(\mu,\Sigma),$$

where

$$\mu = \begin{bmatrix} \mu_1 \\ \vdots \\ \mu_n \end{bmatrix}, \quad \text{and} \quad \Sigma = \begin{bmatrix} \sigma_1^2 & 0 & \cdots & 0 \\ 0 & \sigma_2^2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sigma_n^2 \end{bmatrix}.$$

Fact 2: If we have a r.vector $(X_1,...,X_n)$ with $(X_1,...,X_n) \sim \mathcal{N}(\mu,\Sigma)$, where μ and Σ are as above, then $X_1,...,X_n$ are Independent, and

$$X_k \sim \mathcal{N}(\mu_k, \sigma_k^2), \qquad k = 1, ..., n.$$

Distribution of the Sum of Independent R.V.s

Now, assume Z = X + Y, and assume that now $X \perp \!\!\! \perp Y$.

Distribution of the Sum of Independent R.V.s.

Now, assume Z = X + Y, and assume that now $X \perp \!\!\! \perp Y$.

• If X and Y are **Discrete** and Independent, then

$$\mathbb{P}(Z=x) = \mathbb{P}(X+Y=x) = \sum_{x_i+y_i=x} \mathbb{P}(X=x_i, Y=y_j) =$$

Distribution of the Sum of Independent R.V.s.

Now, assume Z = X + Y, and assume that now $X \perp\!\!\!\perp Y$.

• If X and Y are **Discrete** and Independent, then

$$\mathbb{P}(Z = x) = \mathbb{P}(X + Y = x) = \sum_{x_i + y_j = x} \mathbb{P}(X = x_i, Y = y_j) =$$

$$= \sum_{x_i + y_j = x} \mathbb{P}(X = x_i) \cdot \mathbb{P}(Y = y_j)$$

 $x_i + v_i = x$

Distribution of the Sum of Independent R.V.s

Now, assume Z = X + Y, and assume that now $X \perp\!\!\!\perp Y$.

• If X and Y are **Discrete** and Independent, then

$$\mathbb{P}(Z=x) = \mathbb{P}(X+Y=x) = \sum_{x_i+y_j=x} \mathbb{P}(X=x_i, Y=y_j) =$$

$$=\sum_{x_i+y_j=x}\mathbb{P}(X=x_i)\cdot\mathbb{P}(Y=y_j)=\sum_{x_i}\mathbb{P}(X=x_i)\cdot\mathbb{P}(Y=x-x_i).$$

Distribution of the Sum of Independent R.V.s.

Now, assume Z = X + Y, and assume that now $X \perp \!\!\! \perp Y$.

• If X and Y are **Discrete** and Independent, then

$$\mathbb{P}(Z=x) = \mathbb{P}(X+Y=x) = \sum_{x_i+y_j=x} \mathbb{P}(X=x_i, Y=y_j) =$$

$$\sum_{x_i+y_j=x} \mathbb{P}(X=x_i, Y=y_j) = \sum_{x_i+y_j=x} \mathbb{P}(X=x_i, Y=y_i) = \sum_{x_i+y_i=x} \mathbb{P}(X=x_i, Y=x_i) = \sum_{x_i+y_i=x} \mathbb{P}(X=$$

$$=\sum_{x_i+y_j=x}\mathbb{P}(X=x_i)\cdot\mathbb{P}(Y=y_j)=\sum_{x_i}\mathbb{P}(X=x_i)\cdot\mathbb{P}(Y=x-x_i).$$

• If X and Y are **Jointly Continuous** with the Joint PDF $f_{X,Y}(x,y)$ and Independent, then, for any $x \in \mathbb{R}$,

$$f_{X+Y}(x) = \int_{-\infty}^{\infty} f_{X,Y}(t, x-t) dt =$$



Distribution of the Sum of Independent R.V.s.

Now, assume Z = X + Y, and assume that now $X \perp \!\!\! \perp Y$.

• If X and Y are **Discrete** and Independent, then

$$\mathbb{P}(Z=x) = \mathbb{P}(X+Y=x) = \sum_{x_i+y_j=x} \mathbb{P}(X=x_i, Y=y_j) =$$

$$= \sum_{x_i+y_j=x} \mathbb{P}(X=x_i) \cdot \mathbb{P}(Y=y_j) = \sum_{x_i+y_j=x} \mathbb{P}(X=x_i) \cdot \mathbb{P}(Y=x-x_i).$$

$$\sum_{x_i+y_j=x} \mathbb{E}(x_i \times x_i) \mathbb{E}(x_i \times x_j) \mathbb{E}(x_j \times x_j$$

• If X and Y are **Jointly Continuous** with the Joint PDF $f_{X,Y}(x,y)$ and Independent, then, for any $x \in \mathbb{R}$,

$$f_{X+Y}(x) = \int_{-\infty}^{\infty} f_{X,Y}(t,x-t) dt = \int_{-\infty}^{\infty} f_X(t) \cdot f_Y(x-t) dt.$$

