

# CS 107, Probability, Spring 2019

## Lecture 28

Michael Poghosyan

AUA

29 March 2019

- Normal Distribution
- Functions of Random Variables (aka Transformations of Random Variables)

This could be in one of our Midterms 😊

This could be in one of our Midterms 😊

What is the Probability of choosing the correct answer to this MCh question at random?

This could be in one of our Midterms 😊

What is the Probability of choosing the correct answer to this MCh question at random?

- 1  $\frac{1}{4}$
- 2  $\frac{1}{2}$
- 3  $\frac{1}{1}$
- 4  $\frac{1}{4}$

This could be in one of our Midterms 😊

What is the Probability of choosing the correct answer to this MCh question at random?

- 1  $\frac{1}{4}$
- 2  $\frac{1}{2}$
- 3  $\frac{1}{1}$
- 4  $\frac{1}{4}$

Your answer is not correct! :P

# Normal Distribution

Recall that the Normal Distribution is one of the most important distributions in the Universe 😊

# Normal Distribution

Recall that the Normal Distribution is one of the most important distributions in the Universe ☺

And the PDF of  $X \sim \mathcal{N}(\mu, \sigma^2)$  was

$$f(x) =$$



# Normal Distribution

Recall that the Normal Distribution is one of the most important distributions in the Universe ☺

And the PDF of  $X \sim \mathcal{N}(\mu, \sigma^2)$  was

$$f(x) = \frac{1}{\sigma \cdot \sqrt{2\pi}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}.$$

where  $\mu$  is the Mean of  $X$  and  $\sigma^2$  is the Variance of  $X$ .

# Normal Distribution

Recall that the Normal Distribution is one of the most important distributions in the Universe ☺

And the PDF of  $X \sim \mathcal{N}(\mu, \sigma^2)$  was

$$f(x) = \frac{1}{\sigma \cdot \sqrt{2\pi}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}.$$

where  $\mu$  is the Mean of  $X$  and  $\sigma^2$  is the Variance of  $X$ .

Recall also that if  $X \sim \mathcal{N}(0, 1)$ , then we say that  $X$  is Standard Normal r.v. (or has a Standard Normal Distribution).

# Some Examples

**Example:** Assume that the heights of women in Armenia are Normally distributed with the Mean 158.1cm and Standard Deviation 5.7cm<sup>1</sup>.

---

<sup>1</sup>See <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0018962>

# Some Examples

**Example:** Assume that the heights of women in Armenia are Normally distributed with the Mean 158.1cm and Standard Deviation 5.7cm<sup>1</sup>.

- Calculate the Probability that the (randomly chosen) woman height will be smaller than 158.1cm.

---

<sup>1</sup>See <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0018962>

# Some Examples

**Example:** Assume that the heights of women in Armenia are Normally distributed with the Mean 158.1cm and Standard Deviation 5.7cm<sup>1</sup>.

- Calculate the Probability that the (randomly chosen) woman height will be smaller than 158.1cm.
- Calculate the Probability that the (randomly chosen) woman height will be larger than 170cm.

---

<sup>1</sup>See <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0018962>

# Note: Normal Distribution

Assume  $X \sim \mathcal{N}(\mu, \sigma^2)$ . Then

# Note: Normal Distribution

Assume  $X \sim \mathcal{N}(\mu, \sigma^2)$ . Then

- $\mathbb{P}(\mu - \sigma < X < \mu + \sigma) \approx 0.68$

# Note: Normal Distribution

Assume  $X \sim \mathcal{N}(\mu, \sigma^2)$ . Then

- $\mathbb{P}(\mu - \sigma < X < \mu + \sigma) \approx 0.68$
- $\mathbb{P}(\mu - 2\sigma < X < \mu + 2\sigma) \approx 0.95$




# Note: Normal Distribution

Assume  $X \sim \mathcal{N}(\mu, \sigma^2)$ . Then

- $\mathbb{P}(\mu - \sigma < X < \mu + \sigma) \approx 0.68$
- $\mathbb{P}(\mu - 2\sigma < X < \mu + 2\sigma) \approx 0.95$
- $\mathbb{P}(\mu - 3\sigma < X < \mu + 3\sigma) \approx 0.997$

# Note: Normal Distribution



# Note: Normal Distribution

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Y = X - \mu \sim$$

# Note: Normal Distribution

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Y = X - \mu \sim \mathcal{N}(0, \sigma^2);$$

# Note: Normal Distribution

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Y = X - \mu \sim \mathcal{N}(0, \sigma^2);$$

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Z = \frac{X - \mu}{\sigma} \sim$$

# Note: Normal Distribution

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Y = X - \mu \sim \mathcal{N}(0, \sigma^2);$$

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Z = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1)$$

so  $Z = \frac{X - \mu}{\sigma}$  is a Standard Normal Variable.

# Note: Normal Distribution

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Y = X - \mu \sim \mathcal{N}(0, \sigma^2);$$

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Z = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1)$$

so  $Z = \frac{X - \mu}{\sigma}$  is a Standard Normal Variable. This transformation is called a Standardization, and  $Z$  usually named the  $Z$ -score (of  $X$ ).

# Note: Normal Distribution

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Y = X - \mu \sim \mathcal{N}(0, \sigma^2);$$

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Z = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1)$$

so  $Z = \frac{X - \mu}{\sigma}$  is a Standard Normal Variable. This transformation is called a Standardization, and  $Z$  usually named the  $Z$ -score (of  $X$ ).

- If  $X \sim \mathcal{N}(0, 1)$ , then

$$Y = \mu + \sigma \cdot X \sim$$



# Note: Normal Distribution

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Y = X - \mu \sim \mathcal{N}(0, \sigma^2);$$

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$Z = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1)$$

so  $Z = \frac{X - \mu}{\sigma}$  is a Standard Normal Variable. This transformation is called a Standardization, and  $Z$  usually named the  $Z$ -score (of  $X$ ).

- If  $X \sim \mathcal{N}(0, 1)$ , then

$$Y = \mu + \sigma \cdot X \sim \mathcal{N}(\mu, \sigma^2)$$

# Functions of Random Variables: Making new R.V.s from the old ones

# Functions of Random Variables

Given a r.v.  $X$ , one can form new r.v.s by applying functions on  $X$ .

---

<sup>2</sup>(18+) Measurable!

# Functions of Random Variables

Given a r.v.  $X$ , one can form new r.v.s by applying functions on  $X$ .

**Example:** Say,  $X$  is the amount we will earn if selling 100 AMZN shares at the end of the year.

---

<sup>2</sup>(18+) Measurable!

# Functions of Random Variables

Given a r.v.  $X$ , one can form new r.v.s by applying functions on  $X$ .

**Example:** Say,  $X$  is the amount we will earn if selling 100 AMZN shares at the end of the year. Of course,  $X$  is a r.v.

---

<sup>2</sup>(18+) Measurable!

# Functions of Random Variables

Given a r.v.  $X$ , one can form new r.v.s by applying functions on  $X$ .

**Example:** Say,  $X$  is the amount we will earn if selling 100 AMZN shares at the end of the year. Of course,  $X$  is a r.v. Then the amount of the income tax we will pay,  $Y$ , will be a function of  $X$ .

---

<sup>2</sup>(18+) Measurable!

# Functions of Random Variables

Given a r.v.  $X$ , one can form new r.v.s by applying functions on  $X$ .

**Example:** Say,  $X$  is the amount we will earn if selling 100 AMZN shares at the end of the year. Of course,  $X$  is a r.v. Then the amount of the income tax we will pay,  $Y$ , will be a function of  $X$ . Say, if the tax rate is 10%, then  $Y = 0.1 \cdot X$ , and  $Y$  is again a r.v.

---

<sup>2</sup>(18+) Measurable!

# Functions of Random Variables

Given a r.v.  $X$ , one can form new r.v.s by applying functions on  $X$ .

**Example:** Say,  $X$  is the amount we will earn if selling 100 AMZN shares at the end of the year. Of course,  $X$  is a r.v. Then the amount of the income tax we will pay,  $Y$ , will be a function of  $X$ . Say, if the tax rate is 10%, then  $Y = 0.1 \cdot X$ , and  $Y$  is again a r.v.

Now, if  $X$  is a r.v., and  $g: \mathbb{R} \rightarrow \mathbb{R}$  is some function<sup>2</sup>, then the r.v.

$$Y = g(X)$$

is called the transformation of  $X$ .

---

<sup>2</sup>(18+) Measurable!



# Functions of Random Variables

Given a r.v.  $X$ , one can form new r.v.s by applying functions on  $X$ .

**Example:** Say,  $X$  is the amount we will earn if selling 100 AMZN shares at the end of the year. Of course,  $X$  is a r.v. Then the amount of the income tax we will pay,  $Y$ , will be a function of  $X$ . Say, if the tax rate is 10%, then  $Y = 0.1 \cdot X$ , and  $Y$  is again a r.v.

Now, if  $X$  is a r.v., and  $g: \mathbb{R} \rightarrow \mathbb{R}$  is some function<sup>2</sup>, then the r.v.

$$Y = g(X)$$

is called the transformation of  $X$ .

**Example:** For example,  $Y = X^3$ ,  $Y = \ln(X)$ ,  $Y = \frac{X}{1+X}$ ,  $Z = \sin(X)$ , ... are all (if defined, of course) new r.v.s obtained from the r.v.  $X$ .

---

<sup>2</sup>(18+) Measurable!

# Describing Transformations of $X$

Assume  $X$  is a r.v., and  $Y = g(X)$  is the transformation of  $X$ .

# Describing Transformations of $X$

Assume  $X$  is a r.v., and  $Y = g(X)$  is the transformation of  $X$ .  
The general problem here is

Describe the Distribution of  $Y$ , having the Distribution of  $X$ .

# Describing Transformations of $X$

Assume  $X$  is a r.v., and  $Y = g(X)$  is the transformation of  $X$ .  
The general problem here is

Describe the Distribution of  $Y$ , having the Distribution of  $X$ .

Let us try to see if you can guess:

- Assume  $X$  is Discrete Uniform. What can be said about  $Y = X^2$ ? Say, will  $Y$  be Discrete Uniform?

# Describing Transformations of $X$

Assume  $X$  is a r.v., and  $Y = g(X)$  is the transformation of  $X$ .  
The general problem here is

Describe the Distribution of  $Y$ , having the Distribution of  $X$ .

Let us try to see if you can guess:

- Assume  $X$  is Discrete Uniform. What can be said about  $Y = X^2$ ? Say, will  $Y$  be Discrete Uniform?
- Assume  $X \sim \text{Unif}[-2, 2]$ . What can be said about  $Y = 3X + 1$ ?

# Describing Transformations of $X$

Assume  $X$  is a r.v., and  $Y = g(X)$  is the transformation of  $X$ .  
The general problem here is

Describe the Distribution of  $Y$ , having the Distribution of  $X$ .

Let us try to see if you can guess:

- Assume  $X$  is Discrete Uniform. What can be said about  $Y = X^2$ ? Say, will  $Y$  be Discrete Uniform?
- Assume  $X \sim \text{Unif}[-2, 2]$ . What can be said about  $Y = 3X + 1$ ? What about  $Z = X^2$ ?

# Simulation of Transformed RVs: R Examples

## R Code

```
x <- runif(50000, min = -2, max = 2)
hist(x)
hist(x, freq = F, col = "cyan")
abline(h = 0.25, col = "red", lwd = 2)

y <- 3*x + 1
hist(y, freq = F, col = "lightblue") #shows uniform

z <- x^2
hist(z, freq = F, col = "lightcyan") #shows non-uniform!
```