

# CS 107, Probability, Spring 2019

## Lecture 06

Michael Poghosyan

AUA

30 January 2019

- Classical Probability Models: Finite Sample Spaces with Equally Likely Outcomes = Combinatorial Problems

## Birthday Problem

We have 36 participants in our group of Probability class, including the instructor. What is the probability that at least two participants share the same birthday, i.e., they were born on the same day and month (but maybe not in the same year)?

## Birthday Problem

We have 36 participants in our group of Probability class, including the instructor. What is the probability that at least two participants share the same birthday, i.e., they were born on the same day and month (but maybe not in the same year)?

Answer: the Probability is larger than 83%.

# Equally Likely Outcomes

Recall that when we have the Equally Likely Outcomes (or the uniform discrete distribution) model:

Outcome	$\omega_1$	$\omega_2$	$\omega_3$	$\dots$	$\omega_n$
Probability	$\frac{1}{n}$	$\frac{1}{n}$	$\frac{1}{n}$	$\dots$	$\frac{1}{n}$

then for any Event  $A$ ,

$$\mathbb{P}(A) =$$

# Equally Likely Outcomes

Recall that when we have the Equally Likely Outcomes (or the uniform discrete distribution) model:

Outcome	$\omega_1$	$\omega_2$	$\omega_3$	$\dots$	$\omega_n$
Probability	$\frac{1}{n}$	$\frac{1}{n}$	$\frac{1}{n}$	$\dots$	$\frac{1}{n}$

then for any Event  $A$ ,

$$\mathbb{P}(A) = \frac{\text{number of elements favorable for the event } A}{\text{total number of possible outcomes}}$$

$$= \frac{\#A}{\#\Omega} = \frac{\#A}{n}$$

# Example:

Before the Example - 52 Playing Cards Deck consists of:

- Suits:

# Example:

Before the Example - 52 Playing Cards Deck consists of:

- Suits:
  - **Hearts** = ♥,
  - **Diamonds** = ♦,
  - **Clubs** = ♣,
  - **Spades** = ♠.



# Example:

Before the Example - 52 Playing Cards Deck consists of:

- Suits:
  - **Hearts** = ♥,
  - **Diamonds** = ♦,
  - **Clubs** = ♣,
  - **Spades** = ♠.
- Cards: **Ace, King, Queen, Jack, 2, 3, 4, 5, 6, 7, 8, 9, 10.**

# Example:

Before the Example - 52 Playing Cards Deck consists of:

- Suits:
  - **Hearts** = ♥,
  - **Diamonds** = ♦,
  - **Clubs** = ♣,
  - **Spades** = ♠.
- Cards: **Ace, King, Queen, Jack, 2, 3, 4, 5, 6, 7, 8, 9, 10.**

Now, our Problem: We shuffle our deck and pick at random 2 cards. What is the probability that none of these cards will be hearts?

## Example: Cont'd

- **Problem:** We shuffle our deck and pick at random 2 cards. What is the probability that none of these cards will be hearts?

## Example: Cont'd

- **Problem:** We shuffle our deck and pick at random 2 cards. What is the probability that none of these cards will be hearts?
- Natural Assumption: the probability of choosing any card is the same

## Example: Cont'd

- **Problem:** We shuffle our deck and pick at random 2 cards. What is the probability that none of these cards will be hearts?
- Natural Assumption: the probability of choosing any card is the same = Equiprobable Case!


## Example: Cont'd

- **Problem:** We shuffle our deck and pick at random 2 cards. What is the probability that none of these cards will be hearts?
- Natural Assumption: the probability of choosing any card is the same = Equiprobable Case!
- **Case 1:** We take cards **without replacement**:

## Example: Cont'd

- **Problem:** We shuffle our deck and pick at random 2 cards. What is the probability that none of these cards will be hearts?
- Natural Assumption: the probability of choosing any card is the same = Equiprobable Case!
- **Case 1:** We take cards **without replacement**:


# of all Possible Outcomes:




# Example: Cont'd

- **Problem:** We shuffle our deck and pick at random 2 cards. What is the probability that none of these cards will be hearts?
- Natural Assumption: the probability of choosing any card is the same = Equiprobable Case!
- **Case 1:** We take cards **without replacement**:

# of all Possible Outcomes:



# of all Favorable Outcomes:







# Example: Cont'd

- **Problem:** We shuffle our deck and pick at random 2 cards. What is the probability that none of these cards will be hearts?
- Natural Assumption: the probability of choosing any card is the same = Equiprobable Case!
- **Case 1:** We take cards **without replacement**:

# of all Possible Outcomes:



# of all Favorable Outcomes:



$$\text{Probability} = \frac{\text{\#of favorable outcomes}}{\text{\#of all possible outcomes}} =$$

## Example: Cont'd 2

- **Case 1:** We take cards **without replacement**, almost the same approach in other way:

## Example: Cont'd 2

- **Case 1:** We take cards **without replacement**, almost the same approach in other way: clearly, our outcomes are pairs of cards.

## Example: Cont'd 2

- **Case 1:** We take cards **without replacement**, almost the same approach in other way: clearly, our outcomes are pairs of cards. Each pair is equally probable to be chosen (why?).

## Example: Cont'd 2

- **Case 1:** We take cards **without replacement**, almost the same approach in other way: clearly, our outcomes are pairs of cards. Each pair is equally probable to be chosen (why?). Then

$$\mathbb{P}(\text{no } \heartsuit \text{ in 2 cards}) = \frac{\# \text{pairs without a } \heartsuit}{\# \text{all pairs}} = \frac{\binom{39}{2}}{\binom{52}{2}}$$

## Example: Cont'd 2

- **Case 1:** We take cards **without replacement**, almost the same approach in other way: clearly, our outcomes are pairs of cards. Each pair is equally probable to be chosen (why?). Then

$$\mathbb{P}(\text{no } \heartsuit \text{ in 2 cards}) = \frac{\# \text{pairs without a } \heartsuit}{\# \text{all pairs}} = \frac{\binom{39}{2}}{\binom{52}{2}}$$

- What is the difference between these two approaches?

## Example: Cont'd 2

- **Case 1:** We take cards **without replacement**, almost the same approach in other way: clearly, our outcomes are pairs of cards. Each pair is equally probable to be chosen (why?). Then

$$\mathbb{P}(\text{no } \heartsuit \text{ in 2 cards}) = \frac{\# \text{pairs without a } \heartsuit}{\# \text{all pairs}} = \frac{\binom{39}{2}}{\binom{52}{2}}$$

- What is the difference between these two approaches?
- For the first case, we consider all **ordered** pairs:

$$\{(A\heartsuit, A\diamondsuit), (A\diamondsuit, A\heartsuit), (A\heartsuit, K\clubsuit), (K\clubsuit, A\heartsuit), \dots\}$$

## Example: Cont'd 2

- **Case 1:** We take cards **without replacement**, almost the same approach in other way: clearly, our outcomes are pairs of cards. Each pair is equally probable to be chosen (why?). Then

$$\mathbb{P}(\text{no } \heartsuit \text{ in 2 cards}) = \frac{\# \text{pairs without a } \heartsuit}{\# \text{all pairs}} = \frac{\binom{39}{2}}{\binom{52}{2}}$$

- What is the difference between these two approaches?
- For the first case, we consider all **ordered** pairs:

$$\{(A\heartsuit, A\diamondsuit), (A\diamondsuit, A\heartsuit), (A\heartsuit, K\clubsuit), (K\clubsuit, A\heartsuit), \dots\}$$

- For the second case, we consider all **unordered** pairs:

$$\{(A\heartsuit, A\diamondsuit), (A\heartsuit, K\clubsuit), \dots\}$$



## Example: Cont'd 2

- **Case 1:** We take cards **without replacement**, almost the same approach in other way: clearly, our outcomes are pairs of cards. Each pair is equally probable to be chosen (why?). Then

$$\mathbb{P}(\text{no } \heartsuit \text{ in 2 cards}) = \frac{\# \text{pairs without a } \heartsuit}{\# \text{all pairs}} = \frac{\binom{39}{2}}{\binom{52}{2}}$$

- What is the difference between these two approaches?
- For the first case, we consider all **ordered** pairs:

$$\{(A\heartsuit, A\diamondsuit), (A\diamondsuit, A\heartsuit), (A\heartsuit, K\clubsuit), (K\clubsuit, A\heartsuit), \dots\}$$

- For the second case, we consider all **unordered** pairs:

$$\{(A\heartsuit, A\diamondsuit), (A\heartsuit, K\clubsuit), \dots\}$$

- **Case 2:** We take cards **with replacement**

# Detailed Study of a Simpler Case

**Simplified Problem:** We choose two letters from  $\{a, b, c, d\}$ . What is the probability of having no  $a$  and no  $d$  in the chosen pair?

# Detailed Study of a Simpler Case

**Simplified Problem:** We choose two letters from  $\{a, b, c, d\}$ .  
What is the probability of having no  $a$  and no  $d$  in the chosen pair?

Seems simple, but ...

# Detailed Study of a Simpler Case

**Simplified Problem:** We choose two letters from  $\{a, b, c, d\}$ . What is the probability of having no  $a$  and no  $d$  in the chosen pair?

Seems simple, but ...

**Case 1: choosing without replacement**

# Detailed Study of a Simpler Case

**Simplified Problem:** We choose two letters from  $\{a, b, c, d\}$ . What is the probability of having no  $a$  and no  $d$  in the chosen pair?

Seems simple, but ...

**Case 1: choosing without replacement**

- Ordered Case

# Detailed Study of a Simpler Case

**Simplified Problem:** We choose two letters from  $\{a, b, c, d\}$ . What is the probability of having no  $a$  and no  $d$  in the chosen pair?

Seems simple, but ...

## Case 1: choosing without replacement

- Ordered Case
- Unordered Case

# Detailed Study of a Simpler Case

**Simplified Problem:** We choose two letters from  $\{a, b, c, d\}$ . What is the probability of having no  $a$  and no  $d$  in the chosen pair?

Seems simple, but ...

**Case 1: choosing without replacement**

- Ordered Case
- Unordered Case

**Case 2: choosing with replacement**

# Detailed Study of a Simpler Case

**Simplified Problem:** We choose two letters from  $\{a, b, c, d\}$ . What is the probability of having no  $a$  and no  $d$  in the chosen pair?

Seems simple, but ...

## Case 1: choosing without replacement

- Ordered Case
- Unordered Case

## Case 2: choosing with replacement

- Ordered Case



# Detailed Study of a Simpler Case

**Simplified Problem:** We choose two letters from  $\{a, b, c, d\}$ . What is the probability of having no  $a$  and no  $d$  in the chosen pair?

Seems simple, but ...

## Case 1: choosing without replacement

- Ordered Case
- Unordered Case

## Case 2: choosing with replacement

- Ordered Case
- Unordered Case

# Moral from the previous Example:

# Moral from the previous Example:

- When considering a combinatorial probability problem, try to think first what are your outcomes and what is your Sample Space?

# Moral from the previous Example:

- When considering a combinatorial probability problem, try to think first what are your outcomes and what is your Sample Space?
- If your Sample Space consists of *unordered* pairs, then your Event under the interest also needs to consist of *unordered* pairs.

# Moral from the previous Example:

- When considering a combinatorial probability problem, try to think first what are your outcomes and what is your Sample Space?
- If your Sample Space consists of *unordered* pairs, then your Event under the interest also needs to consist of *unordered* pairs. And if your Sample Space consists of *ordered* pairs, then your Event under the interest also needs to consist of *ordered* pairs.

# Moral from the previous Example:

# Moral from the previous Example:

- If you consider an Experiment of drawing an item several times (say, 2 times) *without replacement*, then Unorderd/Ordered cases, if they are appropriate, will give the same Probability (if calculated correctly), because each outcome  $(a, b)$  in your Unordered Sample Space corresponds exactly to 2 outcomes in the Ordered Space:  $(a, b)$  and  $(b, a)$ .

# Moral from the previous Example:

- If you consider an Experiment of drawing an item several times (say, 2 times) *without replacement*, then Unorderd/Ordered cases, if they are appropriate, will give the same Probability (if calculated correctly), because each outcome  $(a, b)$  in your Unordered Sample Space corresponds exactly to 2 outcomes in the Ordered Space:  $(a, b)$  and  $(b, a)$ . That is, the no. of elements in the Favorable Event (Sample Space) in the Unordered case will be exactly the half of the Favorable Event (Sample Space) in the Ordered case.



# Moral from the previous Example:

- If you consider an Experiment of drawing an item several times (say, 2 times) *without replacement*, then Unorderd/Ordered cases, if they are appropriate, will give the same Probability (if calculated correctly), because each outcome  $(a, b)$  in your Unordered Sample Space corresponds exactly to 2 outcomes in the Ordered Space:  $(a, b)$  and  $(b, a)$ . That is, the no. of elements in the Favorable Event (Sample Space) in the Unordered case will be exactly the half of the Favorable Event (Sample Space) in the Ordered case. And, say, if you are drawing 3 times, you will have for each triple  $(a, b, c)$  in your Unordered SS 6 possible triples in the Ordered triples Sample Space:  $(a, b, c), (a, c, b), (b, a, c), \dots$

# Moral from the previous Example:

# Moral from the previous Example:

- If you consider an Experiment of drawing an item several times (say, 2 times) *with replacement*, then the situation with Unordered/Ordered cases will be completely different. This is because not for each pair in the Unordered Space we will have 2 pairs in the Ordered Space.

# Moral from the previous Example:

- If you consider an Experiment of drawing an item several times (say, 2 times) *with replacement*, then the situation with Unordered/Ordered cases will be completely different. This is because not for each pair in the Unordered Space we will have 2 pairs in the Ordered Space. Say, for the pair of the form  $(a, a)$  (this is a possible outcome, since we draw with replacement), we will not have 2 pairs in the Ordered Space - we will have only the same pair  $(a, a)$ .

## Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

## Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 1: Without replacement - we are not allowed to use the same digit again**

## Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 1: Without replacement - we are not allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{12, 13, 23\}$

# Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 1: Without replacement - we are not allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{12, 13, 23\}$
  - Ordered Sample Space =  $\{12, 13, 21, 23, 31, 32\}$



# Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 1: Without replacement - we are not allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{12, 13, 23\}$
  - Ordered Sample Space =  $\{12, 13, 21, 23, 31, 32\}$
  - Our Event in The UnOrdered SS =  $\{12, 23\}$

# Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 1: Without replacement - we are not allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{12, 13, 23\}$
  - Ordered Sample Space =  $\{12, 13, 21, 23, 31, 32\}$
  - Our Event in The UnOrdered SS =  $\{12, 23\}$
  - Our Event in The Ordered SS =  $\{12, 21, 23, 32\}$

# Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 1: Without replacement - we are not allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{12, 13, 23\}$
  - Ordered Sample Space =  $\{12, 13, 21, 23, 31, 32\}$
  - Our Event in The UnOrdered SS =  $\{12, 23\}$
  - Our Event in The Ordered SS =  $\{12, 21, 23, 32\}$
  - $\mathbb{P}(\text{Contains an Even digit}) = \frac{2}{3} = \frac{4}{6}$ .

## Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

## Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 2: With replacement - we are allowed to use the same digit again**

# Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 2: With replacement - we are allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{11, 12, 13, 22, 23, 33\}$

# Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 2: With replacement - we are allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{11, 12, 13, 22, 23, 33\}$
  - Ordered Sample Space =  $\{11, 12, 13, 21, 22, 23, 31, 32, 33\}$

# Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 2: With replacement - we are allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{11, 12, 13, 22, 23, 33\}$
  - Ordered Sample Space =  $\{11, 12, 13, 21, 22, 23, 31, 32, 33\}$
  - Our Event in The UnOrdered SS =  $\{12, 22, 23\}$



# Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 2: With replacement - we are allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{11, 12, 13, 22, 23, 33\}$
  - Ordered Sample Space =  $\{11, 12, 13, 21, 22, 23, 31, 32, 33\}$
  - Our Event in The UnOrdered SS =  $\{12, 22, 23\}$
  - Our Event in The Ordered SS =  $\{12, 21, 22, 23, 32\}$

# Another Example:

**Problem:** Form all 2-digit numbers from  $\{1, 2, 3\}$ . Consider the Event when the obtained number contains an even digit. Calculate the Probability of that Event. Consider with/without replacement cases.

- **Case 2: With replacement - we are allowed to use the same digit again**
  - UnOrdered Sample Space =  $\{11, 12, 13, 22, 23, 33\}$
  - Ordered Sample Space =  $\{11, 12, 13, 21, 22, 23, 31, 32, 33\}$
  - Our Event in The UnOrdered SS =  $\{12, 22, 23\}$
  - Our Event in The Ordered SS =  $\{12, 21, 22, 23, 32\}$
  - $\mathbb{P}(\text{Contains an Even digit}) =$

# Example:

- **Problem:** We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?

# Example:

- **Problem:** We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?
- Clear assumption: Choosing each ball is equiprobable

# Example:

- **Problem:** We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?
- Clear assumption: Choosing each ball is equiprobable
- **Case 1:** We take balls without a replacement:

# Example:

- **Problem:** We have a box containing 100 balls, from which 25 are white and the rest are black. We choose at random 12 balls. What is the probability of having exactly 4 white balls chosen?
- Clear assumption: Choosing each ball is equiprobable
- **Case 1:** We take balls without a replacement:
- **Case 2:** We take balls with a replacement: