

# AUA CS 108, Statistics, Fall 2019

## Lecture 20

Michael Poghosyan

YSU, AUA

[michael@ysu.am](mailto:michael@ysu.am), [mpoghosyan@aua.am](mailto:mpoghosyan@aua.am)

09 Oct 2019

# Contents

- ▶ Properties of Estimators: MSE
- ▶ Bias, Biased and Unbiased Estimates
- ▶ Consistent Estimators

## Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.

## Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.
- ▶ What is a Statistics,  $v_3$ ?

## Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.
- ▶ What is a Statistics,  $v_3$ ?
- ▶ What is an Estimator?

## Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.
- ▶ What is a Statistics,  $v_3$ ?
- ▶ What is an Estimator?
- ▶ What is an Estimate?

# Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.
- ▶ What is a Statistics,  $v_3$ ?
- ▶ What is an Estimator?
- ▶ What is an Estimate?
- ▶ Give some Estimators for  $Geom(p)$  parameter  $p$ .

## General Setting for the Point Estimation

The general setting for the Point Estimation is the following:



## General Setting for the Point Estimation

The general setting for the Point Estimation is the following:

- ▶ We have a Dataset:  $x : x_1, x_2, \dots, x_n$ ;

# General Setting for the Point Estimation

The general setting for the Point Estimation is the following:

- ▶ We have a Dataset:  $x : x_1, x_2, \dots, x_n$ ;
- ▶ We model the process that generated  $x$  by the Parametric Family of Distributions  $\{\mathcal{F}_\theta, \theta \in \Theta\}$ ;

# General Setting for the Point Estimation

The general setting for the Point Estimation is the following:

- ▶ We have a Dataset:  $x : x_1, x_2, \dots, x_n$ ;
- ▶ We model the process that generated  $x$  by the Parametric Family of Distributions  $\{\mathcal{F}_\theta, \theta \in \Theta\}$ ;
- ▶ We assume the Dataset above is one possible realization of the Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta \quad \text{for some } \theta \in \Theta;$$

## General Setting for the Point Estimation

The general setting for the Point Estimation is the following:

- ▶ We have a Dataset:  $x : x_1, x_2, \dots, x_n$ ;
- ▶ We model the process that generated  $x$  by the Parametric Family of Distributions  $\{\mathcal{F}_\theta, \theta \in \Theta\}$ ;
- ▶ We assume the Dataset above is one possible realization of the Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta \quad \text{for some } \theta \in \Theta;$$

**Problem:** Estimate the unknown Parameter  $\theta$ .

## General Setting for the Point Estimation

The general setting for the Point Estimation is the following:

- ▶ We have a Dataset:  $x : x_1, x_2, \dots, x_n$ ;
- ▶ We model the process that generated  $x$  by the Parametric Family of Distributions  $\{\mathcal{F}_\theta, \theta \in \Theta\}$ ;
- ▶ We assume the Dataset above is one possible realization of the Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta \quad \text{for some } \theta \in \Theta;$$

**Problem:** Estimate the unknown Parameter  $\theta$ .

To estimate  $\theta$ ,

- ▶ We find a good **Estimator**  $\hat{\theta}$ ;

# General Setting for the Point Estimation

The general setting for the Point Estimation is the following:

- ▶ We have a Dataset:  $x : x_1, x_2, \dots, x_n$ ;
- ▶ We model the process that generated  $x$  by the Parametric Family of Distributions  $\{\mathcal{F}_\theta, \theta \in \Theta\}$ ;
- ▶ We assume the Dataset above is one possible realization of the Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta \quad \text{for some } \theta \in \Theta;$$

**Problem:** Estimate the unknown Parameter  $\theta$ .

To estimate  $\theta$ ,

- ▶ We find a good **Estimator**  $\hat{\theta}$ ;
- ▶ We plug the Dataset values into the Estimator to get the **Estimate** for  $\theta$

## Properties of Estimators

## Properties of Estimators

Story about Good and Bad Estimators 😊



# Risk, Mean Squared Error of the Estimator

From the last lecture:

**Definition:** The **Mean Squared Error** or the **Quadratic Risk** of the estimator  $\hat{\theta}_n$  of  $\theta$  is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

# Risk, Mean Squared Error of the Estimator

From the last lecture:

**Definition:** The **Mean Squared Error** or the **Quadratic Risk** of the estimator  $\hat{\theta}_n$  of  $\theta$  is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

**Definition:** We say that the estimator  $\hat{\theta}_n^1$  of  $\theta$  is **preferable** to  $\hat{\theta}_n^2$ , another estimator of  $\theta$ , if

$$MSE(\hat{\theta}_n^1, \theta) \leq MSE(\hat{\theta}_n^2, \theta), \quad \forall \theta \in \Theta,$$

and there exists a  $\theta$  s.t.  $MSE(\hat{\theta}_n^1, \theta) < MSE(\hat{\theta}_n^2, \theta)$ .

## Example

**Example:** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda),$$

and we want to Estimate the unknown  $\lambda$ .

## Example

**Example:** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda),$$

and we want to Estimate the unknown  $\lambda$ . We want to use one of the following Estimators:

$$\hat{\lambda}_1 = X_1 \quad \text{or} \quad \hat{\lambda}_2 = \frac{X_1 + X_n}{2}.$$

## Example

**Example:** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda),$$

and we want to Estimate the unknown  $\lambda$ . We want to use one of the following Estimators:

$$\hat{\lambda}_1 = X_1 \quad \text{or} \quad \hat{\lambda}_2 = \frac{X_1 + X_n}{2}.$$

**Question:** Calculate the Risks of  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$ . Compare these two Estimators: which of  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$  is preferable?

## Example

**Example:** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda),$$

and we want to Estimate the unknown  $\lambda$ . We want to use one of the following Estimators:

$$\hat{\lambda}_1 = X_1 \quad \text{or} \quad \hat{\lambda}_2 = \frac{X_1 + X_n}{2}.$$

**Question:** Calculate the Risks of  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$ . Compare these two Estimators: which of  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$  is preferable?

**Solution:** OTB: Use the fact that for  $X \sim \text{Pois}(\lambda)$ ,

$$\mathbb{E}(X) = \lambda = \text{Var}(X).$$

## Best MSE Estimators

Now, having the idea of the Error of estimation for an Estimator, we can try to find the best Estimator in the sense of MSE, i.e., we can try to solve

Find  $\hat{\theta}$  that minimizes  $MSE(\hat{\theta}, \theta)$ .

## Best MSE Estimators

Now, having the idea of the Error of estimation for an Estimator, we can try to find the best Estimator in the sense of MSE, i.e., we can try to solve

Find  $\hat{\theta}$  that minimizes  $MSE(\hat{\theta}, \theta)$ .

Unfortunately, this Minimization problem has no solution in general.



## Best MSE Estimators

Now, having the idea of the Error of estimation for an Estimator, we can try to find the best Estimator in the sense of MSE, i.e., we can try to solve

Find  $\hat{\theta}$  that minimizes  $MSE(\hat{\theta}, \theta)$ .

Unfortunately, this Minimization problem has no solution in general.

So we move forward considering other Properties of Estimators making them useful in the estimation process.

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

**Definition:** We say that our Estimator  $\hat{\theta}$  (of the parameter  $\theta$ ) is **Unbiased**, if

$$\text{Bias}(\hat{\theta}, \theta) = 0 \quad \text{for any} \quad \theta \in \Theta.$$

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

**Definition:** We say that our Estimator  $\hat{\theta}$  (of the parameter  $\theta$ ) is **Unbiased**, if

$$\text{Bias}(\hat{\theta}, \theta) = 0 \quad \text{for any} \quad \theta \in \Theta.$$

If  $\hat{\theta}$  is not Unbiased, then we say that it is **Biased**.

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

**Definition:** We say that our Estimator  $\hat{\theta}$  (of the parameter  $\theta$ ) is **Unbiased**, if

$$\text{Bias}(\hat{\theta}, \theta) = 0 \quad \text{for any} \quad \theta \in \Theta.$$

If  $\hat{\theta}$  is not Unbiased, then we say that it is **Biased**.

In other words, the Estimator  $\hat{\theta}$  is **Unbiased**, if

$$\mathbb{E}_{\theta}(\hat{\theta}) = \theta \quad \text{for any} \quad \theta \in \Theta.$$

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

**Definition:** We say that our Estimator  $\hat{\theta}$  (of the parameter  $\theta$ ) is **Unbiased**, if

$$\text{Bias}(\hat{\theta}, \theta) = 0 \quad \text{for any} \quad \theta \in \Theta.$$

If  $\hat{\theta}$  is not Unbiased, then we say that it is **Biased**.

In other words, the Estimator  $\hat{\theta}$  is **Unbiased**, if

$$\mathbb{E}_{\theta}(\hat{\theta}) = \theta \quad \text{for any} \quad \theta \in \Theta.$$

**Note:** We require the above hold **for any parameter value** since, if, say, it is correct for *some* values of  $\theta$ , then it can happen that the true value of our unknown  $\theta$  is exactly that value, for which we do not have the equality.

# Bias and Unbiasedness

- ▶ The idea of Unbiased Estimator is the following: if we will calculate Estimates many-many times using our Unbiased Estimator, and then average the obtained Estimates, we will obtain the (almost exact, exact when many-many  $\rightarrow \infty$ ) value of our Parameter.

# Bias and Unbiasedness

- ▶ The idea of Unbiased Estimator is the following: if we will calculate Estimates many-many times using our Unbiased Estimator, and then average the obtained Estimates, we will obtain the (almost exact, exact when many-many  $\rightarrow \infty$ ) value of our Parameter.
- ▶ So we can say about an Unbiased Estimator as: **In average, it is Exact**



# Bias and Unbiasedness

- ▶ The idea of Unbiased Estimator is the following: if we will calculate Estimates many-many times using our Unbiased Estimator, and then average the obtained Estimates, we will obtain the (almost exact, exact when many-many  $\rightarrow \infty$ ) value of our Parameter.
- ▶ So we can say about an Unbiased Estimator as: **In average, it is Exact**
- ▶ Bias can be interpreted, in some sense, as the *accuracy* of the Estimator.

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population.

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution.

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution. So our unknown Parameter here is  $\mu$ .

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution. So our unknown Parameter here is  $\mu$ .

We use size  $n$  Random Sample to Model the situation:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\mu.$$

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution. So our unknown Parameter here is  $\mu$ .

We use size  $n$  Random Sample to Model the situation:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\mu.$$

Now, we choose several Estimators to estimate  $\mu$ :

$$\begin{aligned}\hat{\mu}_1 &= X_1, & \hat{\mu}_2 &= \frac{X_1 + X_3}{2}, & \hat{\mu}_3 &= \frac{X_1 + X_4}{10}, \\ \hat{\mu}_4 &= \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.\end{aligned}$$

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution. So our unknown Parameter here is  $\mu$ .

We use size  $n$  Random Sample to Model the situation:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\mu.$$

Now, we choose several Estimators to estimate  $\mu$ :

$$\hat{\mu}_1 = X_1, \quad \hat{\mu}_2 = \frac{X_1 + X_3}{2}, \quad \hat{\mu}_3 = \frac{X_1 + X_4}{10},$$

$$\hat{\mu}_4 = \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

**Problem:** Check if each Estimator is Biased or Unbiased.

**Solution:** OTB