

AUA CS 108, Statistics, Fall 2019

Lecture 37

Michael Poghosyan

YSU, AUA

michael@ysu.am, mpoghosyan@aua.am

18 Nov 2019

Contents

- ▶ Hypothesis Testing: Error Types, Significance and Power
- ▶ Designing a Test: General Procedure
- ▶ Z-Test

Last Lecture ReCap

- ▶ How to choose the Null Hypothesis?

From the last lecture: Type I and II errors

Assume we are Testing the Hypothesis

$$\mathcal{H}_0 \quad \text{vs} \quad \mathcal{H}_1.$$

Then the following cases can happen:

Test Decision \ Reality	\mathcal{H}_0 is True	\mathcal{H}_0 is False (i.e., \mathcal{H}_1 is True)
Reject \mathcal{H}_0	Type I Error (False Positive)	Correct Decision (True Negative)
Do Not Reject \mathcal{H}_0	Correct Decision (True Positive)	Type II Error (False Negative)

Significance and Power of Test

- ▶ The Probability of Type I Error is called the **Significance Level** of our Test, and is denoted by α :

Significance and Power of Test

- ▶ The Probability of Type I Error is called the **Significance Level** of our Test, and is denoted by α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}).$$

Significance and Power of Test

- ▶ The Probability of Type I Error is called the **Significance Level** of our Test, and is denoted by α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}).$$

- ▶ The Probability of Type II Error is denoted by β :

$$\beta = \mathbb{P}(\text{Type II Error}) = \mathbb{P}(\text{Retain } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is False}).$$

Significance and Power of Test

- ▶ The Probability of Type I Error is called the **Significance Level** of our Test, and is denoted by α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}).$$

- ▶ The Probability of Type II Error is denoted by β :

$$\beta = \mathbb{P}(\text{Type II Error}) = \mathbb{P}(\text{Retain } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is False}).$$

$(1 - \beta)$ is called the **Power** of the Test.

Significance and Power of Test

- ▶ The Probability of Type I Error is called the **Significance Level** of our Test, and is denoted by α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}).$$

- ▶ The Probability of Type II Error is denoted by β :

$$\beta = \mathbb{P}(\text{Type II Error}) = \mathbb{P}(\text{Retain } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is False}).$$

$(1 - \beta)$ is called the **Power** of the Test.

It is easy to see that

$$\text{Power} = 1 - \beta = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is False}).$$

Significance and Power

Here are the Probabilities of correct/incorrect decisions for a Hypothesis testing, on a Table:

Significance and Power

Here are the Probabilities of correct/incorrect decisions for a Hypothesis testing, on a Table:

Probabilities of Correct/InCorrect Decisions:

Test Decision \ Reality	\mathcal{H}_0 is True	\mathcal{H}_0 is False (i.e., \mathcal{H}_1 is True)
Reject \mathcal{H}_0	$\alpha = \mathbf{Significance}$	$1 - \beta = \mathbf{Power}$
Do Not Reject \mathcal{H}_0	$1 - \alpha$	β

Choosing a Test

Test Choice: We want to have a Test with small Error Probabilities, i.e., with small α and β .

Choosing a Test

Test Choice: We want to have a Test with small Error Probabilities, i.e., with small α and β . Unfortunately, making them both small is not always possible. So we first think in controlling α : **our goal will be to choose the desired α and to have a Test of Significance Level α .**

Choosing a Test

Test Choice: We want to have a Test with small Error Probabilities, i.e., with small α and β . Unfortunately, making them both small is not always possible. So we first think in controlling α : **our goal will be to choose the desired α and to have a Test of Significance Level α .**

Common values of α are 0.05, 0.01 and 0.1 (corresponding to 95%, 99% and 90% Confidence Level!).

Example

Example: Assume we have a Patient in a Hospital, and we want to see if he is ill or not (with some illness). Then the Null Hypothesis is:

Example

Example: Assume we have a Patient in a Hospital, and we want to see if he is ill or not (with some illness). Then the Null Hypothesis is: **Patient is healthy**. Our Test is *the Doctor*.

Example

Example: Assume we have a Patient in a Hospital, and we want to see if he is ill or not (with some illness). Then the Null Hypothesis is: **Patient is healthy**. Our Test is *the Doctor*.

- ▶ What is means that the Significance level of our Test, α , is small ?

Example

Example: Assume we have a Patient in a Hospital, and we want to see if he is ill or not (with some illness). Then the Null Hypothesis is: **Patient is healthy**. Our Test is *the Doctor*.

- ▶ What it means that the Significance level of our Test, α , is small ?
- ▶ What it means that the Power of our Test, $1 - \beta$, is high ?

Hypo Testing: Constructing a Test

To test a Hypothesis, one is following the following steps ☺:

Step 1: Choosing a Model: We assume our Data comes from a Parametric Model $\mathcal{F}_\theta, \theta \in \Theta$

Hypo Testing: Constructing a Test

To test a Hypothesis, one is following the following steps ☺:

Step 1: Choosing a Model: We assume our Data comes from a Parametric Model \mathcal{F}_θ , $\theta \in \Theta$

Step 2: We State the Hypotheses: we take Θ_0 and Θ_1 such that $\Theta = \Theta_0 \cup \Theta_1$ and $\Theta_0 \cap \Theta_1 = \emptyset$, and state the Hypothesis:

$$\mathcal{H}_0 : \theta \in \Theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta \in \Theta_1$$

Hypo Testing: Constructing a Test

To test a Hypothesis, one is following the following steps \smile :

Step 1: Choosing a Model: We assume our Data comes from a Parametric Model \mathcal{F}_θ , $\theta \in \Theta$

Step 2: We State the Hypotheses: we take Θ_0 and Θ_1 such that $\Theta = \Theta_0 \cup \Theta_1$ and $\Theta_0 \cap \Theta_1 = \emptyset$, and state the Hypothesis:

$$\mathcal{H}_0 : \theta \in \Theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta \in \Theta_1$$

Step 3: We choose the desirable Significance Level: we choose α (small enough, say, $\alpha = 0.05$).

¹If n is free to choose, then we can find it by taking the desirable Power of the Test.

Hypo Testing: Constructing a Test

To test a Hypothesis, one is following the following steps ☺:

Step 1: Choosing a Model: We assume our Data comes from a Parametric Model \mathcal{F}_θ , $\theta \in \Theta$

Step 2: We State the Hypotheses: we take Θ_0 and Θ_1 such that $\Theta = \Theta_0 \cup \Theta_1$ and $\Theta_0 \cap \Theta_1 = \emptyset$, and state the Hypothesis:

$$\mathcal{H}_0 : \theta \in \Theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta \in \Theta_1$$

Step 3: We choose the desirable Significance Level: we choose α (small enough, say, $\alpha = 0.05$).

Step 4: Assume we know how many Observations we will have¹, say, n .

¹If n is free to choose, then we can find it by taking the desirable Power of the Test.

Hypo Testing: Constructing a Test

To test a Hypothesis, one is following the following steps ☺:

Step 1: Choosing a Model: We assume our Data comes from a Parametric Model \mathcal{F}_θ , $\theta \in \Theta$

Step 2: We State the Hypotheses: we take Θ_0 and Θ_1 such that $\Theta = \Theta_0 \cup \Theta_1$ and $\Theta_0 \cap \Theta_1 = \emptyset$, and state the Hypothesis:

$$\mathcal{H}_0 : \theta \in \Theta_0 \quad \text{vs} \quad \mathcal{H}_1 : \theta \in \Theta_1$$

Step 3: We choose the desirable Significance Level: we choose α (small enough, say, $\alpha = 0.05$).

Step 4: Assume we know how many Observations we will have¹, say, n . Then we take a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta.$$

¹If n is free to choose, then we can find it by taking the desirable Power of the Test.

Hypo Testing: Constructing a Test

Step 5: Next, we choose a Test Statistics,

$$T = T(X_1, X_2, \dots, X_n)$$

such that

Hypo Testing: Constructing a Test

Step 5: Next, we choose a Test Statistics,

$$T = T(X_1, X_2, \dots, X_n)$$

such that

- ▶ We can calculate it, if we will have Observations;

Hypo Testing: Constructing a Test

Step 5: Next, we choose a Test Statistics,

$$T = T(X_1, X_2, \dots, X_n)$$

such that

- ▶ We can calculate it, if we will have Observations;
- ▶ the Distribution (or, at least, the Asymptotic Distribution) of T is known **under** \mathcal{H}_0 .

Hypo Testing: Constructing a Test

Step 5: Next, we choose a Test Statistics,

$$T = T(X_1, X_2, \dots, X_n)$$

such that

- ▶ We can calculate it, if we will have Observations;
- ▶ the Distribution (or, at least, the Asymptotic Distribution) of T is known **under** \mathcal{H}_0 .

Step 6: Then, we specify the **Rejection Region (RR)**, usually of the form

$$RR = \{|T| > c\} \quad \text{or} \quad RR = \{T > c\} \quad \text{or} \quad RR = \{T < c\}.$$

Hypo Testing: Constructing a Test

Step 5: Next, we choose a Test Statistics,

$$T = T(X_1, X_2, \dots, X_n)$$

such that

- ▶ We can calculate it, if we will have Observations;
- ▶ the Distribution (or, at least, the Asymptotic Distribution) of T is known **under** \mathcal{H}_0 .

Step 6: Then, we specify the **Rejection Region (RR)**, usually of the form

$$RR = \{|T| > c\} \quad \text{or} \quad RR = \{T > c\} \quad \text{or} \quad RR = \{T < c\}.$$

Here c is called **the Critical Value** of the Test. We will choose it (soon).

Hypo Testing: Constructing a Test

Step 5: Next, we choose a Test Statistics,

$$T = T(X_1, X_2, \dots, X_n)$$

such that

- ▶ We can calculate it, if we will have Observations;
- ▶ the Distribution (or, at least, the Asymptotic Distribution) of T is known **under** \mathcal{H}_0 .

Step 6: Then, we specify the **Rejection Region (RR)**, usually of the form

$$RR = \{|T| > c\} \quad \text{or} \quad RR = \{T > c\} \quad \text{or} \quad RR = \{T < c\}.$$

Here c is called **the Critical Value** of the Test. We will choose it (soon). We Reject \mathcal{H}_0 if and only if $T \in RR$.

Hypo Testing: Constructing a Test

Now, at this step, we need to choose the Critical Value c .

Hypo Testing: Constructing a Test

Now, at this step, we need to choose the Critical Value c . We choose it to have the desirable Significance Level:

$$\mathbb{P}(\text{Type I Error}) = \alpha,$$

Hypo Testing: Constructing a Test

Now, at this step, we need to choose the Critical Value c . We choose it to have the desirable Significance Level:

$$\mathbb{P}(\text{Type I Error}) = \alpha,$$

i.e.,

$$\alpha = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) = \mathbb{P}(T \in RR \mid \theta \in \Theta_0).$$

Hypo Testing: Constructing a Test

Now, at this step, we need to choose the Critical Value c . We choose it to have the desirable Significance Level:

$$\mathbb{P}(\text{Type I Error}) = \alpha,$$

i.e.,

$$\alpha = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) = \mathbb{P}(T \in RR \mid \theta \in \Theta_0).$$

And usually, c is some order quantile of the Distribution of T .

Hypo Testing: Constructing a Test

Now, at this step, we need to choose the Critical Value c . We choose it to have the desirable Significance Level:

$$\mathbb{P}(\text{Type I Error}) = \alpha,$$

i.e.,

$$\alpha = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) = \mathbb{P}(T \in RR \mid \theta \in \Theta_0).$$

And usually, c is some order quantile of the Distribution of T .

Step 7: We make an observation, x_1, x_2, \dots, x_n , calculate the Test Statistics at that Observation, $T = T(x_1, \dots, x_n)$.

Hypo Testing: Constructing a Test

Now, at this step, we need to choose the Critical Value c . We choose it to have the desirable Significance Level:

$$\mathbb{P}(\text{Type I Error}) = \alpha,$$

i.e.,

$$\alpha = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) = \mathbb{P}(T \in RR \mid \theta \in \Theta_0).$$

And usually, c is some order quantile of the Distribution of T .

Step 7: We make an observation, x_1, x_2, \dots, x_n , calculate the Test Statistics at that Observation, $T = T(x_1, \dots, x_n)$. Now, we will

- Reject \mathcal{H}_0 , if $T(x_1, \dots, x_n) \in RR$;

Hypo Testing: Constructing a Test

Now, at this step, we need to choose the Critical Value c . We choose it to have the desirable Significance Level:

$$\mathbb{P}(\text{Type I Error}) = \alpha,$$

i.e.,

$$\alpha = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) = \mathbb{P}(T \in RR \mid \theta \in \Theta_0).$$

And usually, c is some order quantile of the Distribution of T .

Step 7: We make an observation, x_1, x_2, \dots, x_n , calculate the Test Statistics at that Observation, $T = T(x_1, \dots, x_n)$. Now, we will

- ▶ Reject \mathcal{H}_0 , if $T(x_1, \dots, x_n) \in RR$;
- ▶ Not Reject \mathcal{H}_0 , if $T(x_1, \dots, x_n) \notin RR$.

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known;

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

► Case 1: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

- ▶ Case 1: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$
- ▶ Case 2: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

► Case 1: $\mathcal{H}_0 : \mu = \mu_0$ *vs* $\mathcal{H}_1 : \mu \neq \mu_0$

► Case 2: $\mathcal{H}_0 : \mu = \mu_0$ *vs* $\mathcal{H}_1 : \mu > \mu_0$

► Case 3: $\mathcal{H}_0 : \mu = \mu_0$ *vs* $\mathcal{H}_1 : \mu < \mu_0$

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

► Case 1: $\mathcal{H}_0 : \mu = \mu_0$ *vs* $\mathcal{H}_1 : \mu \neq \mu_0$

► Case 2: $\mathcal{H}_0 : \mu = \mu_0$ *vs* $\mathcal{H}_1 : \mu > \mu_0$

► Case 3: $\mathcal{H}_0 : \mu = \mu_0$ *vs* $\mathcal{H}_1 : \mu < \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

► Case 1: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

► Case 2: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

► Case 3: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Random Sample: We take $X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$;

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

► Case 1: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

► Case 2: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

► Case 3: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Random Sample: We take $X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$;

Test Statistics: Idea:

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

► Case 1: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

► Case 2: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

► Case 3: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Random Sample: We take $X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$;

Test Statistics: Idea:

► what we have to approximate our Parameter (Estimate) μ ?

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

► Case 1: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

► Case 2: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

► Case 3: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Random Sample: We take $X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$;

Test Statistics: Idea:

- what we have to approximate our Parameter (Estimate) μ ? - Easy: \bar{X} ;

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

- ▶ Case 1: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$
- ▶ Case 2: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$
- ▶ Case 3: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Random Sample: We take $X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$;

Test Statistics: Idea:

- ▶ what we have to approximate our Parameter (Estimate) μ ? - Easy: \bar{X} ;
- ▶ In which case we will not believe that $\mu = \mu_0$? - Piece of Cake:

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

- ▶ Case 1: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$
- ▶ Case 2: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$
- ▶ Case 3: $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Random Sample: We take $X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$;

Test Statistics: Idea:

- ▶ what we have to approximate our Parameter (Estimate) μ ? - Easy: \bar{X} ;
- ▶ In which case we will not believe that $\mu = \mu_0$? - Piece of Cake: if \bar{X} will be far from μ_0

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z =$$

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

²Note the difference from the Pivot Z : when constructing a CI for μ , we were taking Z with μ .

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Let us check² that it satisfies the requirements:

²Note the difference from the Pivot Z : when constructing a CI for μ , we were taking Z with μ .

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Let us check² that it satisfies the requirements:

- If we will have the Observation, we can calculate it, for sure;

²Note the difference from the Pivot Z : when constructing a CI for μ , we were taking Z with μ .

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Let us check² that it satisfies the requirements:

- ▶ If we will have the Observation, we can calculate it, for sure;
- ▶ If the Null Hypo is True, then $\mu = \mu_0$. Recall what is the connection between our Statistics, Z and μ :

²Note the difference from the Pivot Z : when constructing a CI for μ , we were taking Z with μ .

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Let us check² that it satisfies the requirements:

- ▶ If we will have the Observation, we can calculate it, for sure;
- ▶ If the Null Hypo is True, then $\mu = \mu_0$. Recall what is the connection between our Statistics, Z and μ : the Distribution of X_k in Z is $\mathcal{N}(\mu, \sigma^2)$.

²Note the difference from the Pivot Z : when constructing a CI for μ , we were taking Z with μ .

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Let us check² that it satisfies the requirements:

- ▶ If we will have the Observation, we can calculate it, for sure;
- ▶ If the Null Hypo is True, then $\mu = \mu_0$. Recall what is the connection between our Statistics, Z and μ : the Distribution of X_k in Z is $\mathcal{N}(\mu, \sigma^2)$. So, under \mathcal{H}_0 , $X_k \sim \mathcal{N}(\mu_0, \sigma^2)$, so

$$\bar{X} \sim \mathcal{N}\left(\mu_0, \frac{\sigma^2}{n}\right)$$

²Note the difference from the Pivot Z : when constructing a CI for μ , we were taking Z with μ .

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Let us check² that it satisfies the requirements:

- ▶ If we will have the Observation, we can calculate it, for sure;
- ▶ If the Null Hypo is True, then $\mu = \mu_0$. Recall what is the connection between our Statistics, Z and μ : the Distribution of X_k in Z is $\mathcal{N}(\mu, \sigma^2)$. So, under \mathcal{H}_0 , $X_k \sim \mathcal{N}(\mu_0, \sigma^2)$, so

$$\bar{X} \sim \mathcal{N}\left(\mu_0, \frac{\sigma^2}{n}\right) \quad \text{and} \quad Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim$$

²Note the difference from the Pivot Z : when constructing a CI for μ , we were taking Z with μ .

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Let us check² that it satisfies the requirements:

- ▶ If we will have the Observation, we can calculate it, for sure;
- ▶ If the Null Hypo is True, then $\mu = \mu_0$. Recall what is the connection between our Statistics, Z and μ : the Distribution of X_k in Z is $\mathcal{N}(\mu, \sigma^2)$. So, under \mathcal{H}_0 , $X_k \sim \mathcal{N}(\mu_0, \sigma^2)$, so

$$\bar{X} \sim \mathcal{N}\left(\mu_0, \frac{\sigma^2}{n}\right) \quad \text{and} \quad Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1).$$

²Note the difference from the Pivot Z : when constructing a CI for μ , we were taking Z with μ .

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: So, we will base our Test-Statistics on \bar{X} : as the name is suggesting, we take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Let us check² that it satisfies the requirements:

- ▶ If we will have the Observation, we can calculate it, for sure;
- ▶ If the Null Hypo is True, then $\mu = \mu_0$. Recall what is the connection between our Statistics, Z and μ : the Distribution of X_k in Z is $\mathcal{N}(\mu, \sigma^2)$. So, under \mathcal{H}_0 , $X_k \sim \mathcal{N}(\mu_0, \sigma^2)$, so

$$\bar{X} \sim \mathcal{N}\left(\mu_0, \frac{\sigma^2}{n}\right) \quad \text{and} \quad Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1).$$

Great!

²Note the difference from the Pivot Z : when constructing a CI for μ , we were taking Z with μ .

Test for the Mean of the Normal, σ is known: Z -Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

In this case we will Reject \mathcal{H}_0 , if

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

In this case we will Reject \mathcal{H}_0 , if Z will be far from 0, i.e., we choose $RR = \{|Z| > c\}$.

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

In this case we will Reject \mathcal{H}_0 , if Z will be far from 0, i.e., we choose $RR = \{|Z| > c\}$. The Critical Value c is yet to be determined.

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

In this case we will Reject \mathcal{H}_0 , if Z will be far from 0, i.e., we choose $RR = \{|Z| > c\}$. The Critical Value c is yet to be determined.

Case 2: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

In this case we will not believe in \mathcal{H}_0 , if

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

In this case we will Reject \mathcal{H}_0 , if Z will be far from 0, i.e., we choose $RR = \{|Z| > c\}$. The Critical Value c is yet to be determined.

Case 2: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

In this case we will not believe in \mathcal{H}_0 , if Z will be far to the **Right** to 0, i.e., we choose $RR = \{Z > c\}$.

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

In this case we will Reject \mathcal{H}_0 , if Z will be far from 0, i.e., we choose $RR = \{|Z| > c\}$. The Critical Value c is yet to be determined.

Case 2: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

In this case we will not believe in \mathcal{H}_0 , if Z will be far to the **Right** to 0, i.e., we choose $RR = \{Z > c\}$. Again, the Critical Value c is yet to be determined.

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

In this case we will Reject \mathcal{H}_0 , if Z will be far from 0, i.e., we choose $RR = \{|Z| > c\}$. The Critical Value c is yet to be determined.

Case 2: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

In this case we will not believe in \mathcal{H}_0 , if Z will be far to the **Right** to 0, i.e., we choose $RR = \{Z > c\}$. Again, the Critical Value c is yet to be determined.

Case 3: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

In this case we will not believe in \mathcal{H}_0 , if

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

In this case we will Reject \mathcal{H}_0 , if Z will be far from 0, i.e., we choose $RR = \{|Z| > c\}$. The Critical Value c is yet to be determined.

Case 2: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

In this case we will not believe in \mathcal{H}_0 , if Z will be far to the **Right** to 0, i.e., we choose $RR = \{Z > c\}$. Again, the Critical Value c is yet to be determined.

Case 3: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

In this case we will not believe in \mathcal{H}_0 , if Z will be far to the **Left** to 0, i.e., we choose $RR = \{Z < -c\}$.

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

In this case we will Reject \mathcal{H}_0 , if Z will be far from 0, i.e., we choose $RR = \{|Z| > c\}$. The Critical Value c is yet to be determined.

Case 2: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

In this case we will not believe in \mathcal{H}_0 , if Z will be far to the **Right** to 0, i.e., we choose $RR = \{Z > c\}$. Again, the Critical Value c is yet to be determined.

Case 3: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

In this case we will not believe in \mathcal{H}_0 , if Z will be far to the **Left** to 0, i.e., we choose $RR = \{Z < -c\}$. Again, the Critical Value c is yet to be determined.

Choosing the Critical Value

Now, let us choose c .

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha =$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha = \mathbb{P}(\text{Type I Error}) =$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) =$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\begin{aligned}\alpha &= \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) = \\ &= \mathbb{P}(|Z| > c \mid \mu = \mu_0) =\end{aligned}$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\begin{aligned}\alpha &= \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) = \\ &= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).\end{aligned}$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) =$$

$$= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).$$

This means, geometrically, that

$$c =$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) =$$

$$= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).$$

This means, geometrically, that

$$c = z_{1-\alpha/2},$$

the $1 - \frac{\alpha}{2}$ -level quantile of the Standard Normal Distribution.

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) =$$

$$= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).$$

This means, geometrically, that

$$c = z_{1-\alpha/2},$$

the $1 - \frac{\alpha}{2}$ -level quantile of the Standard Normal Distribution.

So, finally, we have the Test for the Case 1: given μ_0 , σ , Observations and Significance Level α , calculate $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$.

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) =$$

$$= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).$$

This means, geometrically, that

$$c = z_{1-\alpha/2},$$

the $1 - \frac{\alpha}{2}$ -level quantile of the Standard Normal Distribution.

So, finally, we have the Test for the Case 1: given μ_0 , σ , Observations and Significance Level α , calculate $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$.

► If $|Z| > z_{1-\alpha/2}$,

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\begin{aligned}\alpha &= \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) = \\ &= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).\end{aligned}$$

This means, geometrically, that

$$c = z_{1-\alpha/2},$$

the $1 - \frac{\alpha}{2}$ -level quantile of the Standard Normal Distribution.

So, finally, we have the Test for the Case 1: given μ_0 , σ , Observations and Significance Level α , calculate $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$.

- ▶ If $|Z| > z_{1-\alpha/2}$, **Reject** \mathcal{H}_0 ;
- ▶ If $|Z| \leq z_{1-\alpha/2}$,

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\begin{aligned}\alpha &= \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) = \\ &= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).\end{aligned}$$

This means, geometrically, that

$$c = z_{1-\alpha/2},$$

the $1 - \frac{\alpha}{2}$ -level quantile of the Standard Normal Distribution.

So, finally, we have the Test for the Case 1: given μ_0 , σ , Observations and Significance Level α , calculate $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$.

- ▶ If $|Z| > z_{1-\alpha/2}$, **Reject** \mathcal{H}_0 ;
- ▶ If $|Z| \leq z_{1-\alpha/2}$, **Do Not Reject** \mathcal{H}_0 .

Example

Example: I have generated in **R** a Sample of Size 50 from $\mathcal{N}(3, 2^2)$ and made some rounding:

```
set.seed(20112019)
s.size <- 50; sigma <- 2
obs <- rnorm(s.size, mean = 3, sd = sigma)
obs <- round(obs, digits = 2); obs
```

```
## [1] 1.68 5.48 0.98 3.08 4.79 5.03 1.64 2.35 0.
## [13] 3.86 4.67 1.86 4.38 3.40 4.01 -0.20 3.75 4.
## [25] -0.25 4.82 -1.12 0.44 -1.28 7.98 3.11 1.87 4.
## [37] 0.99 4.25 7.10 7.35 2.64 4.78 3.55 4.55 5.
## [49] 0.78 0.15
```

Example

Example: I have generated in **R** a Sample of Size 50 from $\mathcal{N}(3, 2^2)$ and made some rounding:

```
set.seed(20112019)
s.size <- 50; sigma <- 2
obs <- rnorm(s.size, mean = 3, sd = sigma)
obs <- round(obs, digits = 2); obs
```

```
## [1] 1.68 5.48 0.98 3.08 4.79 5.03 1.64 2.35 0.
## [13] 3.86 4.67 1.86 4.38 3.40 4.01 -0.20 3.75 4.
## [25] -0.25 4.82 -1.12 0.44 -1.28 7.98 3.11 1.87 4.
## [37] 0.99 4.25 7.10 7.35 2.64 4.78 3.55 4.55 5.
## [49] 0.78 0.15
```

I will assume I do not know μ (which is 3, of course), and will just assume my Observation is coming from $\mathcal{N}(\mu, 2^2)$, with some μ .

Example

Example: I have generated in **R** a Sample of Size 50 from $\mathcal{N}(3, 2^2)$ and made some rounding:

```
set.seed(20112019)
s.size <- 50; sigma <- 2
obs <- rnorm(s.size, mean = 3, sd = sigma)
obs <- round(obs, digits = 2); obs
```

```
## [1] 1.68 5.48 0.98 3.08 4.79 5.03 1.64 2.35 0.
## [13] 3.86 4.67 1.86 4.38 3.40 4.01 -0.20 3.75 4.
## [25] -0.25 4.82 -1.12 0.44 -1.28 7.98 3.11 1.87 4.
## [37] 0.99 4.25 7.10 7.35 2.64 4.78 3.55 4.55 5.
## [49] 0.78 0.15
```

I will assume I do not know μ (which is 3, of course), and will just assume my Observation is coming from $\mathcal{N}(\mu, 2^2)$, with some μ . And I will test, at the 5% Significance Level, the following Hypothesis:

$$\mathcal{H}_0 : \mu = 4 \quad \text{vs} \quad \mathcal{H}_1 : \mu \neq 4.$$

Example, Cont'd

First, I calculate Z-statistic:

```
mu0 <- 4  
Z <- (mean(obs) - mu0)/(sigma/sqrt(s.size)); Z  
  
## [1] -3.63665
```

Example, Cont'd

First, I calculate Z-statistic:

```
mu0 <- 4  
Z <- (mean(obs) - mu0)/(sigma/sqrt(s.size)); Z
```

```
## [1] -3.63665
```

Now, I am calculating the quantile $z_{1-\alpha/2}$:

```
a <- 0.05  
z <- qnorm(1-a/2); z
```

```
## [1] 1.959964
```

Example, Cont'd

First, I calculate Z-statistic:

```
mu0 <- 4  
Z <- (mean(obs) - mu0)/(sigma/sqrt(s.size)); Z
```

```
## [1] -3.63665
```

Now, I am calculating the quantile $z_{1-\alpha/2}$:

```
a <- 0.05  
z <- qnorm(1-a/2); z
```

```
## [1] 1.959964
```

Finally, I am checking if $|Z| > z_{1-\alpha/2}$:

```
abs(Z) > z
```

```
## [1] TRUE
```

Example, Cont'd

First, I calculate Z-statistic:

```
mu0 <- 4  
Z <- (mean(obs) - mu0)/(sigma/sqrt(s.size)); Z
```

```
## [1] -3.63665
```

Now, I am calculating the quantile $z_{1-\alpha/2}$:

```
a <- 0.05  
z <- qnorm(1-a/2); z
```

```
## [1] 1.959964
```

Finally, I am checking if $|Z| > z_{1-\alpha/2}$:

```
abs(Z) > z
```

```
## [1] TRUE
```

So the decision is:

Example, Cont'd

First, I calculate Z-statistic:

```
mu0 <- 4  
Z <- (mean(obs) - mu0)/(sigma/sqrt(s.size)); Z  
  
## [1] -3.63665
```

Now, I am calculating the quantile $z_{1-\alpha/2}$:

```
a <- 0.05  
z <- qnorm(1-a/2); z  
  
## [1] 1.959964
```

Finally, I am checking if $|Z| > z_{1-\alpha/2}$:

```
abs(Z) > z
```

```
## [1] TRUE
```

So the decision is: **Reject** \mathcal{H}_0 .

Example, Cont'd

First, I calculate Z-statistic:

```
mu0 <- 4  
Z <- (mean(obs) - mu0)/(sigma/sqrt(s.size)); Z  
  
## [1] -3.63665
```

Now, I am calculating the quantile $z_{1-\alpha/2}$:

```
a <- 0.05  
z <- qnorm(1-a/2); z  
  
## [1] 1.959964
```

Finally, I am checking if $|Z| > z_{1-\alpha/2}$:

```
abs(Z) > z  
  
## [1] TRUE
```

So the decision is: **Reject** \mathcal{H}_0 . In this case we say that the result was **Statistically Significant**.

Example

Example: Now, with the same Observations from the last example, let us test, at the 5% level, the following Hypothesis:

$$\mathcal{H}_0 : \mu = 3.3 \quad \text{vs} \quad \mathcal{H}_0 : \mu \neq 3.3.$$

Example

Example: Now, with the same Observations from the last example, let us test, at the 5% level, the following Hypothesis:

$$\mathcal{H}_0 : \mu = 3.3 \quad \text{vs} \quad \mathcal{H}_0 : \mu \neq 3.3.$$

```
mu0 <- 3.3; a <- 0.05
Z <- (mean(obs) - mu0)/(sigma/sqrt(s.size))
cat("Z-statistics = ", Z)

## Z-statistics = -1.161776

z <- qnorm(1-a/2)
cat("critical value = ", z)

## critical value = 1.959964

if (abs(Z) > z) cat("Reject") else cat("Do Not Reject")

## Do Not Reject
```