

AUA CS 108, Statistics, Fall 2019

Lecture 21

Michael Poghosyan

YSU, AUA

michael@ysu.am, mpoghosyan@aua.am

11 Oct 2019

Contents

- ▶ Bias and Biasedness/Unbiasedness

On Quiz Mistakes

- ▶ if x is the height in Meters and x' is the height in Centimeters, then

$$x' = 100 \cdot x$$

On Quiz Mistakes

- ▶ if x is the height in Meters and x' is the height in Centimeters, then

$$x' = 100 \cdot x$$

- ▶ in our course, we have x , Dataset, set of numbers, and X , a r.v. .

On Quiz Mistakes

- ▶ if x is the height in Meters and x' is the height in Centimeters, then

$$x' = 100 \cdot x$$

- ▶ in our course, we have x , Dataset, set of numbers, and X , a r.v. .The Mean of x is $\bar{x} = \frac{x_1+x_2+\dots+x_n}{n}$, and the Mean of X is $\mathbb{E}(X)$.

On Quiz Mistakes

- ▶ if x is the height in Meters and x' is the height in Centimeters, then

$$x' = 100 \cdot x$$

- ▶ in our course, we have x , Dataset, set of numbers, and X , a r.v. .The Mean of x is $\bar{x} = \frac{x_1+x_2+\dots+x_n}{n}$, and the Mean of X is $\mathbb{E}(X)$.
- ▶ For the CDF of X_n , do not write $F(X_n)$, but write $F_{X_n}(x)$.

On Quiz Mistakes

- ▶ if x is the height in Meters and x' is the height in Centimeters, then

$$x' = 100 \cdot x$$

- ▶ in our course, we have x , Dataset, set of numbers, and X , a r.v. .The Mean of x is $\bar{x} = \frac{x_1+x_2+\dots+x_n}{n}$, and the Mean of X is $\mathbb{E}(X)$.
- ▶ For the CDF of X_n , do not write $F(X_n)$, but write $F_{X_n}(x)$. Similarly, do not write

$$\mathbb{E}(X_n) = \int x_n f(x) dx$$

but, instead,

On Quiz Mistakes

- ▶ if x is the height in Meters and x' is the height in Centimeters, then

$$x' = 100 \cdot x$$

- ▶ in our course, we have x , Dataset, set of numbers, and X , a r.v. .The Mean of x is $\bar{x} = \frac{x_1+x_2+\dots+x_n}{n}$, and the Mean of X is $\mathbb{E}(X)$.
- ▶ For the CDF of X_n , do not write $F(X_n)$, but write $F_{X_n}(x)$. Similarly, do not write

$$\mathbb{E}(X_n) = \int x_n f(x) dx$$

but, instead,

$$\mathbb{E}(X_n) = \int x f_{X_n}(x) dx$$

On Quiz Mistakes

- ▶ if x is the height in Meters and x' is the height in Centimeters, then

$$x' = 100 \cdot x$$

- ▶ in our course, we have x , Dataset, set of numbers, and X , a r.v. .The Mean of x is $\bar{x} = \frac{x_1+x_2+\dots+x_n}{n}$, and the Mean of X is $\mathbb{E}(X)$.
- ▶ For the CDF of X_n , do not write $F(X_n)$, but write $F_{X_n}(x)$. Similarly, do not write

$$\mathbb{E}(X_n) = \int x_n f(x) dx$$

but, instead,

$$\mathbb{E}(X_n) = \int x f_{X_n}(x) dx$$

- ▶ the CDF of $X + \frac{1}{n^2}$ is not $F_X(x) + \frac{1}{n^2}$.

Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.

Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.
- ▶ Define the MSE.

Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.
- ▶ Define the MSE.
- ▶ How we can define that $\hat{\theta}$ is close to θ ?

Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.
- ▶ Define the MSE.
- ▶ How we can define that $\hat{\theta}$ is close to θ ?
- ▶ Define the Bias of an Estimator.

Last Lecture ReCap

- ▶ State the Problem of the Point Estimation.
- ▶ Define the MSE.
- ▶ How we can define that $\hat{\theta}$ is close to θ ?
- ▶ Define the Bias of an Estimator.
- ▶ What is the definition of the Unbiased/Biased Estimator?

Example

Example: Now, let's not forget that we can have **R** in our Midterm 2 or Final 😊. So let's do an experiment with Biased and Unbiased Estimators.

Example

Example: Now, let's not forget that we can have **R** in our Midterm 2 or Final 😊. So let's do an experiment with Biased and Unbiased Estimators.

UnBiased Estimator Case

We consider the Poisson Model:

$$X_1, X_2, \dots, X_{10} \sim \text{Pois}(\lambda)$$

and we want to estimate λ .

Example

Example: Now, let's not forget that we can have **R** in our Midterm 2 or Final 😊. So let's do an experiment with Biased and Unbiased Estimators.

UnBiased Estimator Case

We consider the Poisson Model:

$$X_1, X_2, \dots, X_{10} \sim \text{Pois}(\lambda)$$

and we want to estimate λ . We consider the following Estimator:

$$\hat{\lambda} = \frac{X_1 + X_2 + \dots + X_{10}}{10}.$$

Example

Example: Now, let's not forget that we can have **R** in our Midterm 2 or Final 😊. So let's do an experiment with Biased and Unbiased Estimators.

UnBiased Estimator Case

We consider the Poisson Model:

$$X_1, X_2, \dots, X_{10} \sim \text{Pois}(\lambda)$$

and we want to estimate λ . We consider the following Estimator:

$$\hat{\lambda} = \frac{X_1 + X_2 + \dots + X_{10}}{10}.$$

Easy to see that $\hat{\lambda}$ is an Unbiased Estimator for λ (OTB!).

Example, cont'd

Now, the code


- ▶ observing once: generating a Sample just once and calculating one Estimate:

```
lambda <- 3.21  
x <- rpois(10, lambda = lambda)  
lambda.hat <- mean(x)  
lambda.hat
```

```
## [1] 3.4
```

- ▶ observing many times: generating Samples many times, calculating Estimates, and then averaging:

```
lambda <- 3.21; n <- 10; m <- 200  
x <- rpois(n*m, lambda = lambda)  
x <- as.data.frame(matrix(x, ncol = m))  
lambda.hats <- sapply(x, mean)  
boxplot(lambda.hats, horizontal = T);  
abline(v = lambda, col="red", lwd = 2, lty = 2)
```




```
mean(lambda.hats)
```

```
## [1] 3.172
```

With a Histogram:

```
hist(lambda.hats)
rug(lambda.hats)
abline(v = lambda, col="red", lwd = 2, lty = 2)
```



Example

Biased Estimator Case

Say, let us consider the Exponential Model:

$$X_1, X_2, \dots, X_{10} \sim \text{Exp}(\lambda)$$

and we want to estimate λ .

Example

Biased Estimator Case

Say, let us consider the Exponential Model:

$$X_1, X_2, \dots, X_{10} \sim \text{Exp}(\lambda)$$

and we want to estimate λ . We consider the following Estimator:

$$\hat{\lambda} = \frac{X_1 + X_2 + \dots + X_{10}}{10}.$$

Example

Biased Estimator Case

Say, let us consider the Exponential Model:

$$X_1, X_2, \dots, X_{10} \sim \text{Exp}(\lambda)$$

and we want to estimate λ . We consider the following Estimator:

$$\hat{\lambda} = \frac{X_1 + X_2 + \dots + X_{10}}{10}.$$

Easy to see that $\hat{\lambda}$ is an Biased Estimator for λ (OTB!).

Example, cont'd

Now, the code:


- ▶ observing once: generating a Sample just once and calculating one Estimate:

```
lambda <- 0.3  
x <- rexp(10, rate = lambda)  
lambda.hat <- mean(x)  
lambda.hat
```

```
## [1] 2.3389
```

- ▶ observing many times: generating Samples many times, calculating Estimates, and then averaging:

```
lambda <- 0.3; n <- 10; m <- 2000  
x <- rexp(n*m, rate = lambda)  
x <- as.data.frame(matrix(x, ncol = m))  
lambda.hats <- sapply(x, mean)  
boxplot(lambda.hats, horizontal = T);  
abline(v = lambda, col="red", lwd = 2, lty = 2)
```




```
mean(lambda.hats)
```

```
## [1] 3.380309
```

With a Histogram:

```
hist(lambda.hats)
rug(lambda.hats)
abline(v = lambda, col="red", lwd = 2, lty = 2)
```



Asymptotic Unbiasedness

Some years ago Unbiasedness was a very important, desirable property from any good Estimator.

Asymptotic Unbiasedness

Some years ago Unbiasedness was a very important, desirable property from any good Estimator. Nowadays, it is enough to have *Asymptotic Unbiasedness*:

Asymptotic Unbiasedness

Some years ago Unbiasedness was a very important, desirable property from any good Estimator. Nowadays, it is enough to have *Asymptotic Unbiasedness*:

Definition: Estimator $\hat{\theta}_n$ is called **Asymptotically Unbiased** for θ , if

$$\text{Bias}(\hat{\theta}_n, \theta) \rightarrow 0, \quad \text{for any } \theta \in \Theta.$$

Asymptotic Unbiasedness

Some years ago Unbiasedness was a very important, desirable property from any good Estimator. Nowadays, it is enough to have *Asymptotic Unbiasedness*:

Definition: Estimator $\hat{\theta}_n$ is called **Asymptotically Unbiased** for θ , if

$$\text{Bias}(\hat{\theta}_n, \theta) \rightarrow 0, \quad \text{for any } \theta \in \Theta.$$

Idea: If the Sample size is very large, then the behaviour of our Asymptotic Unbiased Estimator is close to an Unbiased one, $\text{Bias}(\hat{\theta}_n, \theta) \approx 0$

Asymptotic Unbiasedness

Some years ago Unbiasedness was a very important, desirable property from any good Estimator. Nowadays, it is enough to have *Asymptotic Unbiasedness*:

Definition: Estimator $\hat{\theta}_n$ is called **Asymptotically Unbiased** for θ , if

$$\text{Bias}(\hat{\theta}_n, \theta) \rightarrow 0, \quad \text{for any } \theta \in \Theta.$$

Idea: If the Sample size is very large, then the behaviour of our Asymptotic Unbiased Estimator is close to an Unbiased one, $\text{Bias}(\hat{\theta}_n, \theta) \approx 0$

Example: Say, for the Mean μ of the Population,

$$\hat{\mu} = \frac{X_1 + X_2 + \dots + X_n}{n + 1}$$

is a Biased, but Asymptotically Unbiased Estimator. OTB, please!

Bias-Variance Decomposition

This is an important result:

Theorem(Bias-Variance Decomposition of the MSE): If $\hat{\theta}$ is an Estimator for θ , then

$$MSE(\hat{\theta}, \theta) = \left(Bias(\hat{\theta}, \theta) \right)^2 + Var_{\theta}(\hat{\theta}).$$

Bias-Variance Decomposition

This is an important result:

Theorem(Bias-Variance Decomposition of the MSE): If $\hat{\theta}$ is an Estimator for θ , then

$$MSE(\hat{\theta}, \theta) = \left(Bias(\hat{\theta}, \theta) \right)^2 + Var_{\theta}(\hat{\theta}).$$

Proof: OTB