

CS 107, Probability, Spring 2019

Lecture 19

Michael Poghosyan

AUA

01 March 2019

Content

- CDF and its Properties
- Discrete r.v.

Laplace's Law of Succession

We know that the Sun has risen each day for the past 5000 years.

Laplace's Law of Succession

We know that the Sun has risen each day for the past 5000 years.

Question: What is the probability that it will rise tomorrow?

Laplace's Law of Succession

We know that the Sun has risen each day for the past 5000 years.

Question: What is the probability that it will rise tomorrow?

Laplace's answer was:

Laplace's Law of Succession

We know that the Sun has risen each day for the past 5000 years.

Question: What is the probability that it will rise tomorrow?

Laplace's answer was: $\frac{N+1}{N+2}$, where $N = 1,826,213$.

Random Variables, reminder

Assume we have an Experiment with Sample Space Ω .

Random Variables, reminder

Assume we have an Experiment with Sample Space Ω .

- Any function $X : \Omega \rightarrow \mathbb{R}$ is called a Random Variable (r.v.) (on the Experiment);

Random Variables, reminder

Assume we have an Experiment with Sample Space Ω .

- Any function $X : \Omega \rightarrow \mathbb{R}$ is called a Random Variable (r.v.) (on the Experiment);
- Can you give some examples of r.v.?

Random Variables, reminder

Assume we have an Experiment with Sample Space Ω .

- Any function $X: \Omega \rightarrow \mathbb{R}$ is called a Random Variable (r.v.) (on the Experiment);
- Can you give some examples of r.v.?
- If X is a r.v. $A \subset \mathbb{R}$, then $X \in A$ is an Event;

Random Variables, reminder

Assume we have an Experiment with Sample Space Ω .

- Any function $X: \Omega \rightarrow \mathbb{R}$ is called a Random Variable (r.v.) (on the Experiment);
- Can you give some examples of r.v.?
- If X is a r.v. $A \subset \mathbb{R}$, then $X \in A$ is an Event; In particular, for any $a < b$, the followings are Events:
 $a < X < b$, $a \leq X \leq b$, $a < X \leq b$, $a \leq X < b$

Random Variables, reminder

Assume we have an Experiment with Sample Space Ω .

- Any function $X: \Omega \rightarrow \mathbb{R}$ is called a Random Variable (r.v.) (on the Experiment);
- Can you give some examples of r.v.?
- If X is a r.v. $A \subset \mathbb{R}$, then $X \in A$ is an Event; In particular, for any $a < b$, the followings are Events:
 $a < X < b$, $a \leq X \leq b$, $a < X \leq b$, $a \leq X < b$
- Having a Complete Information about a r.v. = being able to calculate $\mathbb{P}(X \in A)$ for any $A \subset \mathbb{R}$;

Random Variables, reminder

Assume we have an Experiment with Sample Space Ω .

- Any function $X: \Omega \rightarrow \mathbb{R}$ is called a Random Variable (r.v.) (on the Experiment);
- Can you give some examples of r.v.?
- If X is a r.v. $A \subset \mathbb{R}$, then $X \in A$ is an Event; In particular, for any $a < b$, the followings are Events:
 $a < X < b$, $a \leq X \leq b$, $a < X \leq b$, $a \leq X < b$
- Having a Complete Information about a r.v. = being able to calculate $\mathbb{P}(X \in A)$ for any $A \subset \mathbb{R}$;
- To be able to calculate $\mathbb{P}(X \in A)$ for any $A \subset \mathbb{R}$, it is enough to know the values of the CDF at any point:

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R}.$$

Properties of the CDF

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Properties of the CDF

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

Properties of the CDF

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;

Properties of the CDF

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;
- $F(-\infty) = 0$ and $F(+\infty) = 1$;

Properties of the CDF

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;
- $F(-\infty) = 0$ and $F(+\infty) = 1$;
- F is an increasing function, i.e., if $x_1 \leq x_2$, then $F(x_1) \leq F(x_2)$;

Properties of the CDF

Assume $F(x)$ is the CDF of the r.v. X :

$$F(x) = F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R},$$

Then:

Properties of CDF

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;
- $F(-\infty) = 0$ and $F(+\infty) = 1$;
- F is an increasing function, i.e., if $x_1 \leq x_2$, then $F(x_1) \leq F(x_2)$;
- F is right-continuous at every point, i.e. $F(x_0+) = F(x_0)$ at any $x_0 \in \mathbb{R}$

Characterization of the CDF

Now we want to answer the inverse question: which functions F can serve as CDFs for some r.v.?

Characterization of the CDF

Now we want to answer the inverse question: which functions F can serve as CDFs for some r.v.? It turns out that the above properties **completely characterize CDFs**:

Characterization of the CDF

Now we want to answer the inverse question: which functions F can serve as CDFs for some r.v.? It turns out that the above properties **completely characterize CDFs**:

Characterization of CDFs

Assume $F: \mathbb{R} \rightarrow \mathbb{R}$ is a function satisfying

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;
- $F(-\infty) = 0$ and $F(+\infty) = 1$;
- F is an increasing function, i.e., if $x_1 \leq x_2$, then $F(x_1) \leq F(x_2)$;
- F is right-continuous at every point, i.e. $F(x_0+) = F(x_0)$ at any $x_0 \in \mathbb{R}$

Characterization of the CDF

Now we want to answer the inverse question: which functions F can serve as CDFs for some r.v.? It turns out that the above properties **completely characterize CDFs**:

Characterization of CDFs

Assume $F: \mathbb{R} \rightarrow \mathbb{R}$ is a function satisfying

- $0 \leq F(x) \leq 1$, for any $x \in \mathbb{R}$;
- $F(-\infty) = 0$ and $F(+\infty) = 1$;
- F is an increasing function, i.e., if $x_1 \leq x_2$, then $F(x_1) \leq F(x_2)$;
- F is right-continuous at every point, i.e. $F(x_0+) = F(x_0)$ at any $x_0 \in \mathbb{R}$

Then there exists an Experiment with a Sample Space Ω , Probability Measure \mathbb{P} and a r.v. $X: \Omega \rightarrow \mathbb{R}$ such that $F(x)$ is the CDF of X : $F(x) = F_X(x)$, $x \in \mathbb{R}$.

Graphical Example of a CDF




Figure: CDF of some r.v. X

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF.

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF. Assume X is a r.v., and $F(x)$ is its CDF. Then:

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF. Assume X is a r.v., and $F(x)$ is its CDF. Then:

Probabilities through CDF

- $\mathbb{P}(X = a) = F(a) - F(a-);$

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF. Assume X is a r.v., and $F(x)$ is its CDF. Then:

Probabilities through CDF

- $\mathbb{P}(X = a) = F(a) - F(a-);$
- $\mathbb{P}(a < X \leq b) = F(b) - F(a);$

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF. Assume X is a r.v., and $F(x)$ is its CDF. Then:

Probabilities through CDF

- $\mathbb{P}(X = a) = F(a) - F(a-);$
- $\mathbb{P}(a < X \leq b) = F(b) - F(a);$
- $\mathbb{P}(a \leq X \leq b) = F(b) - F(a-);$

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF. Assume X is a r.v., and $F(x)$ is its CDF. Then:

Probabilities through CDF

- $\mathbb{P}(X = a) = F(a) - F(a-);$
- $\mathbb{P}(a < X \leq b) = F(b) - F(a);$
- $\mathbb{P}(a \leq X \leq b) = F(b) - F(a-);$
- $\mathbb{P}(a \leq X < b) = F(b-) - F(a-);$

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF. Assume X is a r.v., and $F(x)$ is its CDF. Then:

Probabilities through CDF

- $\mathbb{P}(X = a) = F(a) - F(a-);$
- $\mathbb{P}(a < X \leq b) = F(b) - F(a);$
- $\mathbb{P}(a \leq X \leq b) = F(b) - F(a-);$
- $\mathbb{P}(a \leq X < b) = F(b-) - F(a-);$
- $\mathbb{P}(a < X < b) = F(b-) - F(a);$

Using CDFs to calculate Probabilities

Now we want to calculate Probabilities using the CDF. Assume X is a r.v., and $F(x)$ is its CDF. Then:

Probabilities through CDF

- $\mathbb{P}(X = a) = F(a) - F(a-);$
- $\mathbb{P}(a < X \leq b) = F(b) - F(a);$
- $\mathbb{P}(a \leq X \leq b) = F(b) - F(a-);$
- $\mathbb{P}(a \leq X < b) = F(b-) - F(a-);$
- $\mathbb{P}(a < X < b) = F(b-) - F(a);$

Here it is possible that $a = -\infty$ or/and $b = +\infty$

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 2)$, $\mathbb{P}(2 < X \leq 3)$, $\mathbb{P}(X \leq 5)$, $\mathbb{P}(X > 2)$:

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 2)$, $\mathbb{P}(2 < X \leq 3)$, $\mathbb{P}(X \leq 5)$, $\mathbb{P}(X > 2)$:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$:

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$:

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Problem: For the r.v. X given through its CDF below, which values are more probable:

Reading Probabilities from the graph of CDF

Problem: For the r.v. X given through its CDF below, which values are more probable:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$:

Reading Probabilities from the graph of CDF

Problem: Is the following a CDF of some r.v. X ? If yes, calculate $\mathbb{P}(X = 1)$, $\mathbb{P}(0 < X \leq 3)$, $\mathbb{P}(X \leq 1)$, $\mathbb{P}(X > 0)$:




Figure: CDF of some r.v. X

Reading Probabilities from the graph of CDF

Problem: Below you can find graphs of 2 CDFs: Red is for r.v. X , and Black is for Y . Which one is larger: $\mathbb{P}(2 < X < 4)$ or $\mathbb{P}(2 < Y < 4)$?

Reading Probabilities from the graph of CDF

Problem: Below you can find graphs of 2 CDFs: Red is for r.v. X , and Black is for Y . Which one is larger: $\mathbb{P}(2 < X < 4)$ or $\mathbb{P}(2 < Y < 4)$?




Figure: CDFs of r.v. X and Y

Discrete R.V.s

Discrete Random Variables

In our course, we will consider 2 types of r.v.: **Discrete** and **Continuous**.

¹Right Hand Side

Discrete Random Variables

In our course, we will consider 2 types of r.v.: **Discrete** and **Continuous**. Assume X is a r.v. defined on Ω , i.e., $X: \Omega \rightarrow \mathbb{R}$.

¹Right Hand Side

Discrete Random Variables

In our course, we will consider 2 types of r.v.: **Discrete** and **Continuous**. Assume X is a r.v. defined on Ω , i.e., $X: \Omega \rightarrow \mathbb{R}$.

Discrete Random Variable

We say that the r.v. X is **Discrete**, if the Range of X ,

$Range(X) = \{X(\omega) : \omega \in \Omega\}$ = The set of all possible values of X is finite or countably infinite.

¹Right Hand Side

Discrete Random Variables

In our course, we will consider 2 types of r.v.: **Discrete** and **Continuous**. Assume X is a r.v. defined on Ω , i.e., $X: \Omega \rightarrow \mathbb{R}$.

Discrete Random Variable

We say that the r.v. X is **Discrete**, if the Range of X ,

$Range(X) = \{X(\omega) : \omega \in \Omega\}$ = The set of all possible values of X is finite or countably infinite.

So if X is Discrete, then the Range of X can be written as

$$Range(X) = \{x_1, x_2, x_3, \dots\},$$

where the set on the RHS¹ can be also finite.

¹Right Hand Side

Discrete Random Variables

Example:

- Let X be the number of Heads when tossing 5 coins.
What is the range of X ? Is X discrete?

Discrete Random Variables

Example:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Can you give some more?

Discrete Random Variables

Example:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Can you give some more?
- If Ω is Discrete, then X will be Discrete;

Discrete Random Variables

Example:

- Let X be the number of Heads when tossing 5 coins. What is the range of X ? Is X discrete?
- Can you give some more?
- If Ω is Discrete, then X will be Discrete;
- If Ω is not Discrete, then X CAN BE Discrete.