

# CS 107, Probability, Spring 2019

## Lecture 14

Michael Poghosyan

AUA

18 February 2019

- Repeated, Independent Trials: Binomial Distribution, Cont'd
- Repeated, Independent Trials: Multinomial Distribution

Quiz:

## Quiz:

- a. We are rolling 2 fair dice. What is the probability that the difference between the shown numbers will be 2, if we know that one of the shown numbers was less than 3?

## Quiz:

- a. We are rolling 2 fair dice. What is the probability that the difference between the shown numbers will be 2, if we know that one of the shown numbers was less than 3?
- b. I am choosing 2 persons from our group of 36 participants. We know that 10 of them are female. What is the probability that one of the chosen persons will be a female and the other one - a male?

# Binomial Probabilities formula, again:

## Binomial Probabilities

For any  $k = 0, 1, \dots, n$ ,

$$\mathbb{P}(\text{Exactly } k \text{ successes in } n \text{ trials}) = \binom{n}{k} \cdot p^k \cdot (1 - p)^{n-k}.$$

or, if we will denote the number of Successes in that  $n$  Trials by  $X$ ,

$$\mathbb{P}(X = k) = \binom{n}{k} \cdot p^k \cdot (1 - p)^{n-k}.$$

# Binomial Probabilities formula, again:

## Binomial Probabilities

For any  $k = 0, 1, \dots, n$ ,

$$\mathbb{P}(\text{Exactly } k \text{ successes in } n \text{ trials}) = \binom{n}{k} \cdot p^k \cdot (1 - p)^{n-k}.$$

or, if we will denote the number of Successes in that  $n$  Trials by  $X$ ,

$$\mathbb{P}(X = k) = \binom{n}{k} \cdot p^k \cdot (1 - p)^{n-k}.$$

**Proof:** ONB

# Example:

**Problem:** Some entrance exam test consists of 30 multiple choice problems.



# Example:

**Problem:** Some entrance exam test consists of 30 multiple choice problems. Every correct choice gives 1 points.

# Example:

**Problem:** Some entrance exam test consists of 30 multiple choice problems. Every correct choice gives 1 points. Each problem has suggested 5 choices for the answer, from which only one is correct.

# Example:

**Problem:** Some entrance exam test consists of 30 multiple choice problems. Every correct choice gives 1 points. Each problem has suggested 5 choices for the answer, from which only one is correct. If the person will choose randomly the answers, what is the probability that

# Example:

**Problem:** Some entrance exam test consists of 30 multiple choice problems. Every correct choice gives 1 points. Each problem has suggested 5 choices for the answer, from which only one is correct. If the person will choose randomly the answers, what is the probability that

- He/she will get 25 points?

# Example:

**Problem:** Some entrance exam test consists of 30 multiple choice problems. Every correct choice gives 1 points. Each problem has suggested 5 choices for the answer, from which only one is correct. If the person will choose randomly the answers, what is the probability that

- He/she will get 25 points?
- He/she will get more than 25 points?

# Example:

**Problem:** Some entrance exam test consists of 30 multiple choice problems. Every correct choice gives 1 points. Each problem has suggested 5 choices for the answer, from which only one is correct. If the person will choose randomly the answers, what is the probability that

- He/she will get 25 points?
- He/she will get more than 25 points?
- He/she will get less than 29 points?

# Repeated, Independent Trials: Multinomial Case

Now we consider an Experiment consisting of repetition of a Simple Experiment, which can result in more than 2 events.

# Repeated, Independent Trials: Multinomial Case

Now we consider an Experiment consisting of repetition of a Simple Experiment, which can result in more than 2 events. More precisely,



# Repeated, Independent Trials: Multinomial Case

Now we consider an Experiment consisting of repetition of a Simple Experiment, which can result in more than 2 events. More precisely,

- We have a Simple Experiment (Trial);

# Repeated, Independent Trials: Multinomial Case

Now we consider an Experiment consisting of repetition of a Simple Experiment, which can result in more than 2 events. More precisely,

- We have a Simple Experiment (Trial);
- The Simple Experiment can result in one of the mutually exclusive Events  $A_1, A_2, \dots, A_m$ ;

# Repeated, Independent Trials: Multinomial Case

Now we consider an Experiment consisting of repetition of a Simple Experiment, which can result in more than 2 events. More precisely,

- We have a Simple Experiment (Trial);
- The Simple Experiment can result in one of the mutually exclusive Events  $A_1, A_2, \dots, A_m$ ;
- The probability of having  $A_k$  in one Trial is  $p_k$ , i.e.,  $\mathbb{P}(A_k) = p_k$ , and  $p_1 + p_2 + \dots + p_m = 1$ ;

# Repeated, Independent Trials: Multinomial Case

Now we consider an Experiment consisting of repetition of a Simple Experiment, which can result in more than 2 events. More precisely,

- We have a Simple Experiment (Trial);
- The Simple Experiment can result in one of the mutually exclusive Events  $A_1, A_2, \dots, A_m$ ;
- The probability of having  $A_k$  in one Trial is  $p_k$ , i.e.,  $\mathbb{P}(A_k) = p_k$ , and  $p_1 + p_2 + \dots + p_m = 1$ ;
- We repeat our Trial  $n$  times, independently;

# Repeated, Independent Trials: Multinomial Case

Now we consider an Experiment consisting of repetition of a Simple Experiment, which can result in more than 2 events. More precisely,

- We have a Simple Experiment (Trial);
- The Simple Experiment can result in one of the mutually exclusive Events  $A_1, A_2, \dots, A_m$ ;
- The probability of having  $A_k$  in one Trial is  $p_k$ , i.e.,  $\mathbb{P}(A_k) = p_k$ , and  $p_1 + p_2 + \dots + p_m = 1$ ;
- We repeat our Trial  $n$  times, independently;
- We are interested in the probability that we will have exactly  $k_1$  outcomes from  $A_1$ , exactly  $k_2$  outcomes from  $A_2, \dots$ , and exactly  $k_m$  outcomes from  $A_m$ .

# Repeated, Independent Trials: Multinomial Case

Now we consider an Experiment consisting of repetition of a Simple Experiment, which can result in more than 2 events. More precisely,

- We have a Simple Experiment (Trial);
- The Simple Experiment can result in one of the mutually exclusive Events  $A_1, A_2, \dots, A_m$ ;
- The probability of having  $A_k$  in one Trial is  $p_k$ , i.e.,  $\mathbb{P}(A_k) = p_k$ , and  $p_1 + p_2 + \dots + p_m = 1$ ;
- We repeat our Trial  $n$  times, independently;
- We are interested in the probability that we will have exactly  $k_1$  outcomes from  $A_1$ , exactly  $k_2$  outcomes from  $A_2, \dots$ , and exactly  $k_m$  outcomes from  $A_m$ .
- Clearly, we need to have (we are doing  $n$  trials)

$$k_1 + k_2 + \dots + k_m = n.$$

# Multinomial Case, Remark

Note that the Binomial Case is the particular case of the Multinomial: in that case

# Multinomial Case, Remark

Note that the Binomial Case is the particular case of the Multinomial: in that case

- Each trial can result either in  $A$  (Success) or in  $\bar{A}$  (Failure);



# Multinomial Case, Remark

Note that the Binomial Case is the particular case of the Multinomial: in that case

- Each trial can result either in  $A$  (Success) or in  $\bar{A}$  (Failure);
- So we have  $m = 2$ ,  $A_1 = A$ ,  $A_2 = \bar{A}$ ;

# Multinomial Case, Remark

Note that the Binomial Case is the particular case of the Multinomial: in that case

- Each trial can result either in  $A$  (Success) or in  $\bar{A}$  (Failure);
- So we have  $m = 2$ ,  $A_1 = A$ ,  $A_2 = \bar{A}$ ;
- We have  $p_1 = \mathbb{P}(A_1) = \mathbb{P}(A) = p$ , so  $p_2 = \mathbb{P}(A_2) = \mathbb{P}(\bar{A}) = 1 - p$

# Multinomial Case, Remark

Note that the Binomial Case is the particular case of the Multinomial: in that case

- Each trial can result either in  $A$  (Success) or in  $\bar{A}$  (Failure);
- So we have  $m = 2$ ,  $A_1 = A$ ,  $A_2 = \bar{A}$ ;
- We have  $p_1 = \mathbb{P}(A_1) = \mathbb{P}(A) = p$ , so  $p_2 = \mathbb{P}(A_2) = \mathbb{P}(\bar{A}) = 1 - p$
- We are interested in having exactly  $k$  Successes in  $n$  Trials, that is, we are interested in having exactly  $k_1 = k$  times  $A_1$  (Success) and exactly  $k_2 = n - k_1$  times  $A_2$  (Failure)

# Repeated Indep Trials: Multinomial Probabilities

## Multinomial Probabilities

The probability that exactly  $k_1$  times we will have the event  $A_1$ , exactly  $k_2$  times we will have the event  $A_2$ , ... , exactly  $k_m$  times we will have the event  $A_m$  in the above described  $n$  trials, is

$$P_n(k_1, k_2, \dots, k_m) = \binom{n}{k_1, k_2, \dots, k_m} \cdot p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_m^{k_m},$$

where  $p_k = \mathbb{P}(A_k)$ ,  $k = 1, \dots, m$ .

# Repeated Indep Trials: Multinomial Probabilities

## Multinomial Probabilities

The probability that exactly  $k_1$  times we will have the event  $A_1$ , exactly  $k_2$  times we will have the event  $A_2$ , ... , exactly  $k_m$  times we will have the event  $A_m$  in the above described  $n$  trials, is

$$P_n(k_1, k_2, \dots, k_m) = \binom{n}{k_1, k_2, \dots, k_m} \cdot p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_m^{k_m},$$

where  $p_k = \mathbb{P}(A_k)$ ,  $k = 1, \dots, m$ .

Here

$$\binom{n}{k_1, k_2, \dots, k_m} = \frac{n!}{k_1! \cdot k_2! \cdot \dots \cdot k_m!}$$

# Repeated Indep Trials: Multinomial Probabilities

## Multinomial Probabilities

The probability that exactly  $k_1$  times we will have the event  $A_1$ , exactly  $k_2$  times we will have the event  $A_2$ , ... , exactly  $k_m$  times we will have the event  $A_m$  in the above described  $n$  trials, is

$$P_n(k_1, k_2, \dots, k_m) = \binom{n}{k_1, k_2, \dots, k_m} \cdot p_1^{k_1} \cdot p_2^{k_2} \cdot \dots \cdot p_m^{k_m},$$

where  $p_k = \mathbb{P}(A_k)$ ,  $k = 1, \dots, m$ .

Here

$$\binom{n}{k_1, k_2, \dots, k_m} = \frac{n!}{k_1! \cdot k_2! \cdot \dots \cdot k_m!}$$

Note that in the binomial case, i.e., when  $m = 2$ , we get the Binomial Probabilities formula.