

AUA CS 108, Statistics, Fall 2019

Lecture 29

Michael Poghosyan

YSU, AUA

michael@ysu.am, mpoghosyan@aua.am

30 Oct 2019

Contents

- ▶ The Maximum Likelihood Method (MLE)

Last Lecture ReCap

- ▶ I was asking: *Any Questions?* What was the answer?

The Maximum Likelihood Method

Examples, MLE

Example: Find the MLE Estimator for (μ, σ^2) in the $\mathcal{N}(\mu, \sigma^2)$ Model.

Solution: OTB

Examples, MLE

Example: Find the MLE Estimator for (μ, σ^2) in the $\mathcal{N}(\mu, \sigma^2)$ Model.

Solution: OTB

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

X	0	1	2
$\mathbb{P}(X = x)$	$\frac{\theta}{10}$	$\frac{\theta}{5}$	$1 - \frac{3\theta}{10}$

where $\theta \in [0, \frac{10}{3}]$.

Examples, MLE

Example: Find the MLE Estimator for (μ, σ^2) in the $\mathcal{N}(\mu, \sigma^2)$ Model.

Solution: OTB

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

X	0	1	2
$\mathbb{P}(X = x)$	$\frac{\theta}{10}$	$\frac{\theta}{5}$	$1 - \frac{3\theta}{10}$

where $\theta \in [0, \frac{10}{3}]$. Find the MLE Estimator and MLE Estimate for θ .

Solution: OTB

Examples, MLE


Example: Find the MLE Estimator for θ in the one-Parametric Cauchy Distribution $Cauchy(\theta)$ Model. Here, the PDF of $X \sim Cauchy(\theta)$ is given by

$$f(x|\theta) = \frac{1}{\pi(1 + (x - \theta)^2)}, \quad x \in \mathbb{R},$$

and $\theta \in \mathbb{R}$ is called the *location parameter*.

PDF of Cauchy(0) and Cauchy(2)

```
x <- seq(from = -10, to = 10, by = 0.01)
y1 <- dcauchy(x); y2 <- dcauchy(x, location = 2);
plot(x, y1, type = "l", lwd = 2, xlim = c(-10,10), col = "blue")
par(new = T)
plot(x, y2, type = "l", lwd = 2, xlim = c(-10,10), col = "red")
```



Examples, MLE

Example: Find the MLE Estimator for θ in the one-Parametric Cauchy Distribution $Cauchy(\theta)$ Model. Here, the PDF of $X \sim Cauchy(\theta)$ is given by

$$f(x|\theta) = \frac{1}{\pi(1 + (x - \theta)^2)}, \quad x \in \mathbb{R},$$

and $\theta \in \mathbb{R}$ is called the *location parameter*.

Solution: OTB

Examples, MLE

Example: Find the MLE Estimator for θ in the one-Parametric Cauchy Distribution $Cauchy(\theta)$ Model. Here, the PDF of $X \sim Cauchy(\theta)$ is given by

$$f(x|\theta) = \frac{1}{\pi(1 + (x - \theta)^2)}, \quad x \in \mathbb{R},$$

and $\theta \in \mathbb{R}$ is called the *location parameter*.

Solution: OTB

Note: The MoM will not work in this case, since Moments of this Distribution do not exist.

Examples, MLE

Example: Find the MLE Estimator for θ in the one-Parametric Cauchy Distribution $Cauchy(\theta)$ Model. Here, the PDF of $X \sim Cauchy(\theta)$ is given by

$$f(x|\theta) = \frac{1}{\pi(1 + (x - \theta)^2)}, \quad x \in \mathbb{R},$$

and $\theta \in \mathbb{R}$ is called the *location parameter*.

Solution: OTB

Note: The MoM will not work in this case, since Moments of this Distribution do not exist. Btw, the Median of this Distribution is θ .

Examples, MLE

Example: Find the MLE Estimator for θ in the one-Parametric Cauchy Distribution $Cauchy(\theta)$ Model. Here, the PDF of $X \sim Cauchy(\theta)$ is given by

$$f(x|\theta) = \frac{1}{\pi(1 + (x - \theta)^2)}, \quad x \in \mathbb{R},$$

and $\theta \in \mathbb{R}$ is called the *location parameter*.

Solution: OTB

Note: The MoM will not work in this case, since Moments of this Distribution do not exist. Btw, the Median of this Distribution is θ .

Note: In **R**, the name of Cauchy Distribution is *cauchy*, so you can use, say, `rcauchy` or `dcauchy` .

Examples, MLE

Example: Find the MLE Estimator for θ in the one-Parametric Cauchy Distribution $Cauchy(\theta)$ Model. Here, the PDF of $X \sim Cauchy(\theta)$ is given by

$$f(x|\theta) = \frac{1}{\pi(1 + (x - \theta)^2)}, \quad x \in \mathbb{R},$$

and $\theta \in \mathbb{R}$ is called the *location parameter*.

Solution: OTB

Note: The MoM will not work in this case, since Moments of this Distribution do not exist. Btw, the Median of this Distribution is θ .

Note: In **R**, the name of Cauchy Distribution is *cauchy*, so you can use, say, `rcauchy` or `dcauchy` .

Note: Do you know what is the Distribution of the ratio $\frac{X}{Y}$, when X and Y are Independent Standard Normal Rvs?

Examples, MLE

Example: Find the MLE Estimator for θ in the one-Parametric Cauchy Distribution $Cauchy(\theta)$ Model. Here, the PDF of $X \sim Cauchy(\theta)$ is given by

$$f(x|\theta) = \frac{1}{\pi(1 + (x - \theta)^2)}, \quad x \in \mathbb{R},$$

and $\theta \in \mathbb{R}$ is called the *location parameter*.

Solution: OTB

Note: The MoM will not work in this case, since Moments of this Distribution do not exist. Btw, the Median of this Distribution is θ .

Note: In **R**, the name of Cauchy Distribution is *cauchy*, so you can use, say, `rcauchy` or `dcauchy` .

Note: Do you know what is the Distribution of the ratio $\frac{X}{Y}$, when X and Y are Independent Standard Normal Rvs? $Cauchy(0)$! 😊