

AUA CS108, Statistics, Fall 2020

Lecture 23

Michael Poghosyan

19 Oct 2020

Contents

- ▶ Convergence Types of R.V. Sequences, Some Theorems
- ▶ Limit Theorems

Some Properties

Theorem: Assume $X_n \xrightarrow{a.s.(P, qm, D)} X$ and $Y_n \xrightarrow{a.s.(P, qm, D)} Y$.

Some Properties

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

► $X_n + Y_n \xrightarrow{a.s.(P,qm)} X + Y;$

Some Properties

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

► $X_n + Y_n \xrightarrow{a.s.(P,qm)} X + Y;$

► $X_n \cdot Y_n \xrightarrow{a.s.(P)} X \cdot Y;$

Some Properties

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- ▶ $X_n + Y_n \xrightarrow{a.s.(P,qm)} X + Y$;
- ▶ $X_n \cdot Y_n \xrightarrow{a.s.(P)} X \cdot Y$; and $\alpha \cdot X_n \xrightarrow{qm} \alpha \cdot X$ for any constant α ;

Some Properties

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- ▶ $X_n + Y_n \xrightarrow{a.s.(P,qm)} X + Y$;
- ▶ $X_n \cdot Y_n \xrightarrow{a.s.(P)} X \cdot Y$; and $\alpha \cdot X_n \xrightarrow{qm} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \xrightarrow{a.s.(P,D)} g(X)$

Some Properties

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- ▶ $X_n + Y_n \xrightarrow{a.s.(P,qm)} X + Y$;
- ▶ $X_n \cdot Y_n \xrightarrow{a.s.(P)} X \cdot Y$; and $\alpha \cdot X_n \xrightarrow{qm} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \xrightarrow{a.s.(P,D)} g(X)$

Note: In the general case, if $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{D} Y$, then not necessarily $X_n + Y_n \xrightarrow{D} X + Y$.

Some Properties

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- ▶ $X_n + Y_n \xrightarrow{a.s.(P,qm)} X + Y$;
- ▶ $X_n \cdot Y_n \xrightarrow{a.s.(P)} X \cdot Y$; and $\alpha \cdot X_n \xrightarrow{qm} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \xrightarrow{a.s.(P,D)} g(X)$

Note: In the general case, if $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{D} Y$, then not necessarily $X_n + Y_n \xrightarrow{D} X + Y$.

But we have:

Theorem (Slutsky) Assume $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{P} c$, where $c \in \mathbb{R}$ is a constant. Then

Some Properties

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- ▶ $X_n + Y_n \xrightarrow{a.s.(P,qm)} X + Y$;
- ▶ $X_n \cdot Y_n \xrightarrow{a.s.(P)} X \cdot Y$; and $\alpha \cdot X_n \xrightarrow{qm} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \xrightarrow{a.s.(P,D)} g(X)$

Note: In the general case, if $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{D} Y$, then not necessarily $X_n + Y_n \xrightarrow{D} X + Y$.

But we have:

Theorem (Slutsky) Assume $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{P} c$, where $c \in \mathbb{R}$ is a constant. Then

- ▶ $X_n + Y_n \xrightarrow{D} X + c$;

Some Properties

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- ▶ $X_n + Y_n \xrightarrow{a.s.(P,qm)} X + Y$;
- ▶ $X_n \cdot Y_n \xrightarrow{a.s.(P)} X \cdot Y$; and $\alpha \cdot X_n \xrightarrow{qm} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \xrightarrow{a.s.(P,D)} g(X)$

Note: In the general case, if $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{D} Y$, then not necessarily $X_n + Y_n \xrightarrow{D} X + Y$.


But we have:

Theorem (Slutsky) Assume $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{P} c$, where $c \in \mathbb{R}$ is a constant. Then

- ▶ $X_n + Y_n \xrightarrow{D} X + c$;
- ▶ $X_n \cdot Y_n \xrightarrow{D} c \cdot X$.


Relationship between Convergence Types

Theorem: (Convergence Relationship Diagram)



Relationship between Convergence Types


Theorem: (Convergence Relationship Diagram)



Note: Inverse implications are not always correct.

Relationship between Convergence Types

Theorem: (Convergence Relationship Diagram)



Note: Inverse implications are not always correct. But, say, the following holds: If $X_n \xrightarrow{D} X$ and $X \equiv \text{constant}$, then $X_n \xrightarrow{\mathbb{P}} X$ (X_n and X are defined on the same Probability space).

Note

Note: Mostly, in our course, we will deal with the following type of sequences of r.v.s:

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n},$$

Note

Note: Mostly, in our course, we will deal with the following type of sequences of r.v.s:

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n},$$

and to calculate the limit of this sequence $\bar{X}_1, \bar{X}_2, \dots, \bar{X}_n, \dots$, we will use our famous Limit Theorems: LLN and CLT.

Limit Theorems

Sequence of IID r.v.

Assume X_n is a sequence of **Independent, Identically Distributed (IID)** r.v.s.

Sequence of IID r.v.

Assume X_n is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All X_n -s have the same Distribution.

Sequence of IID r.v.

Assume X_n is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All X_n -s have the same Distribution. In particular, all numerical partial characteristics of X_n coincide.

Sequence of IID r.v.

Assume X_n is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All X_n -s have the same Distribution. In particular, all numerical partial characteristics of X_n coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

Sequence of IID r.v.

Assume X_n is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All X_n -s have the same Distribution. In particular, all numerical partial characteristics of X_n coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

We will use this many-many

Sequence of IID r.v.

Assume X_n is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All X_n -s have the same Distribution. In particular, all numerical partial characteristics of X_n coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

We will use this many-many-many-many-... times.

Sequence of IID r.v.

Assume X_n is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All X_n -s have the same Distribution. In particular, all numerical partial characteristics of X_n coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

We will use this many-many-many-many-... times.

- ▶ X_n -s are independent.

Sequence of IID r.v.

Assume X_n is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All X_n -s have the same Distribution. In particular, all numerical partial characteristics of X_n coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

We will use this many-many-many-many-... times.

- ▶ X_n -s are independent. Say, in particular,

$$\text{Var}(X_1 + X_2 + \dots + X_n) = \text{Var}(X_1) + \text{Var}(X_2) + \dots + \text{Var}(X_n) = n \cdot \text{Var}(X_1).$$

Idea of the LLN and CLT

Assume we have a sequence X_n of IID rvs.

Idea of the LLN and CLT

Assume we have a sequence X_n of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

Idea of the LLN and CLT

Assume we have a sequence X_n of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

or the average

$$\overline{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

Idea of the LLN and CLT

Assume we have a sequence X_n of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

or the average

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

Note: Not an easy task to find the Distribution of S_n or \bar{X}_n .

Idea of the LLN and CLT

Assume we have a sequence X_n of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

or the average

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

Note: Not an easy task to find the Distribution of S_n or \bar{X}_n . Even for $n = 2$.

Idea of the LLN and CLT

Assume we have a sequence X_n of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

or the average

$$\overline{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

Note: Not an easy task to find the Distribution of S_n or \overline{X}_n . Even for $n = 2$. We need Convolutions!

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) =$$

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) =$$

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) =$$

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) =$$

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}.$$

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}.$$

The last property is the mathematical proof of the effectiveness of “7 angam chapir, mek angam ktrir” ☺

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}.$$

The last property is the mathematical proof of the effectiveness of “7 angam chapir, mek angam ktrir” ☺

The interpretation of $\mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1)$ and $\text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}$:

What we know about S_n and \bar{X}_n

Some important known facts about S_n and \bar{X}_n in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}.$$

The last property is the mathematical proof of the effectiveness of “7 angam chapir, mek angam ktrir” ☺

The interpretation of $\mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1)$ and $\text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}$: the values of \bar{X}_n are centered at $\mathbb{E}(X_1)$ and are becoming more and more concentrated around that number as n increases.

Some Particular results about S_n and \overline{X}_n

The following are important cases when we can find exactly the Distribution of S_n and/or \overline{X}_n :

Some Particular results about S_n and \bar{X}_n

The following are important cases when we can find exactly the Distribution of S_n and/or \bar{X}_n :

- ▶ If $X_k \sim \mathcal{N}(\mu, \sigma^2)$, $k = 1, \dots, n$, are Independent, then

$$S_n = X_1 + \dots + X_n \sim$$

Some Particular results about S_n and \bar{X}_n

The following are important cases when we can find exactly the Distribution of S_n and/or \bar{X}_n :

- If $X_k \sim \mathcal{N}(\mu, \sigma^2)$, $k = 1, \dots, n$, are Independent, then

$$S_n = X_1 + \dots + X_n \sim \mathcal{N}(n \cdot \mu, n \cdot \sigma^2)$$

and

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \sim$$

Some Particular results about S_n and \bar{X}_n

The following are important cases when we can find exactly the Distribution of S_n and/or \bar{X}_n :

- If $X_k \sim \mathcal{N}(\mu, \sigma^2)$, $k = 1, \dots, n$, are Independent, then

$$S_n = X_1 + \dots + X_n \sim \mathcal{N}(n \cdot \mu, n \cdot \sigma^2)$$

and

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right).$$

Some Particular results about S_n and \bar{X}_n

The following are important cases when we can find exactly the Distribution of S_n and/or \bar{X}_n :

- ▶ If $X_k \sim \mathcal{N}(\mu, \sigma^2)$, $k = 1, \dots, n$, are Independent, then

$$S_n = X_1 + \dots + X_n \sim \mathcal{N}(n \cdot \mu, n \cdot \sigma^2)$$

and

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right).$$

- ▶ If $X_k \sim \text{Pois}(\lambda)$, $k = 1, \dots, n$, are Independent, then

$$S_n = X_1 + \dots + X_n \sim$$

Some Particular results about S_n and \overline{X}_n

The following are important cases when we can find exactly the Distribution of S_n and/or \overline{X}_n :

- ▶ If $X_k \sim \mathcal{N}(\mu, \sigma^2)$, $k = 1, \dots, n$, are Independent, then

$$S_n = X_1 + \dots + X_n \sim \mathcal{N}(n \cdot \mu, n \cdot \sigma^2)$$

and

$$\overline{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right).$$

- ▶ If $X_k \sim \text{Pois}(\lambda)$, $k = 1, \dots, n$, are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Pois}(n \cdot \lambda).$$

Some Particular results about S_n and \bar{X}_n

- ▶ If $X_k \sim \text{Bernoulli}(p)$, $k = 1, \dots, n$ are Independent, then

$$S_n = X_1 + \dots + X_n \sim$$

Some Particular results about S_n and \bar{X}_n

- ▶ If $X_k \sim \text{Bernoulli}(p)$, $k = 1, \dots, n$ are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Binom}(n, p);$$

Some Particular results about S_n and \overline{X}_n

- ▶ If $X_k \sim \text{Bernoulli}(p)$, $k = 1, \dots, n$ are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Binom}(n, p);$$

- ▶ If $X_k \sim \text{Binom}(m, p)$, $k = 1, \dots, n$, are Independent, then

$$S_n = X_1 + \dots + X_n \sim$$

Some Particular results about S_n and \overline{X}_n

- ▶ If $X_k \sim \text{Bernoulli}(p)$, $k = 1, \dots, n$ are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Binom}(n, p);$$

- ▶ If $X_k \sim \text{Binom}(m, p)$, $k = 1, \dots, n$, are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Binom}(n \cdot m, p).$$

The Weak LLN

Now, what can be said about S_n and \overline{X}_n in the general case?

The Weak LLN

Now, what can be said about S_n and \overline{X}_n in the general case? LLN and CLT help us in this matter, they describe the *asymptotic* properties of these guys:

The Weak LLN

Now, what can be said about S_n and \overline{X}_n in the general case? LLN and CLT help us in this matter, they describe the *asymptotic* properties of these guys:

The Weak Law of Large Numbers, WLLN:

If X_1, X_2, \dots, X_n are IID, with finite $\mathbb{E}(X_1)$ and Variance $\text{Var}(X_1)$, then

$$\overline{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{\mathbb{P}} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

The Weak LLN

Now, what can be said about S_n and \bar{X}_n in the general case? LLN and CLT help us in this matter, they describe the *asymptotic* properties of these guys:

The Weak Law of Large Numbers, WLLN:

If X_1, X_2, \dots, X_n are IID, with finite $\mathbb{E}(X_1)$ and Variance $\text{Var}(X_1)$, then

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{\mathbb{P}} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

i.e., for any $\varepsilon > 0$,

$$\mathbb{P} \left(\left| \frac{X_1 + X_2 + \dots + X_n}{n} - \mathbb{E}(X_1) \right| \geq \varepsilon \right) \rightarrow 0, \quad n \rightarrow +\infty.$$

The Weak LLN

Now, what can be said about S_n and \bar{X}_n in the general case? LLN and CLT help us in this matter, they describe the *asymptotic* properties of these guys:

The Weak Law of Large Numbers, WLLN:

If X_1, X_2, \dots, X_n are IID, with finite $\mathbb{E}(X_1)$ and Variance $\text{Var}(X_1)$, then

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{\mathbb{P}} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

i.e., for any $\varepsilon > 0$,

$$\mathbb{P} \left(\left| \frac{X_1 + X_2 + \dots + X_n}{n} - \mathbb{E}(X_1) \right| \geq \varepsilon \right) \rightarrow 0, \quad n \rightarrow +\infty.$$

Note: This means that for any $\varepsilon > 0$, the chances that \bar{X}_n is far from $\mathbb{E}(X_1)$ more than ε , is very small, if n is large.

The Strong LLN

The Strong Law of Large Numbers, SLLN, Kolmogorov

If X_1, X_2, \dots, X_n are IID, with finite $\mathbb{E}(|X_1|)$, then

$$\frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{a.s.} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

The Strong LLN

The Strong Law of Large Numbers, SLLN, Kolmogorov

If X_1, X_2, \dots, X_n are IID, with finite $\mathbb{E}(|X_1|)$, then

$$\frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{a.s.} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

that is,

$$\mathbb{P} \left(\lim_{n \rightarrow +\infty} \frac{X_1 + X_2 + \dots + X_n}{n} = \mathbb{E}(X_1) \right) = 1.$$

Visualization of the LLN

```
set.seed(111); n <- 1000; expect <- 0.6  
X <- rbinom(n, 1, expect)  
S <- cumsum(X); p <- S/(1:n)  
plot(p, type = "l")  
abline(expect, 0, col = "red", lwd = 2)
```

