

# CS 107, Probability, Spring 2019

## Lecture 21

Michael Poghosyan

AUA

13 March 2019

# Content

- Discrete r.v., PMF
- Continuous r.v., PDF

## Zeno's Paradoxes

- To reach the wall in front of me, I need to pass the half of the way first.

## Zeno's Paradoxes

- To reach the wall in front of me, I need to pass the half of the way first. Then I need to pass the half of the remaining half.

## Zeno's Paradoxes

- To reach the wall in front of me, I need to pass the half of the way first. Then I need to pass the half of the remaining half. Then I need to pass the half of the remaining quarter.

## Zeno's Paradoxes

- To reach the wall in front of me, I need to pass the half of the way first. Then I need to pass the half of the remaining half. Then I need to pass the half of the remaining quarter. Then the half of the remaining eight's part and so on.

## Zeno's Paradoxes

- To reach the wall in front of me, I need to pass the half of the way first. Then I need to pass the half of the remaining half. Then I need to pass the half of the remaining quarter. Then the half of the remaining eight's part and so on. Do you think I will reach the wall 😊 ?

## Zeno's Paradoxes

- To reach the wall in front of me, I need to pass the half of the way first. Then I need to pass the half of the remaining half. Then I need to pass the half of the remaining quarter. Then the half of the remaining eight's part and so on. Do you think I will reach the wall 😊 ?
- Achilles and the tortoise paradox, read the Wiki Page at [https://en.wikipedia.org/wiki/Zeno's\\_paradoxes](https://en.wikipedia.org/wiki/Zeno's_paradoxes)



## Example:

Let  $X$  is the number of heads shown when tossing a fair coin 4 times. Then  $X$  is a r.v. with the PMF:

Values of $X$					
$\mathbb{P}(X = x)$					

## Example:

Let  $X$  is the number of heads shown when tossing a fair coin 4 times. Then  $X$  is a r.v. with the PMF:

Values of $X$					
$\mathbb{P}(X = x)$					

Now,

- Calculate  $\mathbb{P}(X = 2)$ ;

## Example:

Let  $X$  is the number of heads shown when tossing a fair coin 4 times. Then  $X$  is a r.v. with the PMF:

Values of $X$					
$\mathbb{P}(X = x)$					

Now,

- Calculate  $\mathbb{P}(X = 2)$ ;
- Calculate  $\mathbb{P}(X \leq 2.5)$ ;

## Example:

Let  $X$  is the number of heads shown when tossing a fair coin 4 times. Then  $X$  is a r.v. with the PMF:

Values of $X$					
$\mathbb{P}(X = x)$					

Now,

- Calculate  $\mathbb{P}(X = 2)$ ;
- Calculate  $\mathbb{P}(X \leq 2.5)$ ;
- Graph the PMF of  $X$ ;

## Example:

Let  $X$  is the number of heads shown when tossing a fair coin 4 times. Then  $X$  is a r.v. with the PMF:

Values of $X$					
$\mathbb{P}(X = x)$					

Now,

- Calculate  $\mathbb{P}(X = 2)$ ;
- Calculate  $\mathbb{P}(X \leq 2.5)$ ;
- Graph the PMF of  $X$ ;
- Graph the CDF of  $X$ .

# The Graph of CDF of a Discrete r.v

In general, if the r.v.  $X$  is Discrete, given through its PMF:

Values of $X$	$x_1$	$x_2$	$x_3$	...
$\mathbb{P}(X = x)$	$p_1$	$p_2$	$p_3$	...

then the graph of the CDF  $F(x)$  of  $X$  will be

# The Graph of CDF of a Discrete r.v

In general, if the r.v.  $X$  is Discrete, given through its PMF:

Values of $X$	$x_1$	$x_2$	$x_3$	...
$\mathbb{P}(X = x)$	$p_1$	$p_2$	$p_3$	...

then the graph of the CDF  $F(x)$  of  $X$  will be

- a piecewise constant function (step function);

# The Graph of CDF of a Discrete r.v

In general, if the r.v.  $X$  is Discrete, given through its PMF:

Values of $X$	$x_1$	$x_2$	$x_3$	...
$\mathbb{P}(X = x)$	$p_1$	$p_2$	$p_3$	...

then the graph of the CDF  $F(x)$  of  $X$  will be

- a piecewise constant function (step function);
- the jump points will be the points  $x_k$ ,  $k = 1, 2, \dots$ ;



# The Graph of CDF of a Discrete r.v

In general, if the r.v.  $X$  is Discrete, given through its PMF:

Values of $X$	$x_1$	$x_2$	$x_3$	...
$\mathbb{P}(X = x)$	$p_1$	$p_2$	$p_3$	...

then the graph of the CDF  $F(x)$  of  $X$  will be

- a piecewise constant function (step function);
- the jump points will be the points  $x_k$ ,  $k = 1, 2, \dots$ ;
- the jump size at the point  $x_k$  will be  $p_k$ ;

# The Graph of CDF of a Discrete r.v

In general, if the r.v.  $X$  is Discrete, given through its PMF:

Values of $X$	$x_1$	$x_2$	$x_3$	...
$\mathbb{P}(X = x)$	$p_1$	$p_2$	$p_3$	...

then the graph of the CDF  $F(x)$  of  $X$  will be

- a piecewise constant function (step function);
- the jump points will be the points  $x_k$ ,  $k = 1, 2, \dots$ ;
- the jump size at the point  $x_k$  will be  $p_k$ ;
- $F$  will be continuous from the right.

# The Graph of CDF of a Discrete r.v

In general, if the r.v.  $X$  is Discrete, given through its PMF:

Values of $X$	$x_1$	$x_2$	$x_3$	...
$\mathbb{P}(X = x)$	$p_1$	$p_2$	$p_3$	...

then the graph of the CDF  $F(x)$  of  $X$  will be

- a piecewise constant function (step function);
- the jump points will be the points  $x_k$ ,  $k = 1, 2, \dots$ ;
- the jump size at the point  $x_k$  will be  $p_k$ ;
- $F$  will be continuous from the right.

Graph on the board!

# PMF from the CDF

Conversely, if we have the CDF  $F(x)$  of a Discrete r.v.  $X$ , then:

# PMF from the CDF

Conversely, if we have the CDF  $F(x)$  of a Discrete r.v.  $X$ , then:

- We can calculate the PMF by:

$$\mathbb{P}(X = x) =$$

# PMF from the CDF

Conversely, if we have the CDF  $F(x)$  of a Discrete r.v.  $X$ , then:

- We can calculate the PMF by:

$$\mathbb{P}(X = x) = F(x) - F(x-) \quad \forall x \in \mathbb{R}$$

# PMF from the CDF

Conversely, if we have the CDF  $F(x)$  of a Discrete r.v.  $X$ , then:

- We can calculate the PMF by:

$$\mathbb{P}(X = x) = F(x) - F(x-) \quad \forall x \in \mathbb{R}$$

In fact, this will be non-zero only at  $x_k$ , the values from the  $\text{Range}(X)$ ;

# PMF from the CDF

Conversely, if we have the CDF  $F(x)$  of a Discrete r.v.  $X$ , then:

- We can calculate the PMF by:

$$\mathbb{P}(X = x) = F(x) - F(x-) \quad \forall x \in \mathbb{R}$$

In fact, this will be non-zero only at  $x_k$ , the values from the  $\text{Range}(X)$ ;

- Graphically, CDF will be a piecewise constant function



# PMF from the CDF

Conversely, if we have the CDF  $F(x)$  of a Discrete r.v.  $X$ , then:

- We can calculate the PMF by:

$$\mathbb{P}(X = x) = F(x) - F(x-) \quad \forall x \in \mathbb{R}$$

In fact, this will be non-zero only at  $x_k$ , the values from the  $\text{Range}(X)$ ;

- Graphically, CDF will be a piecewise constant function
- Then the possible values  $x_k$  of  $X$  will be the jump points of  $F$ ;

# PMF from the CDF

Conversely, if we have the CDF  $F(x)$  of a Discrete r.v.  $X$ , then:

- We can calculate the PMF by:

$$\mathbb{P}(X = x) = F(x) - F(x-) \quad \forall x \in \mathbb{R}$$

In fact, this will be non-zero only at  $x_k$ , the values from the  $\text{Range}(X)$ ;

- Graphically, CDF will be a piecewise constant function
- Then the possible values  $x_k$  of  $X$  will be the jump points of  $F$ ;
- The Probability  $\mathbb{P}(X = x_k)$  will be the jump size at  $x_k$

# PMF from the CDF

Conversely, if we have the CDF  $F(x)$  of a Discrete r.v.  $X$ , then:

- We can calculate the PMF by:

$$\mathbb{P}(X = x) = F(x) - F(x-) \quad \forall x \in \mathbb{R}$$

In fact, this will be non-zero only at  $x_k$ , the values from the  $\text{Range}(X)$ ;

- Graphically, CDF will be a piecewise constant function
- Then the possible values  $x_k$  of  $X$  will be the jump points of  $F$ ;
- The Probability  $\mathbb{P}(X = x_k)$  will be the jump size at  $x_k$

Give the graph!

# Continuous R.V.

# Intro: Continuous Random Variables

There are many r.v.s that are not Discrete. For example, the lifetime of a light bulb, the height of a (randomly chosen) person, the exact age (not rounded) of a person etc, the waiting time for the quiz grades

# Intro: Continuous Random Variables

There are many r.v.s that are not Discrete. For example, the lifetime of a light bulb, the height of a (randomly chosen) person, the exact age (not rounded) of a person etc, the waiting time for the quiz grades- I mean, metro train

# Intro: Continuous Random Variables

There are many r.v.s that are not Discrete. For example, the lifetime of a light bulb, the height of a (randomly chosen) person, the exact age (not rounded) of a person etc, the waiting time for the quiz grades- I mean, metro train at the bus station 😊 are all examples of non-Discrete r.v.s.

# Intro: Continuous Random Variables

There are many r.v.s that are not Discrete. For example, the lifetime of a light bulb, the height of a (randomly chosen) person, the exact age (not rounded) of a person etc, the waiting time for the quiz grades- I mean, metro train at the bus station 😊 are all examples of non-Discrete r.v.s.

In fact, there is a famous Lebesgue Decomposition Theorem:



# Intro: Continuous Random Variables

## Lebesgue Decomp Theorem, 18+

**Any** r.v.  $X$  can be represented in the form:

$$X = X_{discr} + X_{cont} + X_{sing},$$

where

# Intro: Continuous Random Variables

## Lebesgue Decomp Theorem, 18+

**Any** r.v.  $X$  can be represented in the form:

$$X = X_{discr} + X_{cont} + X_{sing},$$

where

- $X_{discr}$  is a Discrete r.v.;

# Intro: Continuous Random Variables

## Lebesgue Decomp Theorem, 18+

**Any** r.v.  $X$  can be represented in the form:

$$X = X_{discr} + X_{cont} + X_{sing},$$

where

- $X_{discr}$  is a Discrete r.v.;
- $X_{cont}$  is a Continuous r.v.;

# Intro: Continuous Random Variables

## Lebesgue Decomp Theorem, 18+

**Any** r.v.  $X$  can be represented in the form:

$$X = X_{discr} + X_{cont} + X_{sing},$$

where

- $X_{discr}$  is a Discrete r.v.;
- $X_{cont}$  is a Continuous r.v.;
- $X_{sing}$  is a Singular r.v.

# Intro: Continuous Random Variables

## Lebesgue Decomp Theorem, 18+

**Any** r.v.  $X$  can be represented in the form:

$$X = X_{discr} + X_{cont} + X_{sing},$$

where

- $X_{discr}$  is a Discrete r.v.;
- $X_{cont}$  is a Continuous r.v.;
- $X_{sing}$  is a Singular r.v.

We know already what Discrete r.v.s are, the next slides introduce Continuous r.v., and we will not talk about the Singular ones, because they are Singular 😊

# Prelude, Continuous Random Variables

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s:

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ .



# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because each point's weight will be 0.

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because each point's weight will be 0. We define the Weight of some part of our body by

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because each point's weight will be 0. We define the Weight of some part of our body by integrating the Density.

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because each point's weight will be 0. We define the Weight of some part of our body by integrating the Density. In the analogy, we will define the Probability  $\mathbb{P}(a \leq X \leq b)$  by integrating the Density Function over  $[a, b]$ .

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because each point's weight will be 0. We define the Weight of some part of our body by integrating the Density. In the analogy, we will define the Probability  $\mathbb{P}(a \leq X \leq b)$  by integrating the Density Function over  $[a, b]$ .

**LZ** - Zeno's paradox:

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because each point's weight will be 0. We define the Weight of some part of our body by integrating the Density. In the analogy, we will define the Probability  $\mathbb{P}(a \leq X \leq b)$  by integrating the Density Function over  $[a, b]$ .

**LZ** - Zeno's paradox: Consider an arrow in the flight.



# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because each point's weight will be 0. We define the Weight of some part of our body by integrating the Density. In the analogy, we will define the Probability  $\mathbb{P}(a \leq X \leq b)$  by integrating the Density Function over  $[a, b]$ .

**LZ** - Zeno's paradox: Consider an arrow in the flight. At any time instant, the arrow is not moving, is motionless.

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because each point's weight will be 0. We define the Weight of some part of our body by integrating the Density. In the analogy, we will define the Probability  $\mathbb{P}(a \leq X \leq b)$  by integrating the Density Function over  $[a, b]$ .

**LZ** - Zeno's paradox: Consider an arrow in the flight. At any time instant, the arrow is not moving, is motionless. Think as recording the flight and then considering just one frame at a time.

# Prelude, Continuous Random Variables

Now we introduce Continuous r.v.s: unfortunately, we cannot describe them using the probabilities  $\mathbb{P}(X = x)$ , since these probabilities are 0 for any continuous r.v.  $X$ , and any real number  $x$ . So we will not consider the Probability of  $X = x$ , rather Probabilities of the form  $\mathbb{P}(a \leq X \leq b)$ .

The idea is similar to defining the Weight of a physical body: we cannot define it through the weights of points, because each point's weight will be 0. We define the Weight of some part of our body by integrating the Density. In the analogy, we will define the Probability  $\mathbb{P}(a \leq X \leq b)$  by integrating the Density Function over  $[a, b]$ .

**LZ** - Zeno's paradox: Consider an arrow in the flight. At any time instant, the arrow is not moving, is motionless. Think as recording the flight and then considering just one frame at a time. Then how can it move at all?

# Continuous Random Variables

Assume  $X$  is a r.v. and  $F(x)$  is its CDF.

# Continuous Random Variables

Assume  $X$  is a r.v. and  $F(x)$  is its CDF.

## Continuous r.v.

We will say that the r.v.  $X$  is **absolutely continuous** or just **continuous**, if there exists an integrable non-negative function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that

$$F(x) = \int_{-\infty}^x f(t) dt, \quad \forall x \in \mathbb{R}.$$

# Continuous Random Variables

Assume  $X$  is a r.v. and  $F(x)$  is its CDF.

## Continuous r.v.

We will say that the r.v.  $X$  is **absolutely continuous** or just **continuous**, if there exists an integrable non-negative function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that

$$F(x) = \int_{-\infty}^x f(t) dt, \quad \forall x \in \mathbb{R}.$$

Here  $f$  is called the **Probability Density Function (PDF)** of  $X$ .

# Continuous Random Variables

Assume  $X$  is a r.v. and  $F(x)$  is its CDF.

## Continuous r.v.

We will say that the r.v.  $X$  is **absolutely continuous** or just **continuous**, if there exists an integrable non-negative function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that

$$F(x) = \int_{-\infty}^x f(t)dt, \quad \forall x \in \mathbb{R}.$$

Here  $f$  is called the **Probability Density Function (PDF)** of  $X$ . Sometimes we will write  $f_X(x)$  to stress that  $f$  is the PDF of  $X$ .

# Continuous Random Variables

Assume  $X$  is a r.v. and  $F(x)$  is its CDF.

## Continuous r.v.

We will say that the r.v.  $X$  is **absolutely continuous** or just **continuous**, if there exists an integrable non-negative function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that

$$F(x) = \int_{-\infty}^x f(t)dt, \quad \forall x \in \mathbb{R}.$$

Here  $f$  is called the **Probability Density Function (PDF)** of  $X$ . Sometimes we will write  $f_X(x)$  to stress that  $f$  is the PDF of  $X$ .

**Remark:** By a Theorem of *RA ∪ Calc2*, if  $X$  is a continuous r.v., then  $F(x)$  will be continuous at every  $x$ .



# Relation between PDF and CDF for Continuous r.v.

Assume  $F(x)$  is the CDF and  $f(x)$  is the PDF of a r.v.  $X$ .

# Relation between PDF and CDF for Continuous r.v.

Assume  $F(x)$  is the CDF and  $f(x)$  is the PDF of a r.v.  $X$ . Then, by definition,

$$F(x) =$$

# Relation between PDF and CDF for Continuous r.v.

Assume  $F(x)$  is the CDF and  $f(x)$  is the PDF of a r.v.  $X$ . Then, by definition,

$$F(x) = \int_{-\infty}^x f(t)dt, \quad \forall x \in \mathbb{R}.$$

# Relation between PDF and CDF for Continuous r.v.

Assume  $F(x)$  is the CDF and  $f(x)$  is the PDF of a r.v.  $X$ . Then, by definition,

$$F(x) = \int_{-\infty}^x f(t) dt, \quad \forall x \in \mathbb{R}.$$

Hence, using  $RA \cup Calc2$ , we get

$$f(x) =$$

# Relation between PDF and CDF for Continuous r.v.

Assume  $F(x)$  is the CDF and  $f(x)$  is the PDF of a r.v.  $X$ . Then, by definition,

$$F(x) = \int_{-\infty}^x f(t) dt, \quad \forall x \in \mathbb{R}.$$

Hence, using  $RA \cup Calc2$ , we get

$$f(x) = F'(x),$$

# Relation between PDF and CDF for Continuous r.v.

Assume  $F(x)$  is the CDF and  $f(x)$  is the PDF of a r.v.  $X$ . Then, by definition,

$$F(x) = \int_{-\infty}^x f(t)dt, \quad \forall x \in \mathbb{R}.$$

Hence, using  $RA \cup Calc2$ , we get

$$f(x) = F'(x), \quad \text{if } f \text{ is continuous at } x.$$

# Characterization of PDF

What are the characteristic properties of a PDF? The following Theorem give necessary and sufficient conditions in a function for being a PDF for some r.v.

# Characterization of PDF

What are the characteristic properties of a PDF? The following Theorem give necessary and sufficient conditions in a function for being a PDF for some r.v.

## Characterization of PDF

If  $X$  is a continuous r.v. with a PDF  $f(x)$ , then

a.  $f(x) \geq 0$  for all  $x \in \mathbb{R}$ ;

b.  $\int_{-\infty}^{+\infty} f(x) dx = 1.$



# Characterization of PDF

What are the characteristic properties of a PDF? The following Theorem give necessary and sufficient conditions in a function for being a PDF for some r.v.

## Characterization of PDF

If  $X$  is a continuous r.v. with a PDF  $f(x)$ , then

a.  $f(x) \geq 0$  for all  $x \in \mathbb{R}$ ;

b.  $\int_{-\infty}^{+\infty} f(x) dx = 1.$

Inversely, if  $f$  is an integrable function satisfying the above two conditions, then there is a Probability Space and a r.v. on that Space such that  $f$  is the PDF of that r.v.

# Geometric Interpretation of PDF

Recall that

$$\mathbb{P}(X \leq x) = F(x) = \int_{-\infty}^x f(t) dt, \quad \forall x \in \mathbb{R}.$$

# Geometric Interpretation of PDF

Recall that

$$\mathbb{P}(X \leq x) = F(x) = \int_{-\infty}^x f(t)dt, \quad \forall x \in \mathbb{R}.$$

And from  $RA \cup Calc2$  you know that the integral on the RHS is the *Area under the curve*  $y = f(x)$  over the interval  $(-\infty, x]$ .

# Geometric Interpretation of PDF

Recall that

$$\mathbb{P}(X \leq x) = F(x) = \int_{-\infty}^x f(t)dt, \quad \forall x \in \mathbb{R}.$$

And from *RA*  $\cup$  *Calc2* you know that the integral on the RHS is the *Area under the curve*  $y = f(x)$  over the interval  $(-\infty, x]$ . Hence,

$$\mathbb{P}(X \leq x) = F(x) = \text{Area}(\text{region under } y = f(x) \text{ on } (-\infty, x]).$$

# Geometric Interpretation of PDF





Figure: The Shaded Region area is  $\mathbb{P}(X \leq 1) = F(1)$

# Geometric Interpretation of PDF

**Fact:** The area under the PDF graph is 1.



# Using PDF to calculate Probabilities

Recall again that, if  $f$  and  $F$  are the PDF and CDF of r.v.  $X$ , then for any  $x \in \mathbb{R}$ ,

$$\mathbb{P}(X \leq x) = F(x) = \int_{-\infty}^x f(t)dt.$$

# Using PDF to calculate Probabilities

Recall again that, if  $f$  and  $F$  are the PDF and CDF of r.v.  $X$ , then for any  $x \in \mathbb{R}$ ,

$$\mathbb{P}(X \leq x) = F(x) = \int_{-\infty}^x f(t)dt.$$

And we know that

$$\mathbb{P}(a < X \leq b) =$$



# Using PDF to calculate Probabilities

Recall again that, if  $f$  and  $F$  are the PDF and CDF of r.v.  $X$ , then for any  $x \in \mathbb{R}$ ,

$$\mathbb{P}(X \leq x) = F(x) = \int_{-\infty}^x f(t)dt.$$

And we know that

$$\mathbb{P}(a < X \leq b) = F(b) - F(a) =$$

# Using PDF to calculate Probabilities

Recall again that, if  $f$  and  $F$  are the PDF and CDF of r.v.  $X$ , then for any  $x \in \mathbb{R}$ ,

$$\mathbb{P}(X \leq x) = F(x) = \int_{-\infty}^x f(t)dt.$$

And we know that

$$\begin{aligned}\mathbb{P}(a < X \leq b) &= F(b) - F(a) = \int_{-\infty}^b f(x)dx - \int_{-\infty}^a f(x)dx = \\ &= \end{aligned}$$

# Using PDF to calculate Probabilities

Recall again that, if  $f$  and  $F$  are the PDF and CDF of r.v.  $X$ , then for any  $x \in \mathbb{R}$ ,

$$\mathbb{P}(X \leq x) = F(x) = \int_{-\infty}^x f(t)dt.$$

And we know that

$$\begin{aligned}\mathbb{P}(a < X \leq b) &= F(b) - F(a) = \int_{-\infty}^b f(x)dx - \int_{-\infty}^a f(x)dx = \\ &= \int_a^b f(x)dx.\end{aligned}$$

So we have obtained the following relation:

$$\mathbb{P}(a < X \leq b) = \int_a^b f(x)dx.$$

# Using PDF to calculate Probabilities

And, in general, the following properties are true:

# Using PDF to calculate Probabilities

And, in general, the following properties are true:

## Calculation of Probabilities by PDF

If  $X$  is a r.v. with the PDF  $f(x)$  and CDF  $F(x)$ , then

- $\mathbb{P}(X = x) = 0$ ;

# Using PDF to calculate Probabilities

And, in general, the following properties are true:

## Calculation of Probabilities by PDF

If  $X$  is a r.v. with the PDF  $f(x)$  and CDF  $F(x)$ , then


- $\mathbb{P}(X = x) = 0$ ;
- $\mathbb{P}(a < X \leq b) = \mathbb{P}(a \leq X < b) = \mathbb{P}(a \leq X \leq b) =$

$$= \mathbb{P}(a < X < b) = \int_a^b f(x) dx$$

# Example:

**Example:** Below is the graph of the PDF  $f(x)$  for some r.v.  $X$ .

- Which one is larger:  $\mathbb{P}(X = 0)$  or  $\mathbb{P}(X = 2)$  ?



# Example:

- Show geometrically and calculate approximately  $\mathbb{P}(X = 0)$ ,  $\mathbb{P}(X < -1)$ ,  $\mathbb{P}(1 \leq X \leq 2)$

