

# CS 107, Probability, Spring 2019

## Lecture 45

Michael Poghosyan

AUA

the Last One

- Intro to Markov Chains

# Intro to Markov Chains

# Some Problems Concerning Markov Chains

Recall that last time we were considering one of the problems concerning MCs: the Probabilities of Paths

$$\mathbb{P}(X_0 = i_0, X_1 = i_1, \dots, X_n = i_n),$$

which can give us the idea, say, about the best possible scenario for our System for the next  $n$  time instants.

# Some Problems Concerning Markov Chains

The next important problem is to describe the time dynamics of the System, in particular, to find the  $n$ -Step Transition Probabilities

$$p_{ij}^{(n)} = \mathbb{P}(X_n = j | X_0 = i),$$

# Some Problems Concerning Markov Chains

The next important problem is to describe the time dynamics of the System, in particular, to find the  $n$ -Step Transition Probabilities

$$p_{ij}^{(n)} = \mathbb{P}(X_n = j | X_0 = i),$$

i.e., for a fixed  $n$ , to find the Probability that our System, starting at time 0 at the state  $i$ , will be at the State  $j$  after  $n$  steps.

# Some Problems Concerning Markov Chains

The next important problem is to describe the time dynamics of the System, in particular, to find the  $n$ -Step Transition Probabilities

$$p_{ij}^{(n)} = \mathbb{P}(X_n = j | X_0 = i),$$

i.e., for a fixed  $n$ , to find the Probability that our System, starting at time 0 at the state  $i$ , will be at the State  $j$  after  $n$  steps. Let us define the  $n$ -Step Transition Probability Matrix

$$P^{(n)} = \begin{bmatrix} p_{ij}^{(n)} \end{bmatrix}.$$

# Some Problems Concerning Markov Chains

The next important problem is to describe the time dynamics of the System, in particular, to find the  $n$ -Step Transition Probabilities

$$p_{ij}^{(n)} = \mathbb{P}(X_n = j | X_0 = i),$$

i.e., for a fixed  $n$ , to find the Probability that our System, starting at time 0 at the state  $i$ , will be at the State  $j$  after  $n$  steps. Let us define the  $n$ -Step Transition Probability Matrix

$$P^{(n)} = \left[ p_{ij}^{(n)} \right].$$

**Note:** Please note that because our MC is Time-Homogeneous, then, for any  $m \in \mathbb{N}$ ,

$$\mathbb{P}(X_{n+m} = j | X_m = i) = \mathbb{P}(X_n = j | X_0 = i) = p_{ij}^{(n)}.$$



# $n$ -Step Transition Probabilities

Using the Total Probability Formula, one can prove the following theorem:

## Chapman - Kolmogorov Equation

For any  $n \in \mathbb{N}$ ,

$$P^{(n)} = P^n, \quad \text{and, consequently,} \quad P^{(n+m)} = P^{(n)} \cdot P^{(m)}.$$

Here  $P^n$  is the  $n$ -th power of the Transition Probability Matrix  $P$ .

# $n$ -Step Transition Probabilities

Using the Total Probability Formula, one can prove the following theorem:

## Chapman - Kolmogorov Equation

For any  $n \in \mathbb{N}$ ,

$$P^{(n)} = P^n, \quad \text{and, consequently,} \quad P^{(n+m)} = P^{(n)} \cdot P^{(m)}.$$

Here  $P^n$  is the  $n$ -th power of the Transition Probability Matrix  $P$ .

In the terms of the matrix elements, the last equality can be written in the form

$$p_{ij}^{(n+m)} = \sum_{k=1}^N p_{ik}^{(n)} \cdot p_{kj}^{(m)}.$$

**Interpretation:** Give ☺

# Example

**Example: On/Off System** Again we consider our ON/OFF System. It can be in two states, over some discrete time interval: ON and OFF.

# Example

**Example: On/Off System** Again we consider our ON/OFF System. It can be in two states, over some discrete time interval: ON and OFF. If the System is in the State OFF at some time instant, then, at the next instant it will be at the State ON with probability  $p$ .

# Example

**Example: On/Off System** Again we consider our ON/OFF System. It can be in two states, over some discrete time interval: ON and OFF. If the System is in the State OFF at some time instant, then, at the next instant it will be at the State ON with probability  $p$ . Also, if the System is in the State ON at some time instant, then, at the next instant it will be at the State OFF with probability  $q$ .

# Example

**Example: On/Off System** Again we consider our ON/OFF System. It can be in two states, over some discrete time interval: ON and OFF. If the System is in the State OFF at some time instant, then, at the next instant it will be at the State ON with probability  $p$ . Also, if the System is in the State ON at some time instant, then, at the next instant it will be at the State OFF with probability  $q$ .

Describe this System's  $n$ -step Transition Probability Matrix.

# Example

**Solution:** Recall that the (1-step) Transition Matrix of our System is given by

$$P = \begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix}.$$

# Example

**Solution:** Recall that the (1-step) Transition Matrix of our System is given by

$$P = \begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix}.$$

Then, by the Chapman-Kolmogorov Theorem, the  $n$ -step Transition Matrix of our System is

$$P^{(n)} = P^n =$$



# Example

**Solution:** Recall that the (1-step) Transition Matrix of our System is given by

$$P = \begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix}.$$

Then, by the Chapman-Kolmogorov Theorem, the  $n$ -step Transition Matrix of our System is

$$P^{(n)} = P^n = \frac{1}{p+q} \cdot \begin{bmatrix} q & p \\ q & p \end{bmatrix} + \frac{(1-p-q)^n}{p+q} \cdot \begin{bmatrix} p & -p \\ -q & q \end{bmatrix}.$$

# Example

**Solution:** Recall that the (1-step) Transition Matrix of our System is given by

$$P = \begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix}.$$

Then, by the Chapman-Kolmogorov Theorem, the  $n$ -step Transition Matrix of our System is

$$P^{(n)} = P^n = \frac{1}{p+q} \cdot \begin{bmatrix} q & p \\ q & p \end{bmatrix} + \frac{(1-p-q)^n}{p+q} \cdot \begin{bmatrix} p & -p \\ -q & q \end{bmatrix}.$$

For example,

$$\mathbb{P}(X_4 = 1 | X_0 = 0) = p_{01}^{(4)} =$$

# Example

**Solution:** Recall that the (1-step) Transition Matrix of our System is given by

$$P = \begin{bmatrix} 1-p & p \\ q & 1-q \end{bmatrix}.$$

Then, by the Chapman-Kolmogorov Theorem, the  $n$ -step Transition Matrix of our System is

$$P^{(n)} = P^n = \frac{1}{p+q} \cdot \begin{bmatrix} q & p \\ q & p \end{bmatrix} + \frac{(1-p-q)^n}{p+q} \cdot \begin{bmatrix} p & -p \\ -q & q \end{bmatrix}.$$

For example,

$$\mathbb{P}(X_4 = 1 | X_0 = 0) = p_{01}^{(4)} = \frac{p}{p+q} - \frac{p \cdot (1-p-q)^4}{p+q}.$$

# Example

**Example:** This is a Klassik example: Usage of MC in Sociology.

# Example

**Example:** This is a Klassik example: Usage of MC in Sociology. In sociology, sometimes one classifies people by their income into 3 classes: low-income (lower class, 1), middle-income (middle class, 2) and high-income (upper class, 3).

# Example

**Example:** This is a Klassik example: Usage of MC in Sociology. In sociology, sometimes one classifies people by their income into 3 classes: low-income (lower class, 1), middle-income (middle class, 2) and high-income (upper class, 3). Sociologists have found that one of the important features determining the income class of a person is the income class of his/her parents/family.

# Example

**Example:** This is a Klassik example: Usage of MC in Sociology. In sociology, sometimes one classify people by their income into 3 classes: low-income (lower class, 1), middle-income (middle class, 2) and high-income (upper class, 3). Sociologists have found that one of the important features determining the income class of a person is the income class of his/her parents/family. The Transition Probabilities Matrix is given by

		Next Gen		
Current Gen	State	1	2	3
	1	0.65	0.28	0.07
	2	0.15	0.67	0.18
	3	0.12	0.36	0.52

# Example:

Now, the Transition Matrix (TPM) is

$$P = \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$



# Example:

Now, the Transition Matrix (TPM) is

$$P = \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$

Let us consider several problems:

# Example:

Now, the Transition Matrix (TPM) is

$$P = \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$

Let us consider several problems:

**Problem 1:** What is the probability that the grand-child of an income 1 class family will be an income 3 class person?

## Example:

Now, the Transition Matrix (TPM) is

$$P = \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$

Let us consider several problems:

**Problem 1:** What is the probability that the grand-child of an income 1 class family will be an income 3 class person?

Mathematically, or, MC-ally, we need to calculate  $p_{13}^{(2)}$ . To that end we calculate the row1-column3 element of  $P^2$ . (OTB?)

## Example:

Now, the Transition Matrix (TPM) is

$$P = \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$

Let us consider several problems:

**Problem 1:** What is the probability that the grand-child of an income 1 class family will be an income 3 class person?

Mathematically, or, MC-ally, we need to calculate  $p_{13}^{(2)}$ . To that end we calculate the row1-column3 element of  $P^2$ . (OTB?)

**Problem 2:** Given the initial distribution of income classes:

$$\pi_0 = [0.25, 0.6, 0.15],$$

(i.e., the proportion of class 1 families is 0.25, class 2 families is 0.6),

## Example:

Now, the Transition Matrix (TPM) is

$$P = \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$

Let us consider several problems:

**Problem 1:** What is the probability that the grand-child of an income 1 class family will be an income 3 class person?

Mathematically, or, MC-ally, we need to calculate  $p_{13}^{(2)}$ . To that end we calculate the row1-column3 element of  $P^2$ . (OTB?)

**Problem 2:** Given the initial distribution of income classes:

$$\pi_0 = [0.25, 0.6, 0.15],$$

(i.e., the proportion of class 1 families is 0.25, class 2 families is 0.6), find the distribution of the income classes after 2 generations.

# Example:

It is not too hard to see that after 1 generation, the distribution of families into income classes will be

$$\pi_1 = \pi_0 \cdot P =$$

## Example:

It is not too hard to see that after 1 generation, the distribution of families into income classes will be

$$\pi_1 = \pi_0 \cdot P = [0.25, 0.6, 0.15] \cdot \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$

## Example:

It is not too hard to see that after 1 generation, the distribution of families into income classes will be

$$\pi_1 = \pi_0 \cdot P = [0.25, 0.6, 0.15] \cdot \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$

And, to calculate the distribution of families into income classes after 2 generations will be

$$\pi_2 = \pi_0 \cdot P^2.$$



## Example:

It is not too hard to see that after 1 generation, the distribution of families into income classes will be

$$\pi_1 = \pi_0 \cdot P = [0.25, 0.6, 0.15] \cdot \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$

And, to calculate the distribution of families into income classes after 2 generations will be

$$\pi_2 = \pi_0 \cdot P^2.$$

And, in general, the Probability Distribution of the classes after  $n$  steps (generations) will be

$$\pi_n = \pi_0 \cdot P^n.$$

## Example:

It is not too hard to see that after 1 generation, the distribution of families into income classes will be

$$\pi_1 = \pi_0 \cdot P = [0.25, 0.6, 0.15] \cdot \begin{bmatrix} 0.65 & 0.28 & 0.07 \\ 0.15 & 0.67 & 0.18 \\ 0.12 & 0.36 & 0.52 \end{bmatrix}.$$

And, to calculate the distribution of families into income classes after 2 generations will be

$$\pi_2 = \pi_0 \cdot P^2.$$

And, in general, the Probability Distribution of the classes after  $n$  steps (generations) will be

$$\pi_n = \pi_0 \cdot P^n.$$

To be continued ...

# Long Time Behavior of MC

The next important problem is to study the long-time behavior of the MC.

# Long Time Behavior of MC

The next important problem is to study the long-time behavior of the MC. That is, to consider what will happen if  $n \rightarrow +\infty$ .

# Long Time Behavior of MC

The next important problem is to study the long-time behavior of the MC. That is, to consider what will happen if  $n \rightarrow +\infty$ .

**Example:** Consider again our old friend ON/OFF System as a MC. Recall that the  $n$ -step Transition Probability Matrix is

$$P^{(n)} = P^n =$$

# Long Time Behavior of MC

The next important problem is to study the long-time behavior of the MC. That is, to consider what will happen if  $n \rightarrow +\infty$ .

**Example:** Consider again our old friend ON/OFF System as a MC. Recall that the  $n$ -step Transition Probability Matrix is

$$P^{(n)} = P^n = \frac{1}{p+q} \cdot \begin{bmatrix} q & p \\ q & p \end{bmatrix} + \frac{(1-p-q)^n}{p+q} \cdot \begin{bmatrix} p & -p \\ -q & q \end{bmatrix}.$$

# Long Time Behavior of MC

The next important problem is to study the long-time behavior of the MC. That is, to consider what will happen if  $n \rightarrow +\infty$ .

**Example:** Consider again our old friend ON/OFF System as a MC. Recall that the  $n$ -step Transition Probability Matrix is

$$P^{(n)} = P^n = \frac{1}{p+q} \cdot \begin{bmatrix} q & p \\ q & p \end{bmatrix} + \frac{(1-p-q)^n}{p+q} \cdot \begin{bmatrix} p & -p \\ -q & q \end{bmatrix}.$$

Now, if  $n \rightarrow +\infty$ , then (since  $|1-p-q| < 1$ , except the cases when  $p = q = 0$  or  $p = q = 1$ )

$$P^{(\infty)} =$$

# Long Time Behavior of MC

The next important problem is to study the long-time behavior of the MC. That is, to consider what will happen if  $n \rightarrow +\infty$ .

**Example:** Consider again our old friend ON/OFF System as a MC. Recall that the  $n$ -step Transition Probability Matrix is

$$P^{(n)} = P^n = \frac{1}{p+q} \cdot \begin{bmatrix} q & p \\ q & p \end{bmatrix} + \frac{(1-p-q)^n}{p+q} \cdot \begin{bmatrix} p & -p \\ -q & q \end{bmatrix}.$$

Now, if  $n \rightarrow +\infty$ , then (since  $|1-p-q| < 1$ , except the cases when  $p = q = 0$  or  $p = q = 1$ )

$$P^{(\infty)} = \frac{1}{p+q} \cdot \begin{bmatrix} q & p \\ q & p \end{bmatrix}.$$



# Long Time Behavior of MC

The next important problem is to study the long-time behavior of the MC. That is, to consider what will happen if  $n \rightarrow +\infty$ .

**Example:** Consider again our old friend ON/OFF System as a MC. Recall that the  $n$ -step Transition Probability Matrix is

$$P^{(n)} = P^n = \frac{1}{p+q} \cdot \begin{bmatrix} q & p \\ q & p \end{bmatrix} + \frac{(1-p-q)^n}{p+q} \cdot \begin{bmatrix} p & -p \\ -q & q \end{bmatrix}.$$

Now, if  $n \rightarrow +\infty$ , then (since  $|1-p-q| < 1$ , except the cases when  $p = q = 0$  or  $p = q = 1$ )

$$P^{(\infty)} = \frac{1}{p+q} \cdot \begin{bmatrix} q & p \\ q & p \end{bmatrix}.$$

This means, that in the long run, the Probability to be at the State 0 is approximately  $\frac{q}{p+q}$ , and the Probability to be at the State 1 is approximately  $\frac{p}{p+q}$ , irrespective of the initial State!

# The Long Time Behavior of MC

Now, let us consider the general case.

# The Long Time Behavior of MC

Now, let us consider the general case. Assume we have a MC with State Space  $S = \{1, 2, \dots, N\}$ , the TPM  $P = [p_{ij}]$  and IDS (Initial Distribution of States)  $\pi_0 = [\mathbb{P}(X_0 = i)]_{i=1, \dots, N}$ .

# The Long Time Behavior of MC

Now, let us consider the general case. Assume we have a MC with State Space  $S = \{1, 2, \dots, N\}$ , the TPM  $P = [p_{ij}]$  and IDS (Initial Distribution of States)  $\pi_0 = [\mathbb{P}(X_0 = i)]_{i=1, \dots, N}$ .

Then, it can be proved that the distribution of the State Probabilities  $\pi_n$ , after  $n$  steps, will be

$$\pi_n = \pi_0 \cdot P^n, \quad \forall n \in \mathbb{N}.$$

# The Long Time Behavior of MC

Now, let us consider the general case. Assume we have a MC with State Space  $S = \{1, 2, \dots, N\}$ , the TPM  $P = [p_{ij}]$  and IDS (Initial Distribution of States)  $\pi_0 = [\mathbb{P}(X_0 = i)]_{i=1, \dots, N}$ .

Then, it can be proved that the distribution of the State Probabilities  $\pi_n$ , after  $n$  steps, will be

$$\pi_n = \pi_0 \cdot P^n, \quad \forall n \in \mathbb{N}.$$

or, which is the same,

$$\pi_n = \pi_{n-1} \cdot P, \quad \forall n \in \mathbb{N}.$$

# The Long Time Behavior of MC

Now, let us consider the general case. Assume we have a MC with State Space  $S = \{1, 2, \dots, N\}$ , the TPM  $P = [p_{ij}]$  and IDS (Initial Distribution of States)  $\pi_0 = [\mathbb{P}(X_0 = i)]_{i=1, \dots, N}$ .

Then, it can be proved that the distribution of the State Probabilities  $\pi_n$ , after  $n$  steps, will be

$$\pi_n = \pi_0 \cdot P^n, \quad \forall n \in \mathbb{N}.$$

or, which is the same,

$$\pi_n = \pi_{n-1} \cdot P, \quad \forall n \in \mathbb{N}.$$

Now, it turns out, that, under some conditions on the TPM  $P$ , the sequence  $\pi_n$  will converge to some vector  $\pi$ ,  $\pi_n \rightarrow \pi$ ,

# The Long Time Behavior of MC

Now, let us consider the general case. Assume we have a MC with State Space  $S = \{1, 2, \dots, N\}$ , the TPM  $P = [p_{ij}]$  and IDS (Initial Distribution of States)  $\pi_0 = [\mathbb{P}(X_0 = i)]_{i=1, \dots, N}$ .

Then, it can be proved that the distribution of the State Probabilities  $\pi_n$ , after  $n$  steps, will be

$$\pi_n = \pi_0 \cdot P^n, \quad \forall n \in \mathbb{N}.$$

or, which is the same,

$$\pi_n = \pi_{n-1} \cdot P, \quad \forall n \in \mathbb{N}.$$

Now, it turns out, that, under some conditions on the TPM  $P$ , the sequence  $\pi_n$  will converge to some vector  $\pi$ ,  $\pi_n \rightarrow \pi$ , which is called the **Stationary (or the Equilibrium) Distribution of MC**.

# The Long Time Behavior of MC

To find the Stationary Distribution  $\pi$ , we use

$$\pi_n = \pi_{n-1} \cdot P, \quad \forall n \in \mathbb{N},$$

and pass to the limit  $n \rightarrow \infty$ :



# The Long Time Behavior of MC

To find the Stationary Distribution  $\pi$ , we use

$$\pi_n = \pi_{n-1} \cdot P, \quad \forall n \in \mathbb{N},$$

and pass to the limit  $n \rightarrow \infty$ : as  $\pi_n \rightarrow \pi$ , we'll get

# The Long Time Behavior of MC

To find the Stationary Distribution  $\pi$ , we use

$$\pi_n = \pi_{n-1} \cdot P, \quad \forall n \in \mathbb{N},$$

and pass to the limit  $n \rightarrow \infty$ : as  $\pi_n \rightarrow \pi$ , we'll get

$$\pi = \pi \cdot P.$$

# The Long Time Behavior of MC

To find the Stationary Distribution  $\pi$ , we use

$$\pi_n = \pi_{n-1} \cdot P, \quad \forall n \in \mathbb{N},$$

and pass to the limit  $n \rightarrow \infty$ : as  $\pi_n \rightarrow \pi$ , we'll get

$$\pi = \pi \cdot P.$$

**Important Note 1:** It is important that the Stationary Distribution  $\pi$  is *independent* of the initial Distribution  $\pi_0$ .

# The Long Time Behavior of MC

To find the Stationary Distribution  $\pi$ , we use

$$\pi_n = \pi_{n-1} \cdot P, \quad \forall n \in \mathbb{N},$$

and pass to the limit  $n \rightarrow \infty$ : as  $\pi_n \rightarrow \pi$ , we'll get

$$\pi = \pi \cdot P.$$

**Important Note 1:** It is important that the Stationary Distribution  $\pi$  is *independent* of the initial Distribution  $\pi_0$ . So for any Initial Distribution of States Probabilities, after some time the Distribution will stabilize to the same Distribution.

# The Long Time Behavior of MC

# The Long Time Behavior of MC

**Important Note 2:** To calculate the Stationary Distribution of States, we need to solve the above equation

$$\pi = \pi \cdot P.$$

# The Long Time Behavior of MC

**Important Note 2:** To calculate the Stationary Distribution of States, we need to solve the above equation

$$\pi = \pi \cdot P.$$

Here  $\pi$  is an  $N$ -dim vector,

$$\pi = [\mathbb{P}(X_\infty = 1), \dots, \mathbb{P}(X_\infty = N)],$$

and  $P$  is the  $N \times N$  TPM, and we have a SLE to solve.

# The Long Time Behavior of MC

**Important Note 2:** To calculate the Stationary Distribution of States, we need to solve the above equation

$$\pi = \pi \cdot P.$$

Here  $\pi$  is an  $N$ -dim vector,

$$\pi = [\mathbb{P}(X_\infty = 1), \dots, \mathbb{P}(X_\infty = N)],$$

and  $P$  is the  $N \times N$  TPM, and we have a SLE to solve. But it turns out that the rank of this system is less than  $N$ , so there are infinitely many solutions to this SLE.



# The Long Time Behavior of MC

**Important Note 2:** To calculate the Stationary Distribution of States, we need to solve the above equation

$$\pi = \pi \cdot P.$$

Here  $\pi$  is an  $N$ -dim vector,

$$\pi = \left[ \mathbb{P}(X_\infty = 1), \dots, \mathbb{P}(X_\infty = N) \right],$$

and  $P$  is the  $N \times N$  TPM, and we have a SLE to solve. But it turns out that the rank of this system is less than  $N$ , so there are infinitely many solutions to this SLE. But we need to remember that  $\pi$  needs to be a Probability Distribution, so the sum of its elements needs to be 1. Adding this to our SLE gives a unique solution.

# Example

In details, assume that the Stationary Distribution is given by

$$\pi = [x_1, x_2, \dots, x_N].$$

# Example

In details, assume that the Stationary Distribution is given by

$$\pi = [x_1, x_2, \dots, x_N].$$

Then, to find  $x_k$ -s, we need to solve

$$\begin{cases} [x_1, x_2, \dots, x_N] = [x_1, x_2, \dots, x_N] \cdot P \\ x_1 + x_2 + \dots + x_N = 1 \end{cases}$$

# Example

In details, assume that the Stationary Distribution is given by

$$\pi = [x_1, x_2, \dots, x_N].$$

Then, to find  $x_k$ -s, we need to solve

$$\begin{cases} [x_1, x_2, \dots, x_N] = [x_1, x_2, \dots, x_N] \cdot P \\ x_1 + x_2 + \dots + x_N = 1 \end{cases}$$

**Example:** GOF ON/OFF System. Find the Stationary Distribution of this MC.

**Solution:** OTB

# Example

In details, assume that the Stationary Distribution is given by

$$\pi = [x_1, x_2, \dots, x_N].$$

Then, to find  $x_k$ -s, we need to solve

$$\begin{cases} [x_1, x_2, \dots, x_N] = [x_1, x_2, \dots, x_N] \cdot P \\ x_1 + x_2 + \dots + x_N = 1 \end{cases}$$

**Example:** GOF ON/OFF System. Find the Stationary Distribution of this MC.

**Solution:** OTB

**Example:** In the Income Classes Example, find the Stationary Distribution of States and give an interpretation.

**Solution:** OTB

# Existence/Uniqueness of Stationary Distribution

It is not a fact that the Stationary Distribution Exists for any MC, or, in the case it exists, it is not a fact that it is unique.

# Existence/Uniqueness of Stationary Distribution

It is not a fact that the Stationary Distribution Exists for any MC, or, in the case it exists, it is not a fact that it is unique. The next Theorem states a condition under which the Stationary Distribution exists and is unique.

# Existence/Uniqueness of Stationary Distribution

It is not a fact that the Stationary Distribution Exists for any MC, or, in the case it exists, it is not a fact that it is unique. The next Theorem states a condition under which the Stationary Distribution exists and is unique.

We will say that our MC is Irreducible, if for any  $i, j \in S$ , there exists an  $n \in \mathbb{N}$  such that

$$p_{ij}^{(n)} > 0.$$



# Existence/Uniqueness of Stationary Distribution

It is not a fact that the Stationary Distribution Exists for any MC, or, in the case it exists, it is not a fact that it is unique. The next Theorem states a condition under which the Stationary Distribution exists and is unique.

We will say that our MC is Irreducible, if for any  $i, j \in S$ , there exists an  $n \in \mathbb{N}$  such that

$$p_{ij}^{(n)} > 0.$$

If  $X_1, X_2, \dots$  is a MC with Finite State Space, and if it is Irreducible, then it has a unique Stationary Distribution.

# Existence/Uniqueness of Stationary Distribution

It is not a fact that the Stationary Distribution Exists for any MC, or, in the case it exists, it is not a fact that it is unique. The next Theorem states a condition under which the Stationary Distribution exists and is unique.

We will say that our MC is Irreducible, if for any  $i, j \in S$ , there exists an  $n \in \mathbb{N}$  such that

$$p_{ij}^{(n)} > 0.$$

If  $X_1, X_2, \dots$  is a MC with Finite State Space, and if it is Irreducible, then it has a unique Stationary Distribution.

**Idea:** If our MC is Irreducible, then the initial State Probabilities are thoroughly mixed.

# PRTS! URAA!

