

# AUA CS 108, Statistics, Fall 2019

## Lecture 14

Michael Poghosyan

YSU, AUA

[michael@ysu.am](mailto:michael@ysu.am), [mpoghosyan@aua.am](mailto:mpoghosyan@aua.am)

25 Sep 2019

# Contents

- ▶ Important Discrete Distributions
- ▶ Important Continuous Distributions
- ▶ Convergence Types of R.V. Sequences

# Last Lecture ReCap

- ▶ Why we use the Sample Covariance?

# Last Lecture ReCap

- ▶ Why we use the Sample Covariance?
- ▶ Why we use the Sample Correlation Coefficient?

# Important Discrete Distributions

# Bernoulli Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )

## Bernoulli Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Bernoulli}(p)$ ;

## Bernoulli Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Bernoulli}(p)$ ;
- ▶ Support:  $\{0, 1\}$



## Bernoulli Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Bernoulli}(p)$ ;
- ▶ Support:  $\{0, 1\}$
- ▶ PMF:

Values of $X$	0	1
$\mathbb{P}(X = x)$	$1 - p$	$p$

## Bernoulli Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Bernoulli}(p)$ ;
- ▶ Support:  $\{0, 1\}$
- ▶ PMF:

Values of $X$	0	1
$\mathbb{P}(X = x)$	$1 - p$	$p$

**Note:** This can be written in the form:

$$f(x) = f(x; p) = f(x|p) = p^x \cdot (1 - p)^{1-x}, \quad x \in \{0, 1\}.$$

## Bernoulli Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Bernoulli}(p)$ ;
- ▶ Support:  $\{0, 1\}$
- ▶ PMF:

Values of $X$	0	1
$\mathbb{P}(X = x)$	$1 - p$	$p$

**Note:** This can be written in the form:

$$f(x) = f(x; p) = f(x|p) = p^x \cdot (1 - p)^{1-x}, \quad x \in \{0, 1\}.$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$ .

## Bernoulli Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Bernoulli}(p)$ ;
- ▶ Support:  $\{0, 1\}$
- ▶ PMF:

Values of $X$	0	1
$\mathbb{P}(X = x)$	$1 - p$	$p$

**Note:** This can be written in the form:

$$f(x) = f(x; p) = f(x|p) = p^x \cdot (1 - p)^{1-x}, \quad x \in \{0, 1\}.$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$ .
- ▶ Models: Models binary output, “success-failure” type Experiments, a lot of examples.

## Bernoulli Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Bernoulli}(p)$ ;
- ▶ Support:  $\{0, 1\}$
- ▶ PMF:

Values of $X$	0	1
$\mathbb{P}(X = x)$	$1 - p$	$p$

**Note:** This can be written in the form:

$$f(x) = f(x; p) = f(x|p) = p^x \cdot (1 - p)^{1-x}, \quad x \in \{0, 1\}.$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$ .
- ▶ Models: Models binary output, “success-failure” type Experiments, a lot of examples.
- ▶ **R** name: `binom` with the parameters `size=1` and `prob`

## Bernoulli Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Bernoulli}(p)$ ;
- ▶ Support:  $\{0, 1\}$
- ▶ PMF:

Values of $X$	0	1
$\mathbb{P}(X = x)$	$1 - p$	$p$

**Note:** This can be written in the form:

$$f(x) = f(x; p) = f(x|p) = p^x \cdot (1 - p)^{1-x}, \quad x \in \{0, 1\}.$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = p$ ,  $\text{Var}(X) = p(1 - p)$ .
- ▶ Models: Models binary output, “success-failure” type Experiments, a lot of examples.
- ▶ **R** name: `binom` with the parameters `size=1` and `prob`

# Binomial Distribution

- ▶ Parameters:  $n \in \mathbb{N}$ ,  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )

## Binomial Distribution

- ▶ Parameters:  $n \in \mathbb{N}$ ,  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Binom}(n, p)$ ;



## Binomial Distribution

- ▶ Parameters:  $n \in \mathbb{N}$ ,  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Binom}(n, p)$ ;
- ▶ Support:  $\{0, 1, 2, \dots, n\}$

# Binomial Distribution

- ▶ Parameters:  $n \in \mathbb{N}$ ,  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Binom}(n, p)$ ;
- ▶ Support:  $\{0, 1, 2, \dots, n\}$
- ▶ PMF:

Values of $X$	0	1	...	$k$	...	$n$
$\mathbb{P}(X = x)$	$\binom{n}{0} p^0 (1-p)^{n-0}$	$\binom{n}{1} p^1 (1-p)^{n-1}$	...	$\binom{n}{k} p^k (1-p)^{n-k}$	...	$\binom{n}{n} p^n (1-p)^{n-n}$

# Binomial Distribution

- ▶ Parameters:  $n \in \mathbb{N}$ ,  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Binom}(n, p)$ ;
- ▶ Support:  $\{0, 1, 2, \dots, n\}$
- ▶ PMF:

Values of $X$	0	1	...	$k$	...	$n$
$\mathbb{P}(X = x)$	$\binom{n}{0} p^0 (1-p)^{n-0}$	$\binom{n}{1} p^1 (1-p)^{n-1}$	...	$\binom{n}{k} p^k (1-p)^{n-k}$	...	$\binom{n}{n} p^n (1-p)^{n-n}$

- ▶ Mean and Variance:  $\mathbb{E}(X) = n \cdot p$ ,  $\text{Var}(X) = n \cdot p(1-p)$ .

# Binomial Distribution

- ▶ Parameters:  $n \in \mathbb{N}$ ,  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Binom}(n, p)$ ;
- ▶ Support:  $\{0, 1, 2, \dots, n\}$
- ▶ PMF:

Values of $X$	0	1	...	$k$	...	$n$
$\mathbb{P}(X = x)$	$\binom{n}{0} p^0 (1-p)^{n-0}$	$\binom{n}{1} p^1 (1-p)^{n-1}$	...	$\binom{n}{k} p^k (1-p)^{n-k}$	...	$\binom{n}{n} p^n (1-p)^{n-n}$

- ▶ Mean and Variance:  $\mathbb{E}(X) = n \cdot p$ ,  $\text{Var}(X) = n \cdot p(1-p)$ .
- ▶ Models: Models the independent repetition of the *Bernoulli*( $p$ ) Experiment.

# Binomial Distribution

- ▶ Parameters:  $n \in \mathbb{N}$ ,  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Binom}(n, p)$ ;
- ▶ Support:  $\{0, 1, 2, \dots, n\}$
- ▶ PMF:

Values of $X$	0	1	...	$k$	...	$n$
$\mathbb{P}(X = x)$	$\binom{n}{0} p^0 (1-p)^{n-0}$	$\binom{n}{1} p^1 (1-p)^{n-1}$	...	$\binom{n}{k} p^k (1-p)^{n-k}$	...	$\binom{n}{n} p^n (1-p)^{n-n}$

- ▶ Mean and Variance:  $\mathbb{E}(X) = n \cdot p$ ,  $\text{Var}(X) = n \cdot p(1-p)$ .
- ▶ Models: Models the independent repetition of the *Bernoulli*( $p$ ) Experiment.
- ▶ **R** name: `binom` with the parameters `size` and `prob`

# Binomial Distribution

- ▶ Parameters:  $n \in \mathbb{N}$ ,  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Binom}(n, p)$ ;
- ▶ Support:  $\{0, 1, 2, \dots, n\}$
- ▶ PMF:

Values of $X$	0	1	...	$k$	...	$n$
$\mathbb{P}(X = x)$	$\binom{n}{0} p^0 (1-p)^{n-0}$	$\binom{n}{1} p^1 (1-p)^{n-1}$	...	$\binom{n}{k} p^k (1-p)^{n-k}$	...	$\binom{n}{n} p^n (1-p)^{n-n}$

- ▶ Mean and Variance:  $\mathbb{E}(X) = n \cdot p$ ,  $\text{Var}(X) = n \cdot p(1-p)$ .
- ▶ Models: Models the independent repetition of the *Bernoulli*( $p$ ) Experiment.
- ▶ **R** name: `binom` with the parameters `size` and `prob`
- ▶ Additional: If  $X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$  are independent, then  $X_1 + X_2 + \dots + X_n \sim \text{Binom}(n, p)$ .

# Binomial Distribution

- ▶ Parameters:  $n \in \mathbb{N}$ ,  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Binom}(n, p)$ ;
- ▶ Support:  $\{0, 1, 2, \dots, n\}$
- ▶ PMF:

Values of $X$	0	1	...	$k$	...	$n$
$\mathbb{P}(X = x)$	$\binom{n}{0} p^0 (1-p)^{n-0}$	$\binom{n}{1} p^1 (1-p)^{n-1}$	...	$\binom{n}{k} p^k (1-p)^{n-k}$	...	$\binom{n}{n} p^n (1-p)^{n-n}$

- ▶ Mean and Variance:  $\mathbb{E}(X) = n \cdot p$ ,  $\text{Var}(X) = n \cdot p(1-p)$ .
- ▶ Models: Models the independent repetition of the *Bernoulli*( $p$ ) Experiment.
- ▶ **R** name: `binom` with the parameters `size` and `prob`
- ▶ Additional: If  $X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$  are independent, then  $X_1 + X_2 + \dots + X_n \sim \text{Binom}(n, p)$ .
- ▶ Example:

## Geometric Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )



## Geometric Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Geom}(p)$ ;

## Geometric Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Geom}(p)$ ;
- ▶ Support:  $\{1, 2, 3, \dots\}$  (or, sometimes,  $\{0, 1, 2, 3, \dots\}$  )

# Geometric Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Geom}(p)$ ;
- ▶ Support:  $\{1, 2, 3, \dots\}$  (or, sometimes,  $\{0, 1, 2, 3, \dots\}$  )
- ▶ PMF:

Values of $X$	1	2	3	...
$\mathbb{P}(X = x)$	$p$	$p(1 - p)$	$p(1 - p)^2$	...

# Geometric Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Geom}(p)$ ;
- ▶ Support:  $\{1, 2, 3, \dots\}$  (or, sometimes,  $\{0, 1, 2, 3, \dots\}$  )
- ▶ PMF:

Values of $X$	1	2	3	...
$\mathbb{P}(X = x)$	$p$	$p(1 - p)$	$p(1 - p)^2$	...

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{1}{p}$ ,  $\text{Var}(X) = \frac{1 - p}{p^2}$ .

## Geometric Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Geom}(p)$ ;
- ▶ Support:  $\{1, 2, 3, \dots\}$  (or, sometimes,  $\{0, 1, 2, 3, \dots\}$  )
- ▶ PMF:

Values of $X$	1	2	3	...
$\mathbb{P}(X = x)$	$p$	$p(1 - p)$	$p(1 - p)^2$	...

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{1}{p}$ ,  $\text{Var}(X) = \frac{1 - p}{p^2}$ .
- ▶ Models: Models the independent repetition of the *Bernoulli*( $p$ ) Experiment until the *First Success*.

## Geometric Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Geom}(p)$ ;
- ▶ Support:  $\{1, 2, 3, \dots\}$  (or, sometimes,  $\{0, 1, 2, 3, \dots\}$  )
- ▶ PMF:

Values of $X$	1	2	3	...
$\mathbb{P}(X = x)$	$p$	$p(1 - p)$	$p(1 - p)^2$	...

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{1}{p}$ ,  $\text{Var}(X) = \frac{1 - p}{p^2}$ .
- ▶ Models: Models the independent repetition of the *Bernoulli*( $p$ ) Experiment until the *First Success*.
- ▶ **R** name: `geom` with the parameter `prob`

## Geometric Distribution

- ▶ Parameter:  $p \in [0, 1]$  (usually,  $p \in (0, 1)$ )
- ▶ Notation:  $X \sim \text{Geom}(p)$ ;
- ▶ Support:  $\{1, 2, 3, \dots\}$  (or, sometimes,  $\{0, 1, 2, 3, \dots\}$  )
- ▶ PMF:

Values of $X$	1	2	3	...
$\mathbb{P}(X = x)$	$p$	$p(1 - p)$	$p(1 - p)^2$	...

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{1}{p}$ ,  $\text{Var}(X) = \frac{1 - p}{p^2}$ .
- ▶ Models: Models the independent repetition of the *Bernoulli*( $p$ ) Experiment until the *First Success*.
- ▶ **R** name: geom with the parameter prob
- ▶ Example:

```
rgeom(10,prob = 0.3)
```

```
## [1] 1 2 6 0 2 4 4 3 1 4
```

# Poisson Distribution

- ▶ Parameter:  $\lambda > 0$



## Poisson Distribution

- ▶ Parameter:  $\lambda > 0$
- ▶ Notation:  $X \sim \text{Pois}(\lambda)$ ;

# Poisson Distribution

- ▶ Parameter:  $\lambda > 0$
- ▶ Notation:  $X \sim \text{Pois}(\lambda)$ ;
- ▶ Support:  $\{0, 1, 2, 3, \dots\}$

# Poisson Distribution

- ▶ Parameter:  $\lambda > 0$
- ▶ Notation:  $X \sim \text{Pois}(\lambda)$ ;
- ▶ Support:  $\{0, 1, 2, 3, \dots\}$
- ▶ PMF:

Values of $X$	0	1	2	...
$\mathbb{P}(X = x)$	$e^{-\lambda} \frac{\lambda^0}{0!}$	$e^{-\lambda} \frac{\lambda^1}{1!}$	$e^{-\lambda} \frac{\lambda^2}{2!}$	...

# Poisson Distribution

- ▶ Parameter:  $\lambda > 0$
- ▶ Notation:  $X \sim \text{Pois}(\lambda)$ ;
- ▶ Support:  $\{0, 1, 2, 3, \dots\}$
- ▶ PMF:

Values of $X$	0	1	2	...
$\mathbb{P}(X = x)$	$e^{-\lambda} \frac{\lambda^0}{0!}$	$e^{-\lambda} \frac{\lambda^1}{1!}$	$e^{-\lambda} \frac{\lambda^2}{2!}$	...

- ▶ Mean and Variance:  $\mathbb{E}(X) = \lambda$ ,  $\text{Var}(X) = \lambda$ .

# Poisson Distribution

- ▶ Parameter:  $\lambda > 0$
- ▶ Notation:  $X \sim \text{Pois}(\lambda)$ ;
- ▶ Support:  $\{0, 1, 2, 3, \dots\}$
- ▶ PMF:

Values of $X$	0	1	2	...
$\mathbb{P}(X = x)$	$e^{-\lambda} \frac{\lambda^0}{0!}$	$e^{-\lambda} \frac{\lambda^1}{1!}$	$e^{-\lambda} \frac{\lambda^2}{2!}$	...

- ▶ Mean and Variance:  $\mathbb{E}(X) = \lambda$ ,  $\text{Var}(X) = \lambda$ .
- ▶ Models: Models the number of success in a fixed interval. Say, number of calls, number of customers, number of clicks, number of page visits, ...

# Poisson Distribution

- ▶ Parameter:  $\lambda > 0$
- ▶ Notation:  $X \sim \text{Pois}(\lambda)$ ;
- ▶ Support:  $\{0, 1, 2, 3, \dots\}$
- ▶ PMF:

Values of $X$	0	1	2	...
$\mathbb{P}(X = x)$	$e^{-\lambda} \frac{\lambda^0}{0!}$	$e^{-\lambda} \frac{\lambda^1}{1!}$	$e^{-\lambda} \frac{\lambda^2}{2!}$	...

- ▶ Mean and Variance:  $\mathbb{E}(X) = \lambda$ ,  $\text{Var}(X) = \lambda$ .
- ▶ Models: Models the number of success in a fixed interval. Say, number of calls, number of customers, number of clicks, number of page visits, ...  $\lambda$  is the average number of calls, customers, clicks, page visits, ...

# Poisson Distribution

- ▶ Parameter:  $\lambda > 0$
- ▶ Notation:  $X \sim \text{Pois}(\lambda)$ ;
- ▶ Support:  $\{0, 1, 2, 3, \dots\}$
- ▶ PMF:

Values of $X$	0	1	2	...
$\mathbb{P}(X = x)$	$e^{-\lambda} \frac{\lambda^0}{0!}$	$e^{-\lambda} \frac{\lambda^1}{1!}$	$e^{-\lambda} \frac{\lambda^2}{2!}$	...

- ▶ Mean and Variance:  $\mathbb{E}(X) = \lambda$ ,  $\text{Var}(X) = \lambda$ .
- ▶ Models: Models the number of success in a fixed interval. Say, number of calls, number of customers, number of clicks, number of page visits, ...  $\lambda$  is the average number of calls, customers, clicks, page visits,...
- ▶ **R** name: `pois` with the parameter `lambda`

# Poisson Distribution

- ▶ Parameter:  $\lambda > 0$
- ▶ Notation:  $X \sim \text{Pois}(\lambda)$ ;
- ▶ Support:  $\{0, 1, 2, 3, \dots\}$
- ▶ PMF:

Values of $X$	0	1	2	...
$\mathbb{P}(X = x)$	$e^{-\lambda} \frac{\lambda^0}{0!}$	$e^{-\lambda} \frac{\lambda^1}{1!}$	$e^{-\lambda} \frac{\lambda^2}{2!}$	...

- ▶ Mean and Variance:  $\mathbb{E}(X) = \lambda$ ,  $\text{Var}(X) = \lambda$ .
- ▶ Models: Models the number of success in a fixed interval. Say, number of calls, number of customers, number of clicks, number of page visits, ...  $\lambda$  is the average number of calls, customers, clicks, page visits,...
- ▶ **R** name: `pois` with the parameter `lambda`
- ▶ Example:

```
mpois(10, lambda = 2)
```



# Important Continuous Distributions

# Uniform Distribution

- ▶ Parameters:  $a, b$  ( $a < b$ )

## Uniform Distribution

- ▶ Parameters:  $a, b$  ( $a < b$ )
- ▶ Notation:  $X \sim \text{Unif}[a, b];$

# Uniform Distribution

- ▶ Parameters:  $a, b$  ( $a < b$ )
- ▶ Notation:  $X \sim \text{Unif}[a, b]$ ;
- ▶ Support:  $[a, b]$

# Uniform Distribution

- ▶ Parameters:  $a, b$  ( $a < b$ )
- ▶ Notation:  $X \sim \text{Unif}[a, b]$ ;
- ▶ Support:  $[a, b]$
- ▶ PDF:

$$f(x) = f(x|a, b) = \begin{cases} \frac{1}{b-a}, & x \in [a, b] \\ 0, & \text{otherwise} \end{cases}$$

# Uniform Distribution

- ▶ Parameters:  $a, b$  ( $a < b$ )
- ▶ Notation:  $X \sim \text{Unif}[a, b]$ ;
- ▶ Support:  $[a, b]$
- ▶ PDF:

$$f(x) = f(x|a, b) = \begin{cases} \frac{1}{b-a}, & x \in [a, b] \\ 0, & \text{otherwise} \end{cases}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ .

# Uniform Distribution

- ▶ Parameters:  $a, b$  ( $a < b$ )
- ▶ Notation:  $X \sim \text{Unif}[a, b]$ ;
- ▶ Support:  $[a, b]$
- ▶ PDF:

$$f(x) = f(x|a, b) = \begin{cases} \frac{1}{b-a}, & x \in [a, b] \\ 0, & \text{otherwise} \end{cases}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ .
- ▶ Models: Usually we think about the Uniform Distribution when talking about *picking a random number from an interval*

## Uniform Distribution

- ▶ Parameters:  $a, b$  ( $a < b$ )
- ▶ Notation:  $X \sim \text{Unif}[a, b]$ ;
- ▶ Support:  $[a, b]$
- ▶ PDF:

$$f(x) = f(x|a, b) = \begin{cases} \frac{1}{b-a}, & x \in [a, b] \\ 0, & \text{otherwise} \end{cases}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ .
- ▶ Models: Usually we think about the Uniform Distribution when talking about *picking a random number from an interval*
- ▶ **R** name: `unif` with the parameters `min = 0` and `max = 1`



## Uniform Distribution

- ▶ Parameters:  $a, b$  ( $a < b$ )
- ▶ Notation:  $X \sim \text{Unif}[a, b]$ ;
- ▶ Support:  $[a, b]$
- ▶ PDF:

$$f(x) = f(x|a, b) = \begin{cases} \frac{1}{b-a}, & x \in [a, b] \\ 0, & \text{otherwise} \end{cases}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{a+b}{2}$ ,  $\text{Var}(X) = \frac{(b-a)^2}{12}$ .
- ▶ Models: Usually we think about the Uniform Distribution when talking about *picking a random number from an interval*
- ▶ **R** name: `unif` with the parameters `min = 0` and `max = 1`
- ▶ Example:

```
runif(10, min = 2, max = 5)
```

## Exponential Distribution

- ▶ Parameter:  $\lambda > 0$  (rate) (or, sometimes,  $\beta = \frac{1}{\lambda}$ , scale)

## Exponential Distribution

- ▶ Parameter:  $\lambda > 0$  (rate) (or, sometimes,  $\beta = \frac{1}{\lambda}$ , scale)
- ▶ Notation:  $X \sim \text{Exp}(\lambda)$  (or  $\text{Exp}(\beta)$ );

## Exponential Distribution

- ▶ Parameter:  $\lambda > 0$  (rate) (or, sometimes,  $\beta = \frac{1}{\lambda}$ , scale)
- ▶ Notation:  $X \sim \text{Exp}(\lambda)$  (or  $\text{Exp}(\beta)$ );
- ▶ Support:  $[0, +\infty)$

# Exponential Distribution

- ▶ Parameter:  $\lambda > 0$  (rate) (or, sometimes,  $\beta = \frac{1}{\lambda}$ , scale)
- ▶ Notation:  $X \sim \text{Exp}(\lambda)$  (or  $\text{Exp}(\beta)$ );
- ▶ Support:  $[0, +\infty)$
- ▶ PDF:

$$f(x|\lambda) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

# Exponential Distribution

- ▶ Parameter:  $\lambda > 0$  (rate) (or, sometimes,  $\beta = \frac{1}{\lambda}$ , scale)
- ▶ Notation:  $X \sim \text{Exp}(\lambda)$  (or  $\text{Exp}(\beta)$ );
- ▶ Support:  $[0, +\infty)$
- ▶ PDF:

$$f(x|\lambda) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ .

# Exponential Distribution

- ▶ Parameter:  $\lambda > 0$  (rate) (or, sometimes,  $\beta = \frac{1}{\lambda}$ , scale)
- ▶ Notation:  $X \sim \text{Exp}(\lambda)$  (or  $\text{Exp}(\beta)$ );
- ▶ Support:  $[0, +\infty)$

- ▶ PDF:

$$f(x|\lambda) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ .
- ▶ Models: time elapsed until the occurrence of certain event, or the time between events (waiting times), when that time is random.

## Exponential Distribution

- ▶ Parameter:  $\lambda > 0$  (rate) (or, sometimes,  $\beta = \frac{1}{\lambda}$ , scale)
- ▶ Notation:  $X \sim \text{Exp}(\lambda)$  (or  $\text{Exp}(\beta)$ );
- ▶ Support:  $[0, +\infty)$

- ▶ PDF:

$$f(x|\lambda) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ .
- ▶ Models: time elapsed until the occurrence of certain event, or the time between events (waiting times), when that time is random.  $\lambda$  is the average “arrival rate”, the reciprocal of the average time between the events,  
$$\lambda = \frac{1}{\text{average time between events}}.$$



## Exponential Distribution

- ▶ Parameter:  $\lambda > 0$  (rate) (or, sometimes,  $\beta = \frac{1}{\lambda}$ , scale)
- ▶ Notation:  $X \sim \text{Exp}(\lambda)$  (or  $\text{Exp}(\beta)$ );
- ▶ Support:  $[0, +\infty)$

- ▶ PDF:

$$f(x|\lambda) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ .
- ▶ Models: time elapsed until the occurrence of certain event, or the time between events (waiting times), when that time is random.  $\lambda$  is the average “arrival rate”, the reciprocal of the average time between the events,  
$$\lambda = \frac{1}{\text{average time between events}}.$$
- ▶ **R** name: `exp` with the parameter `rate = 1`

# Exponential Distribution

- ▶ Parameter:  $\lambda > 0$  (rate) (or, sometimes,  $\beta = \frac{1}{\lambda}$ , scale)
- ▶ Notation:  $X \sim \text{Exp}(\lambda)$  (or  $\text{Exp}(\beta)$ );
- ▶ Support:  $[0, +\infty)$

- ▶ PDF:

$$f(x|\lambda) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \frac{1}{\lambda}$ ,  $\text{Var}(X) = \frac{1}{\lambda^2}$ .
- ▶ Models: time elapsed until the occurrence of certain event, or the time between events (waiting times), when that time is random.  $\lambda$  is the average “arrival rate”, the reciprocal of the average time between the events,

$$\lambda = \frac{1}{\text{average time between events}}.$$

- ▶ **R** name: `exp` with the parameter `rate = 1`

# Normal (Gaussian) Distribution

- ▶ Parameters:  $\mu$  (mean) and  $\sigma^2$  (variance);

# Normal (Gaussian) Distribution

- ▶ Parameters:  $\mu$  (mean) and  $\sigma^2$  (variance);
- ▶ Notation:  $X \sim \mathcal{N}(\mu, \sigma^2)$ ;

# Normal (Gaussian) Distribution

- ▶ Parameters:  $\mu$  (mean) and  $\sigma^2$  (variance);
- ▶ Notation:  $X \sim \mathcal{N}(\mu, \sigma^2)$ ;
- ▶ Support:  $\mathbb{R}$

# Normal (Gaussian) Distribution

- ▶ Parameters:  $\mu$  (mean) and  $\sigma^2$  (variance);
- ▶ Notation:  $X \sim \mathcal{N}(\mu, \sigma^2)$ ;
- ▶ Support:  $\mathbb{R}$
- ▶ PDF:

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

# Normal (Gaussian) Distribution

- ▶ Parameters:  $\mu$  (mean) and  $\sigma^2$  (variance);
- ▶ Notation:  $X \sim \mathcal{N}(\mu, \sigma^2)$ ;
- ▶ Support:  $\mathbb{R}$
- ▶ PDF:

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \mu$ ,  $\text{Var}(X) = \sigma^2$ .

## Normal (Gaussian) Distribution

- ▶ Parameters:  $\mu$  (mean) and  $\sigma^2$  (variance);
- ▶ Notation:  $X \sim \mathcal{N}(\mu, \sigma^2)$ ;
- ▶ Support:  $\mathbb{R}$
- ▶ PDF:

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \mu$ ,  $\text{Var}(X) = \sigma^2$ .
- ▶ Models: Different things. Important is because of the CLT



## Normal (Gaussian) Distribution

- ▶ Parameters:  $\mu$  (mean) and  $\sigma^2$  (variance);
- ▶ Notation:  $X \sim \mathcal{N}(\mu, \sigma^2)$ ;
- ▶ Support:  $\mathbb{R}$
- ▶ PDF:

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \mu$ ,  $\text{Var}(X) = \sigma^2$ .
- ▶ Models: Different things. Important is because of the CLT
- ▶ **R** name: `norm` with the parameters `mean = 0`, `sd = 1`

# Normal (Gaussian) Distribution

- ▶ Parameters:  $\mu$  (mean) and  $\sigma^2$  (variance);
- ▶ Notation:  $X \sim \mathcal{N}(\mu, \sigma^2)$ ;
- ▶ Support:  $\mathbb{R}$
- ▶ PDF:

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

- ▶ Mean and Variance:  $\mathbb{E}(X) = \mu$ ,  $\text{Var}(X) = \sigma^2$ .
- ▶ Models: Different things. Important is because of the CLT
- ▶ **R** name: `norm` with the parameters `mean = 0`, `sd = 1`
- ▶ Example:

```
rnorm(10, mean = 2, sd = 3)
```

```
## [1] -0.4924022  4.0012314  6.4835360  2.3935659  4.9664
## [7]  2.7634180  4.5808330  2.4298559 -1.0900586
```

# Normal (Gaussian) Distribution

## Note:

- ▶ **R** is using  $\mathcal{N}(\text{mean}, \text{sd})$  format
- ▶ **In Math** we are using the  $\mathcal{N}(\text{mean}, \text{variance})$  format

# Normal (Gaussian) Distribution

## Note:

- ▶ **R** is using  $\mathcal{N}(\text{mean}, \text{sd})$  format
- ▶ **In Math** we are using the  $\mathcal{N}(\text{mean}, \text{variance})$  format

So if you want to generate a sample of size 100 from  $\mathcal{N}(2, 9)$ , use the command `rnorm(100, mean = 2, sd = 3)`.

# Normal (Gaussian) Distribution

## Note:

- ▶ **R** is using  $\mathcal{N}(\text{mean}, \text{sd})$  format
- ▶ **In Math** we are using the  $\mathcal{N}(\text{mean}, \text{variance})$  format

So if you want to generate a sample of size 100 from  $\mathcal{N}(2, 9)$ , use the command `rnorm(100, mean = 2, sd = 3)`.

## Additional Properties:

- ▶ If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $Z = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1)$  and

$$\begin{aligned}\mathbb{P}(a < X < b) &= \mathbb{P}\left(\frac{a - \mu}{\sigma} < \frac{X - \mu}{\sigma} < \frac{b - \mu}{\sigma}\right) = \\ &= \Phi\left(\frac{b - \mu}{\sigma}\right) - \Phi\left(\frac{a - \mu}{\sigma}\right).\end{aligned}$$

# Normal (Gaussian) Distribution

- ▶ If  $Z \sim \mathcal{N}(0, 1)$ , then  $X = \mu + \sigma \cdot Z \sim \mathcal{N}(\mu, \sigma^2)$

# Normal (Gaussian) Distribution

- ▶ If  $Z \sim \mathcal{N}(0, 1)$ , then  $X = \mu + \sigma \cdot Z \sim \mathcal{N}(\mu, \sigma^2)$
- ▶ If  $X_k \sim \mathcal{N}(\mu_k, \sigma_k^2)$  for  $k = 1, \dots, n$  are independent, then  $X_1 + X_2 + \dots + X_n \sim \mathcal{N}(\mu_1 + \mu_2 + \dots + \mu_n, \sigma_1^2 + \sigma_2^2 + \dots + \sigma_n^2)$ .

# Normal (Gaussian) Distribution

- ▶ If  $Z \sim \mathcal{N}(0, 1)$ , then  $X = \mu + \sigma \cdot Z \sim \mathcal{N}(\mu, \sigma^2)$
- ▶ If  $X_k \sim \mathcal{N}(\mu_k, \sigma_k^2)$  for  $k = 1, \dots, n$  are independent, then  $X_1 + X_2 + \dots + X_n \sim \mathcal{N}(\mu_1 + \mu_2 + \dots + \mu_n, \sigma_1^2 + \sigma_2^2 + \dots + \sigma_n^2)$ .
- ▶ If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$\mathbb{P}(-\sigma < X - \mu < \sigma) \approx 0.6827,$$

$$\mathbb{P}(-2\sigma < X - \mu < 2\sigma) \approx 0.9545$$

and

$$\mathbb{P}(-3\sigma < X - \mu < 3\sigma) \approx 0.9973.$$



## Additions

- ▶ See many other Distributions at [Wiki](#) or in different textbooks.

# Additions

- ▶ See many other Distributions at [Wiki](#) or in different textbooks.
- ▶ Relationships among probability distributions: [Wiki](#)

# Additions

- ▶ See many other Distributions at [Wiki](#) or in different textbooks.
- ▶ Relationships among probability distributions: [Wiki](#)
- ▶ Another page for the Relationship: [L. Leemis Page](#)