

AUA CS108, Statistics, Fall 2020

Lecture 10

Michael Poghosyan

16 Sep 2020

Contents

- ▶ BoxPlot
- ▶ Sample Quantiles
- ▶ Theoretical Quantiles
- ▶ Q-Q Plots

Example (Quartiles and IQR)

Example: Find the Quartiles and IQR of

$x : 1, 1, 2, 3, 1, 1, 3, 4, 5, 2$

Quartiles and IQR

Remark: Note that the Quartiles Q_1 , Q_2 , Q_3 are not always Datapoints.

Quartiles and IQR

Remark: Note that the Quartiles Q_1, Q_2, Q_3 are not always Datapoints.

Note: Recall the idea of Quartiles: the points Q_1, Q_2, Q_3 on the real axis divide our Dataset into (almost) four equal-length portions:

- ▶ almost 25% of our Datapoints are to the left to Q_1

Quartiles and IQR

Remark: Note that the Quartiles Q_1, Q_2, Q_3 are not always Datapoints.

Note: Recall the idea of Quartiles: the points Q_1, Q_2, Q_3 on the real axis divide our Dataset into (almost) four equal-length portions:

- ▶ almost 25% of our Datapoints are to the left to Q_1
- ▶ almost 25% of our Datapoints are between Q_1 and Q_2

Quartiles and IQR

Remark: Note that the Quartiles Q_1 , Q_2 , Q_3 are not always Datapoints.

Note: Recall the idea of Quartiles: the points Q_1 , Q_2 , Q_3 on the real axis divide our Dataset into (almost) four equal-length portions:

- ▶ almost 25% of our Datapoints are to the left to Q_1
- ▶ almost 25% of our Datapoints are between Q_1 and Q_2
- ▶ almost 25% of our Datapoints are between Q_2 and Q_3

Quartiles and IQR

Remark: Note that the Quartiles Q_1 , Q_2 , Q_3 are not always Datapoints.

Note: Recall the idea of Quartiles: the points Q_1 , Q_2 , Q_3 on the real axis divide our Dataset into (almost) four equal-length portions:

- ▶ almost 25% of our Datapoints are to the left to Q_1
- ▶ almost 25% of our Datapoints are between Q_1 and Q_2
- ▶ almost 25% of our Datapoints are between Q_2 and Q_3
- ▶ almost 25% of our Datapoints are to the right to Q_3

Quartiles and IQR

Remark: Note that the Quartiles Q_1 , Q_2 , Q_3 are not always Datapoints.

Note: Recall the idea of Quartiles: the points Q_1 , Q_2 , Q_3 on the real axis divide our Dataset into (almost) four equal-length portions:

- ▶ almost 25% of our Datapoints are to the left to Q_1
- ▶ almost 25% of our Datapoints are between Q_1 and Q_2
- ▶ almost 25% of our Datapoints are between Q_2 and Q_3
- ▶ almost 25% of our Datapoints are to the right to Q_3

Note: The interval $[Q_1, Q_3]$ contains almost the half of the Datapoints.

Quartiles and IQR

Remark: Note that the Quartiles Q_1 , Q_2 , Q_3 are not always Datapoints.

Note: Recall the idea of Quartiles: the points Q_1 , Q_2 , Q_3 on the real axis divide our Dataset into (almost) four equal-length portions:

- ▶ almost 25% of our Datapoints are to the left to Q_1
- ▶ almost 25% of our Datapoints are between Q_1 and Q_2
- ▶ almost 25% of our Datapoints are between Q_2 and Q_3
- ▶ almost 25% of our Datapoints are to the right to Q_3

Note: The interval $[Q_1, Q_3]$ contains almost the half of the Datapoints. So the IQR shows the Spread of the middle half of our Dataset, it is a measure of the Spread/Variability.

Quartiles in R

In **R**, one can use the commands `quantile(x, 0.25)` and `quantile(x, 0.75)` to find Q_1 and Q_3 .

Quartiles in R

In R, one can use the commands `quantile(x, 0.25)` and `quantile(x, 0.75)` to find Q_1 and Q_3 . For example,

```
x <- 1:10  
quantile(x,0.25)
```

```
## 25%
```

```
## 3.25
```

Quartiles in R

In **R**, one can use the commands `quantile(x, 0.25)` and `quantile(x, 0.75)` to find Q_1 and Q_3 . For example,

```
x <- 1:10  
quantile(x,0.25)
```

```
## 25%  
## 3.25
```

If you will not give a parameter to `quantile`, **R** will calculate 0% (minimum datapoint), 25%, 50%, 75% and 100% (maximum datapoint) quartiles:

```
x <- 1:10  
quantile(x)
```

```
##      0%      25%      50%      75%     100%  
##  1.00   3.25   5.50   7.75  10.00
```

Quartiles in R

Also, you can use the following commands:

```
x <- 1:10  
fivenum(x)
```

```
## [1] 1.0 3.0 5.5 8.0 10.0
```

```
summary(x)
```

##	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
##	1.00	3.25	5.50	5.50	7.75	10.00

Quartiles in R

Also, you can use the following commands:

```
x <- 1:10  
fivenum(x)
```

```
## [1] 1.0 3.0 5.5 8.0 10.0
```

```
summary(x)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.  
##      1.00   3.25   5.50   5.50   7.75   10.00
```

To calculate the IQR in **R**, we can use the IQR command:

```
x <- 1:10  
IQR(x)
```

```
## [1] 4.5
```

Note

Note: Please note that **R** is not using our definition of the Quartiles, so sometimes we will get different results when calculating by a hand or by **R**.

BoxPlot

BoxPlot (or Box and Whiskers Plot) is another very common method of visualization.

BoxPlot

BoxPlot (or Box and Whiskers Plot) is another very common method of visualization. To draw the BoxPlot, we calculate the following:

BoxPlot

BoxPlot (or Box and Whiskers Plot) is another very common method of visualization. To draw the BoxPlot, we calculate the following:

- ▶ The Quartiles $Q_1, Q_2 = \textit{Median}, Q_3$

BoxPlot

BoxPlot (or Box and Whiskers Plot) is another very common method of visualization. To draw the BoxPlot, we calculate the following:

- ▶ The Quartiles $Q_1, Q_2 = \textit{Median}, Q_3$
- ▶ the Lower and Upper Fences
 $W_1 = \min\{x_i : x_i \geq Q_1 - 1.5 \cdot IQR\}$ and
 $W_2 = \max\{x_i : x_i \leq Q_3 + 1.5 \cdot IQR\},$

BoxPlot

BoxPlot (or Box and Whiskers Plot) is another very common method of visualization. To draw the BoxPlot, we calculate the following:

- ▶ The Quartiles $Q_1, Q_2 = \text{Median}, Q_3$
- ▶ the Lower and Upper Fences
 $W_1 = \min\{x_i : x_i \geq Q_1 - 1.5 \cdot IQR\}$ and
 $W_2 = \max\{x_i : x_i \leq Q_3 + 1.5 \cdot IQR\}$, i.e., the first and last observations lying in

$$\left[Q_1 - \frac{3}{2}IQR, Q_3 + \frac{3}{2}IQR \right];$$

BoxPlot

BoxPlot (or Box and Whiskers Plot) is another very common method of visualization. To draw the BoxPlot, we calculate the following:

- ▶ The Quartiles $Q_1, Q_2 = \textit{Median}, Q_3$
- ▶ the Lower and Upper Fences
 $W_1 = \min\{x_i : x_i \geq Q_1 - 1.5 \cdot IQR\}$ and
 $W_2 = \max\{x_i : x_i \leq Q_3 + 1.5 \cdot IQR\}$, i.e., the first and last observations lying in

$$\left[Q_1 - \frac{3}{2}IQR, Q_3 + \frac{3}{2}IQR \right];$$

the lines joining that fences to corresponding quartiles are the *Whiskers*;

BoxPlot

BoxPlot (or Box and Whiskers Plot) is another very common method of visualization. To draw the BoxPlot, we calculate the following:

- ▶ The Quartiles $Q_1, Q_2 = \text{Median}, Q_3$
- ▶ the Lower and Upper Fences
 $W_1 = \min\{x_i : x_i \geq Q_1 - 1.5 \cdot IQR\}$ and
 $W_2 = \max\{x_i : x_i \leq Q_3 + 1.5 \cdot IQR\}$, i.e., the first and last observations lying in

$$\left[Q_1 - \frac{3}{2}IQR, Q_3 + \frac{3}{2}IQR \right];$$

the lines joining that fences to corresponding quartiles are the *Whiskers*;

- ▶ the set of all Outliers

$$O = \left\{ x_i : x_i \notin \left[Q_1 - \frac{3}{2}IQR, Q_3 + \frac{3}{2}IQR \right] \right\}$$

BoxPlot, Example

Then we draw the points W_1, Q_1, Q_2, Q_3, W_2 on the real line and add all outliers, and make a box over $[Q_1, Q_3]$.

BoxPlot, Example

Then we draw the points W_1, Q_1, Q_2, Q_3, W_2 on the real line and add all outliers, and make a box over $[Q_1, Q_3]$.

Example: Draw the Boxplot of

$$x : 0, -2, 2, 1, 5, 6, 4, 1, 2, 1, 12$$

BoxPlot, Example

Then we draw the points W_1, Q_1, Q_2, Q_3, W_2 on the real line and add all outliers, and make a box over $[Q_1, Q_3]$.

Example: Draw the Boxplot of


$$x : 0, -2, 2, 1, 5, 6, 4, 1, 2, 1, 12$$

Solution: OTB

BoxPlot, Example

Now, using **R**:


```
x <- c(0, -2, 2, 1, 5, 6, 4, 1, 2, 1, 12)
boxplot(x)
```



BoxPlot, Example

Another view:



```
x <- c(0, -2, 2, 1, 5, 6, 4, 1, 2, 1, 12)
boxplot(x, horizontal = T, col = "magenta")
```



BoxPlot, Example

Here are some Datasets' Histograms along with the BoxPlots:



```
x <- rnorm(1000, mean = -3, sd = 2)
par(mfrow=c(2,1)); hist(x)
boxplot(x, horizontal = T, col = "cyan")
```



BoxPlot, Example

Here are some Datasets' Histograms along with the BoxPlots:



```
x <- rexp(1000, rate = 0.2)
par(mfrow=c(2,1)); hist(x)
boxplot(x, horizontal = T, col = "lightgreen")
```



BoxPlot, Example


Here are some Datasets' Histograms along with the BoxPlots:

```
x <- rbeta(1000, shape1 = 4, shape2 = 2)
par(mfrow=c(2,1)); hist(x)
boxplot(x, horizontal = T, col = "red")
```





BoxPlot, Example

Here is the BoxPlot of our Quiz 01 grades: can you describe the result?



BoxPlot, Example

And here is the BoxPlot of our Quiz 01 grades along with the Histogram:



##	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
##	7.50	61.25	82.50	74.63	95.00	100.00