

# AUA CS108, Statistics, Fall 2020

## Lecture 28

Michael Poghosyan

30 Oct 2020

# Contents

- ▶ Properties of Estimators: MSE
- ▶ Bias and Unbiasedness

# Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator  $\hat{\theta}$  to estimate  $\theta$ .

# Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator  $\hat{\theta}$  to estimate  $\theta$ .

**Definition:** The **Mean Squared Error** or the **Quadratic Risk** of the estimator  $\hat{\theta}_n$  of  $\theta$  is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

# Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator  $\hat{\theta}$  to estimate  $\theta$ .

**Definition:** The **Mean Squared Error** or the **Quadratic Risk** of the estimator  $\hat{\theta}_n$  of  $\theta$  is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

**Note:** *MSE* calculates how close are, in the Quadratic Mean sense, possible values of the Estimator  $\hat{\theta}$  to the actual (unknown) value of  $\theta$ .

# Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator  $\hat{\theta}$  to estimate  $\theta$ .

**Definition:** The **Mean Squared Error** or the **Quadratic Risk** of the estimator  $\hat{\theta}_n$  of  $\theta$  is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

**Note:** *MSE* calculates how close are, in the Quadratic Mean sense, possible values of the Estimator  $\hat{\theta}$  to the actual (unknown) value of  $\theta$ . The smaller the value of *MSE*, the better, of course.

# Risk, Mean Squared Error of the Estimator

Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta,$$

and we use the Estimator  $\hat{\theta}$  to estimate  $\theta$ .

**Definition:** The **Mean Squared Error** or the **Quadratic Risk** of the estimator  $\hat{\theta}_n$  of  $\theta$  is

$$MSE(\hat{\theta}_n, \theta) = Risk(\hat{\theta}_n, \theta) = \mathbb{E}_\theta[(\hat{\theta}_n - \theta)^2].$$

**Note:** *MSE* calculates how close are, in the Quadratic Mean sense, possible values of the Estimator  $\hat{\theta}$  to the actual (unknown) value of  $\theta$ . The smaller the value of *MSE*, the better, of course.

**Definition:** We say that the estimator  $\hat{\theta}_n^1$  of  $\theta$  is **preferable** to  $\hat{\theta}_n^2$ , another estimator of  $\theta$ , if

$$MSE(\hat{\theta}_n^1, \theta) \leq MSE(\hat{\theta}_n^2, \theta), \quad \forall \theta \in \Theta,$$

and there exists a  $\theta$  s.t.  $MSE(\hat{\theta}_n^1, \theta) < MSE(\hat{\theta}_n^2, \theta)$ .

## Useful Identity

A very useful identity for calculating  $\mathbb{E}(X^2)$  for a r.v.  $X$  is the following:



## Useful Identity

A very useful identity for calculating  $\mathbb{E}(X^2)$  for a r.v.  $X$  is the following: because

$$\text{Var}(X) = \mathbb{E}(X^2) - [\mathbb{E}(X)]^2,$$

then

## Useful Identity

A very useful identity for calculating  $\mathbb{E}(X^2)$  for a r.v.  $X$  is the following: because

$$\text{Var}(X) = \mathbb{E}(X^2) - \left[\mathbb{E}(X)\right]^2,$$

then

$$\mathbb{E}(X^2) = \text{Var}(X) + \left[\mathbb{E}(X)\right]^2.$$

## Example

**Example:** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda),$$

and we want to Estimate the unknown  $\lambda$ .

## Example

**Example:** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda),$$

and we want to Estimate the unknown  $\lambda$ . We want to use one of the following Estimators:

$$\hat{\lambda}_1 = X_1 \quad \text{or} \quad \hat{\lambda}_2 = \frac{X_1 + X_n}{2}.$$

## Example

**Example:** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda),$$

and we want to Estimate the unknown  $\lambda$ . We want to use one of the following Estimators:

$$\hat{\lambda}_1 = X_1 \quad \text{or} \quad \hat{\lambda}_2 = \frac{X_1 + X_n}{2}.$$

**Question:** Calculate the Risks of  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$ . Compare these two Estimators: which of  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$  is preferable?

## Example

**Example:** Assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Pois}(\lambda),$$

and we want to Estimate the unknown  $\lambda$ . We want to use one of the following Estimators:

$$\hat{\lambda}_1 = X_1 \quad \text{or} \quad \hat{\lambda}_2 = \frac{X_1 + X_n}{2}.$$

**Question:** Calculate the Risks of  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$ . Compare these two Estimators: which of  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$  is preferable?

**Solution:** OTB: Use the fact that for  $X \sim \text{Pois}(\lambda)$ ,

$$\mathbb{E}(X) = \lambda = \text{Var}(X).$$

## Best MSE Estimators

Now, having the idea of the Error of estimation for an Estimator, we can try to find the best Estimator in the sense of MSE, i.e., we can try to solve

Find  $\hat{\theta}$  among all Estimators that minimizes  $MSE(\hat{\theta}, \theta)$ .

## Best MSE Estimators

Now, having the idea of the Error of estimation for an Estimator, we can try to find the best Estimator in the sense of MSE, i.e., we can try to solve

Find  $\hat{\theta}$  among all Estimators that minimizes  $MSE(\hat{\theta}, \theta)$ .

Unfortunately, this Minimization problem has no solution in general.



## Best MSE Estimators

Now, having the idea of the Error of estimation for an Estimator, we can try to find the best Estimator in the sense of MSE, i.e., we can try to solve

Find  $\hat{\theta}$  among all Estimators that minimizes  $MSE(\hat{\theta}, \theta)$ .

Unfortunately, this Minimization problem has no solution in general.

So we move forward considering other Properties of Estimators making them useful in the estimation process.

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

**Definition:** We say that our Estimator  $\hat{\theta}$  (of the parameter  $\theta$ ) is **Unbiased**, if

$$\text{Bias}(\hat{\theta}, \theta) = 0 \quad \text{for any} \quad \theta \in \Theta.$$

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

**Definition:** We say that our Estimator  $\hat{\theta}$  (of the parameter  $\theta$ ) is **Unbiased**, if

$$\text{Bias}(\hat{\theta}, \theta) = 0 \quad \text{for any} \quad \theta \in \Theta.$$

If  $\hat{\theta}$  is not Unbiased, then we say that it is **Biased**.

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

**Definition:** We say that our Estimator  $\hat{\theta}$  (of the parameter  $\theta$ ) is **Unbiased**, if

$$\text{Bias}(\hat{\theta}, \theta) = 0 \quad \text{for any} \quad \theta \in \Theta.$$

If  $\hat{\theta}$  is not Unbiased, then we say that it is **Biased**.

In other words, the Estimator  $\hat{\theta}$  is **Unbiased**, if

$$\mathbb{E}_{\theta}(\hat{\theta}) = \theta \quad \text{for any} \quad \theta \in \Theta.$$

## Bias, Biased and Unbiased Estimators

**Definition:** The **Bias** of Estimator  $\hat{\theta}$  of  $\theta$  is

$$\text{Bias}(\hat{\theta}, \theta) = \mathbb{E}_{\theta}(\hat{\theta} - \theta) = \mathbb{E}_{\theta}(\hat{\theta}) - \theta.$$

**Definition:** We say that our Estimator  $\hat{\theta}$  (of the parameter  $\theta$ ) is **Unbiased**, if

$$\text{Bias}(\hat{\theta}, \theta) = 0 \quad \text{for any} \quad \theta \in \Theta.$$

If  $\hat{\theta}$  is not Unbiased, then we say that it is **Biased**.

In other words, the Estimator  $\hat{\theta}$  is **Unbiased**, if

$$\mathbb{E}_{\theta}(\hat{\theta}) = \theta \quad \text{for any} \quad \theta \in \Theta.$$

**Note:** We require the above hold **for any parameter value** since, if, say, it is correct for *some* values of  $\theta$ , then it can happen that the true value of our unknown  $\theta$  is exactly that value, for which we do not have the equality.

# Bias and Unbiasedness

- ▶ The idea of Unbiased Estimator is the following: if we will calculate Estimates many-many times using our Unbiased Estimator, and then average the obtained Estimates, we will obtain the (almost exact, exact when many-many  $\rightarrow \infty$ ) value of our Parameter.

# Bias and Unbiasedness

- ▶ The idea of Unbiased Estimator is the following: if we will calculate Estimates many-many times using our Unbiased Estimator, and then average the obtained Estimates, we will obtain the (almost exact, exact when many-many  $\rightarrow \infty$ ) value of our Parameter.
- ▶ So we can say about an Unbiased Estimator as: **In average, it is Exact**



# Bias and Unbiasedness

- ▶ The idea of Unbiased Estimator is the following: if we will calculate Estimates many-many times using our Unbiased Estimator, and then average the obtained Estimates, we will obtain the (almost exact, exact when many-many  $\rightarrow \infty$ ) value of our Parameter.
- ▶ So we can say about an Unbiased Estimator as: **In average, it is Exact**
- ▶ Bias can be interpreted, in some sense, as the *accuracy* of the Estimator.

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population.

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution.

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution. So our unknown Parameter here is  $\mu$ .

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution. So our unknown Parameter here is  $\mu$ .

We use size  $n$  Random Sample to Model the situation:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\mu.$$

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution. So our unknown Parameter here is  $\mu$ .

We use size  $n$  Random Sample to Model the situation:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\mu.$$

Now, we choose several Estimators to estimate  $\mu$ :

$$\begin{aligned}\hat{\mu}_1 &= X_1, & \hat{\mu}_2 &= \frac{X_1 + X_3}{2}, & \hat{\mu}_3 &= \frac{X_1 + X_4}{10}, \\ \hat{\mu}_4 &= \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.\end{aligned}$$

## Example

**Example:** Assume we have a Sample of size  $n$  from some Population, and we want to estimate the Mean of that Population. Let  $\mathcal{F}_\mu$  be the Distribution of our Population, and  $\mu$  be the Mean (Expectation) of that Distribution. So our unknown Parameter here is  $\mu$ .

We use size  $n$  Random Sample to Model the situation:

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\mu.$$

Now, we choose several Estimators to estimate  $\mu$ :

$$\hat{\mu}_1 = X_1, \quad \hat{\mu}_2 = \frac{X_1 + X_3}{2}, \quad \hat{\mu}_3 = \frac{X_1 + X_4}{10},$$
$$\hat{\mu}_4 = \bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

**Problem:** Check if each Estimator is Biased or Unbiased.

**Solution:** OTB

## Example

**Example:** Let's do an experiment with Biased and Unbiased Estimators, with **R**.



## Example

**Example:** Let's do an experiment with Biased and Unbiased Estimators, with **R**.

### UnBiased Estimator Case

We consider the Poisson Model:

$$X_1, X_2, \dots, X_{10} \sim \text{Pois}(\lambda)$$

and we want to estimate  $\lambda$ .

## Example

**Example:** Let's do an experiment with Biased and Unbiased Estimators, with **R**.

### UnBiased Estimator Case

We consider the Poisson Model:

$$X_1, X_2, \dots, X_{10} \sim \text{Pois}(\lambda)$$

and we want to estimate  $\lambda$ . We consider the following Estimator:

$$\hat{\lambda} = \frac{X_1 + X_2 + \dots + X_{10}}{10}.$$

## Example

**Example:** Let's do an experiment with Biased and Unbiased Estimators, with **R**.

### UnBiased Estimator Case

We consider the Poisson Model:

$$X_1, X_2, \dots, X_{10} \sim \text{Pois}(\lambda)$$

and we want to estimate  $\lambda$ . We consider the following Estimator:

$$\hat{\lambda} = \frac{X_1 + X_2 + \dots + X_{10}}{10}.$$

Easy to see that  $\hat{\lambda}$  is an Unbiased Estimator for  $\lambda$  (OTB!).

## Example, cont'd

Now, the code


- ▶ observing once: generating a Sample just once and calculating one Estimate:

```
lambda <- 3.21  
x <- rpois(10, lambda = lambda)  
lambda.hat <- mean(x)  
lambda.hat
```

```
## [1] 3
```

- ▶ observing many times: generating Samples many times, calculating Estimates, and then averaging:

```
lambda <- 3.21; n <- 10; m <- 2000  
x <- rpois(n*m, lambda = lambda)  
x <- as.data.frame(matrix(x, ncol = m))  
lambda.hats <- sapply(x, mean)  
boxplot(lambda.hats, horizontal = T);  
abline(v = lambda, col="red", lwd = 2, lty = 2)
```



```
mean(lambda.hats)
```

```
## [1] 3.22645
```

With a Histogram:

```
hist(lambda.hats)
rug(lambda.hats)
abline(v = lambda, col="red", lwd = 2, lty = 2)
```

