

CS 107, Probability, Spring 2019

Lecture 27

Michael Poghosyan

AUA

27 March 2019

- Examples of Important Continuous R.V.s

The Galton Board

See <https://www.youtube.com/watch?v=6YDHBfVivIs>

The Galton Board

See <https://www.youtube.com/watch?v=6YDHBfVivIs>
and
<https://www.youtube.com/watch?v=UCmPmkHqHXk>

The Galton Board

See <https://www.youtube.com/watch?v=6YDHBfVIvIs>
and
<https://www.youtube.com/watch?v=UCmPmkHqHXk>
and, maybe, <https://galtonboard.com/>

Exponential Distribution

Exponential Distribution

We say that the r.v. X has an Exponential Distribution with the parameter $\lambda > 0$ (rate), and we write $X \sim \text{Exp}(\lambda)$, if its PDF is given by

$$f(x) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0, \\ 0, & x < 0 \end{cases}$$

Exponential Distribution

Exponential Distribution

We say that the r.v. X has an Exponential Distribution with the parameter $\lambda > 0$ (rate), and we write $X \sim \text{Exp}(\lambda)$, if its PDF is given by

$$f(x) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0, \\ 0, & x < 0 \end{cases}$$

Exercise: Check that this function is a PDF of some r.v.

Exponential Distribution

Exponential Distribution

We say that the r.v. X has an Exponential Distribution with the parameter $\lambda > 0$ (rate), and we write $X \sim \text{Exp}(\lambda)$, if its PDF is given by

$$f(x) = \begin{cases} \lambda \cdot e^{-\lambda x}, & x \geq 0, \\ 0, & x < 0 \end{cases}$$

Exercise: Check that this function is a PDF of some r.v.

Exercise: Prove that the CDF of $X \sim \text{Exp}(\lambda)$ is given by:

$$F(x) = \begin{cases} 1 - e^{-\lambda x}, & x \geq 0, \\ 0, & x < 0 \end{cases}$$

Exponential Distribution Examples

We use the Exponential Distribution to model the waiting times for some events to happen, the time until the next event will happen.

Exponential Distribution Examples

We use the Exponential Distribution to model the waiting times for some events to happen, the time until the next event will happen.

- Say, the waiting time X until the next phone call at the GG taxi service during the time interval 9AM and 10AM can be modeled as an Exponential r.v., $X \sim \text{Exp}(\lambda)$.

Exponential Distribution Examples

We use the Exponential Distribution to model the waiting times for some events to happen, the time until the next event will happen.

- Say, the waiting time X until the next phone call at the GG taxi service during the time interval 9AM and 10AM can be modeled as an Exponential r.v., $X \sim \text{Exp}(\lambda)$.
- Say, the time between clicks on the webpage in some fixed time interval can be modeled as an Exponential r.v.;

Exponential Distribution Examples

We use the Exponential Distribution to model the waiting times for some events to happen, the time until the next event will happen.

- Say, the waiting time X until the next phone call at the GG taxi service during the time interval 9AM and 10AM can be modeled as an Exponential r.v., $X \sim \text{Exp}(\lambda)$.
- Say, the time between clicks on the webpage in some fixed time interval can be modeled as an Exponential r.v.;
- The time between goals scored in a World Cup soccer match;

Exponential Distribution Examples

We use the Exponential Distribution to model the waiting times for some events to happen, the time until the next event will happen.

- Say, the waiting time X until the next phone call at the GG taxi service during the time interval 9AM and 10AM can be modeled as an Exponential r.v., $X \sim \text{Exp}(\lambda)$.
- Say, the time between clicks on the webpage in some fixed time interval can be modeled as an Exponential r.v.;
- The time between goals scored in a World Cup soccer match; Time between meteors greater than 1 meter diameter striking earth;

Exponential Distribution Examples

We use the Exponential Distribution to model the waiting times for some events to happen, the time until the next event will happen.

- Say, the waiting time X until the next phone call at the GG taxi service during the time interval 9AM and 10AM can be modeled as an Exponential r.v., $X \sim \text{Exp}(\lambda)$.
- Say, the time between clicks on the webpage in some fixed time interval can be modeled as an Exponential r.v.;
- The time between goals scored in a World Cup soccer match; Time between meteors greater than 1 meter diameter striking earth; Time between successive failures of a machine;

Exponential Distribution Examples

We use the Exponential Distribution to model the waiting times for some events to happen, the time until the next event will happen.

- Say, the waiting time X until the next phone call at the GG taxi service during the time interval 9AM and 10AM can be modeled as an Exponential r.v., $X \sim \text{Exp}(\lambda)$.
- Say, the time between clicks on the webpage in some fixed time interval can be modeled as an Exponential r.v.;
- The time between goals scored in a World Cup soccer match; Time between meteors greater than 1 meter diameter striking earth; Time between successive failures of a machine; Time until the discovery of a new Bitcoin block,

Exponential Distribution Examples

We use the Exponential Distribution to model the waiting times for some events to happen, the time until the next event will happen.

- Say, the waiting time X until the next phone call at the GG taxi service during the time interval 9AM and 10AM can be modeled as an Exponential r.v., $X \sim \text{Exp}(\lambda)$.
- Say, the time between clicks on the webpage in some fixed time interval can be modeled as an Exponential r.v.;
- The time between goals scored in a World Cup soccer match; Time between meteors greater than 1 meter diameter striking earth; Time between successive failures of a machine; Time until the discovery of a new Bitcoin block, Time (over the short term) a storekeeper must wait before the arrival of their next customer;

Exponential Distribution Examples

We use the Exponential Distribution to model the waiting times for some events to happen, the time until the next event will happen.

- Say, the waiting time X until the next phone call at the GG taxi service during the time interval 9AM and 10AM can be modeled as an Exponential r.v., $X \sim \text{Exp}(\lambda)$.
- Say, the time between clicks on the webpage in some fixed time interval can be modeled as an Exponential r.v.;
- The time between goals scored in a World Cup soccer match; Time between meteors greater than 1 meter diameter striking earth; Time between successive failures of a machine; Time until the discovery of a new Bitcoin block, Time (over the short term) a storekeeper must wait before the arrival of their next customer; Time between late arrivals to Prob class ...

The Rate Parameter; Exponential and Poisson Distributions

The Rate Parameter; Exponential and Poisson Distributions

Note: The rate parameter λ is the inverse of the average waiting time. That is, if the average time between two phone calls is $2.5min$, then we can model the waiting time as $X \sim Exp(1/2.5)$.

The Rate Parameter; Exponential and Poisson Distributions

Note: The rate parameter λ is the inverse of the average waiting time. That is, if the average time between two phone calls is $2.5min$, then we can model the waiting time as $X \sim Exp(1/2.5)$.

Difference between the Poisson and Exponential Distribution is the following:

The Rate Parameter; Exponential and Poisson Distributions

Note: The rate parameter λ is the inverse of the average waiting time. That is, if the average time between two phone calls is $2.5min$, then we can model the waiting time as $X \sim Exp(1/2.5)$.

Difference between the Poisson and Exponential Distribution is the following:

Exponential Distribution is calculating the **time between two events**, and the **time until the first event will happen**, which is continuous, and Poisson Distribution is calculating the **Number of events (occurrences)**, which is discrete.

Exponential Distributions: Example

Example: Assume $X \sim \text{Exp}(2.6)$. Calculate $\mathbb{P}(X = 2.6)$, $\mathbb{P}(X \leq 2)$, $\mathbb{P}(2 \leq X \leq 3.5)$, $\mathbb{P}(X > 3|X > 1)$.

Memoryless Property

The following property is very important characteristic of the Exponential Distribution: this distribution is the only continuous distribution with the Memoryless Property:

Memoryless Property

The following property is very important characteristic of the Exponential Distribution: this distribution is the only continuous distribution with the Memoryless Property: $X \sim \text{Exp}(\lambda)$ iff

$$\mathbb{P}(X > t + s | X > t) = \mathbb{P}(X > s), \quad \forall t, s \in [0, +\infty).$$

Memoryless Property

The following property is very important characteristic of the Exponential Distribution: this distribution is the only continuous distribution with the Memoryless Property: $X \sim \text{Exp}(\lambda)$ iff

$$\mathbb{P}(X > t + s | X > t) = \mathbb{P}(X > s), \quad \forall t, s \in [0, +\infty).$$

Interpretation: If X is the waiting time for smthng, then $\mathbb{P}(X > s)$ is the Probability that the waiting time will be more than s .

Memoryless Property

The following property is very important characteristic of the Exponential Distribution: this distribution is the only continuous distribution with the Memoryless Property: $X \sim \text{Exp}(\lambda)$ iff

$$\mathbb{P}(X > t + s | X > t) = \mathbb{P}(X > s), \quad \forall t, s \in [0, +\infty).$$

Interpretation: If X is the waiting time for smthng, then $\mathbb{P}(X > s)$ is the Probability that the waiting time will be more than s . Now, $\mathbb{P}(X > t + s | X > t)$ is the Probability that the waiting time will be more than $t + s$, if you have waited already t units of time.

Memoryless Property

The following property is very important characteristic of the Exponential Distribution: this distribution is the only continuous distribution with the Memoryless Property: $X \sim \text{Exp}(\lambda)$ iff

$$\mathbb{P}(X > t + s | X > t) = \mathbb{P}(X > s), \quad \forall t, s \in [0, +\infty).$$

Interpretation: If X is the waiting time for smthng, then $\mathbb{P}(X > s)$ is the Probability that the waiting time will be more than s . Now, $\mathbb{P}(X > t + s | X > t)$ is the Probability that the waiting time will be more than $t + s$, if you have waited already t units of time. That is, waiting for *another* s units is independent how long you have already waited!

Memoryless Property

Analogously, the only Discrete Distribution sharing the Memoryless Property:

$$\mathbb{P}(X > m + n | X > m) = \mathbb{P}(X > n), \quad \forall m, n \in \{0, 1, 2, \dots\}$$

is the

Memoryless Property

Analogously, the only Discrete Distribution sharing the Memoryless Property:

$$\mathbb{P}(X > m + n | X > m) = \mathbb{P}(X > n), \quad \forall m, n \in \{0, 1, 2, \dots\}$$

is the Geometric Distribution (which can be described as the waiting time for the Bernoulli process).

Exponential Distribution: R Examples

R Code

```
### Exponential Distribution
lambda <- 2.4
x <- seq(from = -1, to = 5, by = 0.01)
y <- dexp(x, rate = lambda)
#plot the PDF
plot(x,y, type = "l", lwd = 3, main = "Exponential Distribution")

z <- pexp(x, rate = lambda)
#plot the CDF
plot(x,z, type = "l", lwd = 3, main = "Exponential Distribution")

##Comparing of 2 exponential PDFs
curve(dexp(x,rate = 1), -1, 5, ylim = c(0,2))
par(new = T)
curve(dexp(x,rate = 2), -1, 5, ylim = c(0,2), col = "red")
```

R Example: Moodle Page Visits Model




Figure: The histogram is for the time between our Probability Moodle visits by students in hours. Fitted red line is the Exponential Distribution with the $\lambda = \frac{1}{\text{average time between visits}}$

Normal Distribution

Normal Distribution is one of the most important distributions in Probability, in the World and in the Universe 😊

Normal Distribution

Normal Distribution is one of the most important distributions in Probability, in the World and in the Universe ☺

We start by the definition:

Normal Distribution

Normal Distribution is one of the most important distributions in Probability, in the World and in the Universe ☺

We start by the definition:

The Normal (Gaussian) Distribution

We say that the r.v. X has a Normal (or Gaussian) Distribution with the Mean μ and the Standard Deviation σ (or, with the Mean μ and Variance σ^2), and we will write

$$X \sim \mathcal{N}(\mu, \sigma^2),$$

Normal Distribution

Normal Distribution is one of the most important distributions in Probability, in the World and in the Universe ☺

We start by the definition:

The Normal (Gaussian) Distribution

We say that the r.v. X has a Normal (or Gaussian) Distribution with the Mean μ and the Standard Deviation σ (or, with the Mean μ and Variance σ^2), and we will write

$$X \sim \mathcal{N}(\mu, \sigma^2),$$

if its PDF is given by

$$f(x) = \frac{1}{\sigma \cdot \sqrt{2\pi}} \cdot e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}.$$

Standard Normal Distribution

For example, if X is a Normal r.v. with the Mean 0 and Variance 1, i.e.

$$X \sim \mathcal{N}(0, 1),$$

then we call X to be a **Standard Normal r.v.**

Standard Normal Distribution

For example, if X is a Normal r.v. with the Mean 0 and Variance 1, i.e.

$$X \sim \mathcal{N}(0, 1),$$

then we call X to be a **Standard Normal r.v.**

The PDF of the Standard Normal r.v. is given by

$$\varphi(x) = \frac{1}{\sqrt{2\pi}} \cdot e^{-\frac{x^2}{2}}, \quad x \in \mathbb{R}.$$

Standard Normal Distribution

For example, if X is a Normal r.v. with the Mean 0 and Variance 1, i.e.

$$X \sim \mathcal{N}(0, 1),$$

then we call X to be a **Standard Normal r.v.**

The PDF of the Standard Normal r.v. is given by

$$\varphi(x) = \frac{1}{\sqrt{2\pi}} \cdot e^{-\frac{x^2}{2}}, \quad x \in \mathbb{R}.$$

The CDF of the Standard Normal r.v. is given by

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \cdot \int_{-\infty}^x e^{-\frac{t^2}{2}} dt, \quad x \in \mathbb{R}.$$

Standard Normal Distribution

For example, if X is a Normal r.v. with the Mean 0 and Variance 1, i.e.

$$X \sim \mathcal{N}(0, 1),$$

then we call X to be a **Standard Normal r.v.**

The PDF of the Standard Normal r.v. is given by

$$\varphi(x) = \frac{1}{\sqrt{2\pi}} \cdot e^{-\frac{x^2}{2}}, \quad x \in \mathbb{R}.$$

The CDF of the Standard Normal r.v. is given by

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \cdot \int_{-\infty}^x e^{-\frac{t^2}{2}} dt, \quad x \in \mathbb{R}.$$

Exercise: Prove that $\varphi(x)$ is a PDF of some r.v.

Standard Normal Distribution

In 1997, the Nobel Prize in Economic Sciences was awarded to Robert Merton and Myron Scholes for the **Black-Scholes options pricing model**.

Standard Normal Distribution

In 1997, the Nobel Prize in Economic Sciences was awarded to Robert Merton and Myron Scholes for the **Black-Scholes options pricing model**. And the Black-Scholes formula involves the CDF of the Standard Normal Distribution: see, e.g., https://en.wikipedia.org/wiki/Black-Scholes_model.

Normal Distribution: R Examples

```
#Standard Normal Distribution
t <- seq(from=-4,to=4, by=0.01)
y <- dnorm(t, mean = 0, sd = 1) #PDF
plot(t,y, type = "l", lwd = 2, main = "PDF of the Standard
z <- pnorm(t, mean = 0, sd = 1) #CDF
plot(t,z, type = "l", lwd = 2, main = "CDF of the Standard
#Other Mean and SD Normal PDFs
t <- seq(from=-2,to=6, by=0.01)
y <- dnorm(t, mean = 2, sd = 1)
plot(t,y, type = "l", lwd = 2, ylim = c(0,1), main = "PDF of
u <- dnorm(t, mean = 2, sd = 0.5)
par(new = T)
plot(t,u, type = "l", lwd = 2, col = "red", ylim = c(0,1),
w <- dnorm(t, mean = 2, sd = 1.5)
par(new = T)
plot(t,w, type = "l", lwd = 2, col = "green", ylim = c(0,1),
```

Examples of Normal Distribution

You can find many examples of Normal Distribution in the real world.



Figure: Hallgrímskirkja, church in Reykjavík, Iceland

Other examples of Normal Distribution

Other examples of Normal Distribution



Figure: Normal Table

Other examples of Normal Distribution

Other examples of Normal Distribution



Other examples of Normal Distribution

Other examples of Normal Distribution



Other examples of Normal Distribution

Surely you know what us **curving**.

Other examples of Normal Distribution

Surely you know what us **curving**. And, of course, you know that the Probability that we will do curving for Probability is 0
😊.

Other examples of Normal Distribution

Surely you know what us **curving**. And, of course, you know that the Probability that we will do curving for Probability is 0 😊.

But, what is curving, in fact? And what is the **curve** we are talking about?

Other examples of Normal Distribution

Surely you know what us **curving**. And, of course, you know that the Probability that we will do curving for Probability is 0 😊.

But, what is curving, in fact? And what is the **curve** we are talking about? Of course, our new friend the Bell-Shaped Gaussian Curve!

See <https://www.youtube.com/watch?v=vqNExEhXHvc>

Now, Serious Things

Many phenomena can be modeled by the Normal Distribution.
See, for example,

<https://galtonboard.com/probabilityexamplesinlife>

Now, Serious Things

Many phenomena can be modeled by the Normal Distribution.
See, for example,

<https://galtonboard.com/probabilityexamplesinlife>

The importance of the Normal Distribution is justified by the Central Limit Theorem, which we will cover close to the end of our course, and you will use a lot in Statistics.

Now, Some Examples

Example: Assume $X \sim \mathcal{N}(0, 1)$. Calculate $\mathbb{P}(X \in \{0, 1, 2\})$, $\mathbb{P}(-1 < X < 1)$, $\mathbb{P}(X < 0)$, $\mathbb{P}(X > 3)$