# CS 107, Probability, Spring 2019 Lecture 02

Michael Poghosyan

AUA

18 January 2019

#### Content

- Experiment, Outcomes and the Sample Space
- Events, Operations with Events

#### LZ

Assume our mobile phone Weather App says that there is a 50% chance of snow for this Saturday, and also 50% chance of snow this Sunday.

Is it true that it will snow for sure (i.e., with probability 1) this weekend?

• A random (or probabilistic) Experiment is a situation, where we are uncertain about the result.

- A random (or probabilistic) Experiment is a situation, where we are uncertain about the result.
- An **Outcome** is a possible result of an Experiment.

- A random (or probabilistic) Experiment is a situation, where we are uncertain about the result.
- An **Outcome** is a possible result of an Experiment.

So if our "Experiment" has just one Outcome, then it is not a random Experiment, it is not an Experiment in our sense.

- A random (or probabilistic) Experiment is a situation, where we are uncertain about the result.
- An Outcome is a possible result of an Experiment.

So if our "Experiment" has just one Outcome, then it is not a random Experiment, it is not an Experiment in our sense.

 The set of all Outcomes of an Experiment is called the Sample Space of that Experiment:

```
\Omega = the Sample Space of the Experiment =
```

= the set of all outcomes of our Experiment

• Our Experiment: we are tossing a (fair) coin.

- Our Experiment: we are tossing a (fair) coin.
- **Heads** is one of the outcomes.

- Our Experiment: we are tossing a (fair) coin.
- **Heads** is one of the outcomes.
- The Sample Space in this Example is:

$$\Omega = \mathsf{Sample} \; \mathsf{Space} = \{\mathsf{Heads}, \; \mathsf{Tails}\}$$

- Our Experiment: we are tossing a (fair) coin.
- Heads is one of the outcomes.
- The Sample Space in this Example is:

$$\Omega = \mathsf{Sample} \; \mathsf{Space} = \{\mathsf{Heads}, \; \mathsf{Tails}\} = \{\mathsf{H}, \; \mathsf{T}\}$$

• Experiment: we are rolling a (fair) die.

- Experiment: we are rolling a (fair) die.
- One of the outcomes is •.

- Experiment: we are rolling a (fair) die.
- One of the outcomes is
- The Sample Space in this Example is:

 Experiment: we are watching (interested) if the produced detail is defective.

- Experiment: we are watching (interested) if the produced detail is defective.
- One of the outcomes is that it is Not Defective.

- Experiment: we are watching (interested) if the produced detail is defective.
- One of the outcomes is that it is Not Defective.
- The Sample Space in this Example is:

• Experiment: we are interested in the remaining lifetime (in years) of a person (for insurance reasons, say).

- Experiment: we are interested in the remaining lifetime (in years) of a person (for insurance reasons, say).
- One of the outcomes is 30.1.

- Experiment: we are interested in the remaining lifetime (in years) of a person (for insurance reasons, say).
- One of the outcomes is 30.1.
- The Sample Space in this Example is:

 Experiment: we are watching for the Google Stock closing Price for the next Friday.

- Experiment: we are watching for the Google Stock closing Price for the next Friday.
- One of the outcomes is 1092.31\$.

- Experiment: we are watching for the Google Stock closing Price for the next Friday.
- One of the outcomes is 1092.31\$.
- The Sample Space in this Example is:

• Experiment: we are rolling 2 dice.

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

#### Case 1: Order is important. Example:

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

**Case 1: Order is important**. **Example:** 2 persons are rolling dice - wins the person obtaining the largest number.

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

**Case 1: Order is important**. **Example:** 2 persons are rolling dice - wins the person obtaining the largest number.

• Some outcomes are (1,1), (2,3), (3,2)

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

**Case 1: Order is important**. **Example:** 2 persons are rolling dice - wins the person obtaining the largest number.

- Some outcomes are (1,1),(2,3),(3,2)
- Say, first person rolls a black die, the second one white:

$$(\bullet, \bullet), (\bullet, \bullet), (\bullet, \bullet)$$

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

**Case 1: Order is important**. **Example:** 2 persons are rolling dice - wins the person obtaining the largest number.

- Some outcomes are (1,1),(2,3),(3,2)
- Say, first person rolls a black die, the second one white:

$$(\bullet, \bullet), (\bullet, \bullet), (\bullet, \bullet)$$

• The Sample Space is:



- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

Case 2: Order is not important. Example:

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

Case 2: Order is not important. Example: 2 persons are playing Nardi at Besedka.

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

Case 2: Order is not important. Example: 2 persons are playing Nardi at Besedka.

• Some outcomes are (1,1),(2,3), panj-u-du, dord-djhar

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

Case 2: Order is not important. Example: 2 persons are playing Nardi at Besedka.

- Some outcomes are (1,1),(2,3), panj-u-du, dord-djhar
- In this case rolling (2,3) or (3,2) are the same

- Experiment: we are rolling 2 dice.
- Important: we need to specify (depends on the problem we are solving) if the order is important for us or not.

Case 2: Order is not important. Example: 2 persons are playing Nardi at Besedka.

- Some outcomes are (1,1),(2,3), panj-u-du, dord-djhar
- In this case rolling (2,3) or (3,2) are the same
- The Sample Space is:

#### **Events**

Usually in the Probability Theory (and in real-life situations), we are interested not in particular outcomes and their probabilities, but also in some groups of outcomes and their probabilities.



<sup>&</sup>lt;sup>1</sup>Some Restrictions may apply!

#### **Events**

Usually in the Probability Theory (and in real-life situations), we are interested not in particular outcomes and their probabilities, but also in some groups of outcomes and their probabilities.

An **Event** in some Experiment is some collection of Outcomes of that Experiment. Mathematically, every subset of the Sample Space is called an **Event**<sup>1</sup>.



<sup>&</sup>lt;sup>1</sup>Some Restrictions may apply!

• Experiment: Rolling a die

- Experiment: Rolling a die
- Sample Space =  $\Omega = \{1, 2, 3, 4, 5, 6\}$

- Experiment: Rolling a die
- Sample Space =  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- Some Events:
  - The Result is Odd

- Experiment: Rolling a die
- Sample Space =  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- Some Events:
  - The Result is  $Odd = \{1, 3, 5\}$
  - The Result is larger than 2

- Experiment: Rolling a die
- Sample Space =  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- Some Events:
  - The Result is  $Odd = \{1, 3, 5\}$
  - The Result is larger than  $2 = \{3, 4, 5, 6\}$
  - Any Result

- Experiment: Rolling a die
- Sample Space =  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- Some Events:
  - The Result is  $Odd = \{1, 3, 5\}$
  - The Result is larger than  $2 = \{3, 4, 5, 6\}$
  - Any Result =  $\Omega$
  - No Result

- Experiment: Rolling a die
- Sample Space =  $\Omega = \{1, 2, 3, 4, 5, 6\}$
- Some Events:
  - The Result is  $Odd = \{1, 3, 5\}$
  - The Result is larger than  $2 = \{3, 4, 5, 6\}$
  - Any Result =  $\Omega$
  - No Result  $= \emptyset$

Experiment: Waiting Time (in minutes) for the Metro train

- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome:

- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24.

- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$

- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space

- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space  $= \Omega = [0, 20]$

- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space  $= \Omega = [0, 20]$
- It is not interesting to have the probability of one outcome: say, what is the probability that the waiting time will be 3.24312456231? Exactly, I mean. The answer is

- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space  $= \Omega = [0, 20]$
- It is not interesting to have the probability of one outcome: say, what is the probability that the waiting time will be 3.24312456231? Exactly, I mean. The answer is 0.

- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space  $= \Omega = [0, 20]$
- It is not interesting to have the probability of one outcome: say, what is the probability that the waiting time will be 3.24312456231? Exactly, I mean. The answer is 0.
- So in this case we are interested in events' probabilities rather than in particular outcome probability.

- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space =  $\Omega = [0, 20]$
- It is not interesting to have the probability of one outcome: say, what is the probability that the waiting time will be 3.24312456231? Exactly, I mean. The answer is 0.
- So in this case we are interested in events' probabilities rather than in particular outcome probability.
- Some Events:
  - The WT is larger than 3



- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space  $= \Omega = [0, 20]$
- It is not interesting to have the probability of one outcome: say, what is the probability that the waiting time will be 3.24312456231? Exactly, I mean. The answer is 0.
- So in this case we are interested in events' probabilities rather than in particular outcome probability.
- Some Events:
  - The WT is larger than 3 = (3, 20]
  - The WT is between 2 and 5, included



- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space  $= \Omega = [0, 20]$
- It is not interesting to have the probability of one outcome: say, what is the probability that the waiting time will be 3.24312456231? Exactly, I mean. The answer is 0.
- So in this case we are interested in events' probabilities rather than in particular outcome probability.
- Some Events:
  - The WT is larger than 3 = (3, 20]
  - The WT is between 2 and 5, included = [2, 5]
  - The WT is anything



- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space  $= \Omega = [0, 20]$
- It is not interesting to have the probability of one outcome: say, what is the probability that the waiting time will be 3.24312456231? Exactly, I mean. The answer is 0.
- So in this case we are interested in events' probabilities rather than in particular outcome probability.
- Some Events:
  - The WT is larger than 3 = (3, 20]
  - The WT is between 2 and 5, included = [2, 5]
  - The WT is anything  $= \Omega$
  - No Result.



- Experiment: Waiting Time (in minutes) for the Metro train
- An example of an outcome: 3.24. Another one:  $\pi$
- Sample Space  $= \Omega = [0, 20]$
- It is not interesting to have the probability of one outcome: say, what is the probability that the waiting time will be 3.24312456231? Exactly, I mean. The answer is 0.
- So in this case we are interested in events' probabilities rather than in particular outcome probability.
- Some Events:
  - The WT is larger than 3 = (3, 20]
  - $\bullet$  The WT is between 2 and 5, included = [2,5]
  - The WT is anything  $= \Omega$
  - No Result = Ø



Assume we have an Experiment with Sample Space  $\Omega$ . And assume A and B are two Events in this experiment, i.e.,

$$A \subset \Omega$$
 and  $B \subset \Omega$ .

Then we can obtain new Events from A and B:

•  $A^c = \Omega \setminus A$  - the complement of A, the negation of A;

Assume we have an Experiment with Sample Space  $\Omega$ . And assume A and B are two Events in this experiment, i.e.,

$$A \subset \Omega$$
 and  $B \subset \Omega$ .

Then we can obtain new Events from A and B:

- $A^c = \Omega \setminus A$  the complement of A, the negation of A;
- $A \cap B$  the intersection of A and B, the Event "A and B";

Assume we have an Experiment with Sample Space  $\Omega$ . And assume A and B are two Events in this experiment, i.e.,

$$A \subset \Omega$$
 and  $B \subset \Omega$ .

Then we can obtain new Events from A and B:

- $A^c = \Omega \setminus A$  the complement of A, the negation of A;
- $A \cap B$  the intersection of A and B, the Event "A and B";
- $A \cup B$  the union of A and B, the Event "A or B";

Assume we have an Experiment with Sample Space  $\Omega$ . And assume A and B are two Events in this experiment, i.e.,

$$A \subset \Omega$$
 and  $B \subset \Omega$ .

Then we can obtain new Events from A and B:

- $A^c = \Omega \setminus A$  the complement of A, the negation of A;
- $A \cap B$  the intersection of A and B, the Event "A and B";
- $A \cup B$  the union of A and B, the Event "A or B";
- $A \setminus B$  the Event "A but not B" ...

Assume we have an Experiment with Sample Space  $\Omega$ . And assume A and B are two Events in this experiment, i.e.,

$$A \subset \Omega$$
 and  $B \subset \Omega$ .

Then we can obtain new Events from A and B:

- $A^c = \Omega \setminus A$  the complement of A, the negation of A;
- $A \cap B$  the intersection of A and B, the Event "A and B";
- $A \cup B$  the union of A and B, the Event "A or B";
- $A \setminus B$  the Event "A but not B" ...

And having Events  $A_1, A_2, ..., A_k, ...$  in our Experiment, i.e.,  $A_k \subset \Omega$ , we can form

$$\bigcup_k A_k, \qquad \bigcap_k A_k, \dots$$



• What is an Experiment?

- What is an Experiment?
- What is an Outcome?

- What is an Experiment?
- What is an Outcome?
- What is the Sample Space?

- What is an Experiment?
- What is an Outcome?
- What is the Sample Space?
- What is an Event?

- What is an Experiment?
- What is an Outcome?
- What is the Sample Space?
- What is an Event?
- Is an Outcome an Event?

- What is an Experiment?
- What is an Outcome?
- What is the Sample Space?
- What is an Event?
- Is an Outcome an Event?
- If the Sample Space of an Experiment has cardinality n,  $\#\Omega = |\Omega| = card(\Omega) = n$  (i.e., the number of elements in  $\Omega$  is n), how many different Events our Experiment have?