

AUA CS 108, Statistics, Fall 2019

Lecture 31

Michael Poghosyan

YSU, AUA

michael@ysu.am, mpoghosyan@aua.am

4 Nov 2019

Contents

- ▶ Confidence Intervals (CI)
- ▶ Confidence Intervals by Chebyshev Inequality

Last Lecture ReCap

- ▶ What are the remarkable properties of MLE?

Last Lecture ReCap

- ▶ What are the remarkable properties of MLE?
- ▶ What is a Random Interval?

CI Problem Setting

Assume we have a Random Sample from a Parametric Model \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

CI Problem Setting

Assume we have a Random Sample from a Parametric Model \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

We take $\alpha \in (0, 1)$, and call the number $1 - \alpha$ the **confidence level**.

CI Problem Setting

Assume we have a Random Sample from a Parametric Model \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

We take $\alpha \in (0, 1)$, and call the number $1 - \alpha$ the **confidence level**.

Problem: Using the Random Sample, construct an interval containing the Unknown Parameter value with Probability not less than $1 - \alpha$.

CI Problem Setting

Assume we have a Random Sample from a Parametric Model \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

We take $\alpha \in (0, 1)$, and call the number $1 - \alpha$ the **confidence level**.

Problem: Using the Random Sample, construct an interval containing the Unknown Parameter value with Probability not less than $1 - \alpha$.

The usual values of the confidence level are 90%, 95%, 99%, so the usual values of α are 0.1, 0.05 and 0.01.

Definition: Assume $0 < \alpha < 1$, and let $L = L(x_1, \dots, x_n, \alpha)$, $U = U(x_1, \dots, x_n, \alpha)$ be two functions with $L(x_1, \dots, x_n, \alpha) \leq U(x_1, \dots, x_n, \alpha)$ for all $(x_1, \dots, x_n, \alpha)$.

Definition: Assume $0 < \alpha < 1$, and let $L = L(x_1, \dots, x_n, \alpha)$, $U = U(x_1, \dots, x_n, \alpha)$ be two functions with $L(x_1, \dots, x_n, \alpha) \leq U(x_1, \dots, x_n, \alpha)$ for all $(x_1, \dots, x_n, \alpha)$. The random interval

$$(L, U) = \left(L(X_1, \dots, X_n, \alpha), U(X_1, \dots, X_n, \alpha) \right)$$

is called a **confidence interval (or confidence interval estimator) for θ of confidence level $1 - \alpha$** , if for any $\theta \in \Theta$,

$$\mathbb{P}(L < \theta < U) \geq 1 - \alpha.$$

In the case we have a realization/observation of X_1, \dots, X_n , say, x_1, \dots, x_n , then the interval

$$\left(L(x_1, \dots, x_n, \alpha), U(x_1, \dots, x_n, \alpha) \right)$$

will be an **interval estimate for θ for the confidence level $(1 - \alpha)$** .

CI

In the case we have a realization/observation of X_1, \dots, X_n , say, x_1, \dots, x_n , then the interval

$$\left(L(x_1, \dots, x_n, \alpha), U(x_1, \dots, x_n, \alpha) \right)$$

will be an **interval estimate for θ for the confidence level $(1 - \alpha)$** .

Going back to our CI, CI of the confidence level $1 - \alpha$ is a Random Interval that contains θ in more than $(1 - \alpha) \cdot 100\%$ of cases.

CI, Interpretation

Note: It is important to understand, that in the CI definition

$$\mathbb{P}(L < \theta < U) \geq 1 - \alpha$$

θ is not our r.v., θ is our unknown constant Parameter, so we do not read this as “with high Probability, θ is in (L, U) ”.

CI, Interpretation

Note: It is important to understand, that in the CI definition

$$\mathbb{P}(L < \theta < U) \geq 1 - \alpha$$

θ is not our r.v., θ is our unknown constant Parameter, so we do not read this as “with high Probability, θ is in (L, U) ”. Instead, the Interval (L, U) is Random, so we read this as **with high Probability, the interval (L, U) will contain θ .**

¹But not different θ -s !!

CI, Interpretation

Note: It is important to understand, that in the CI definition

$$\mathbb{P}(L < \theta < U) \geq 1 - \alpha$$

θ is not our r.v., θ is our unknown constant Parameter, so we do not read this as “with high Probability, θ is in (L, U) ”. Instead, the Interval (L, U) is Random, so we read this as **with high Probability, the interval (L, U) will contain θ .**

So, if we will have/generate different observations, we will have different Intervals¹ (L, U) , and we want to have that most of the time that interval contains our unknown Parameter value.

¹But not different θ -s !!

CI, R Simulation

Example: Consider an example: our Model is $Exp(\lambda)$, and we have an observation from it. Let us take a Random Sample for the general case: X_1, X_2, \dots, X_n from $Exp(\lambda)$.

CI, R Simulation

Example: Consider an example: our Model is $Exp(\lambda)$, and we have an observation from it. Let us take a Random Sample for the general case: X_1, X_2, \dots, X_n from $Exp(\lambda)$. Then good **Estimator** for λ is

CI, R Simulation

Example: Consider an example: our Model is $Exp(\lambda)$, and we have an observation from it. Let us take a Random Sample for the general case: X_1, X_2, \dots, X_n from $Exp(\lambda)$. Then good **Estimator** for λ is

$$\hat{\lambda} = \frac{1}{\overline{X}}.$$

CI, R Simulation

Example: Consider an example: our Model is $Exp(\lambda)$, and we have an observation from it. Let us take a Random Sample for the general case: X_1, X_2, \dots, X_n from $Exp(\lambda)$. Then good **Estimator** for λ is

$$\hat{\lambda} = \frac{1}{\bar{X}}.$$

Now, let us take as CI

CI, R Simulation

Example: Consider an example: our Model is $Exp(\lambda)$, and we have an observation from it. Let us take a Random Sample for the general case: X_1, X_2, \dots, X_n from $Exp(\lambda)$. Then good **Estimator** for λ is


$$\hat{\lambda} = \frac{1}{\bar{X}}.$$

Now, let us take as CI

$$\left(\frac{1}{\bar{X}} - 0.1, \frac{1}{\bar{X}} + 0.1 \right)$$

and do some simulations:

CI, R Simulation



CI, R Simulation, Code

#CI Idea, Exponential Model

```
lambda <- 0.41
```

```
conf.level <- 0.95; a = 1 - conf.level
```

```
sample.size <- 50; no.of.intervals <- 100
```

```
epsilon <- 0.1
```

```
plot.new()
```

```
plot.window(xlim = c(0,no.of.intervals), ylim = c(0.2,0.8))
```

```
axis(1); axis(2)
```

```
title("Exponential Model, CI, (1/mean - 0.1, 1/mean + 0.1)")
```

```
for(i in 1:no.of.intervals){
```

```
  x <- rexp(sample.size, rate = lambda)
```

```
  lo <- 1/mean(x) - epsilon; up <- 1/mean(x) + epsilon
```

```
  if(lo > lambda || up < lambda){
```

```
    segments(c(i), c(lo), c(i), c(up), col = "red")
```

```
  }
```

```
  else{
```

```
    segments(c(i), c(lo), c(i), c(up))
```

```
  }
```

```
}
```

```
abline(h = lambda, lwd = 2, col = "blue")
```

Methods to obtain Confidence Intervals

We will consider several methods to construct CIs:

- ▶ Chebyshev Inequality Based;
- ▶ Pivotal Quantity Based

Methods to obtain Confidence Intervals

We will consider several methods to construct CIs:

- ▶ Chebyshev Inequality Based;
- ▶ Pivotal Quantity Based

And then we will talk about Asymptotic CIs.

Prob Refresher, Chebyshev Inequality

Recall the Cheby Inequality: If X is a r.v. with finite Mean $\mathbb{E}(X)$ and Variance $Var(X)$, then for any $\varepsilon > 0$,

$$\mathbb{P}\left(|X - \mathbb{E}(X)| \geq \varepsilon\right) \leq \frac{Var(X)}{\varepsilon^2},$$

Prob Refresher, Chebyshev Inequality

Recall the Cheby Inequality: If X is a r.v. with finite Mean $\mathbb{E}(X)$ and Variance $Var(X)$, then for any $\varepsilon > 0$,

$$\mathbb{P}\left(|X - \mathbb{E}(X)| \geq \varepsilon\right) \leq \frac{Var(X)}{\varepsilon^2},$$

or, which is the same,

$$\mathbb{P}\left(|X - \mathbb{E}(X)| < \varepsilon\right) \geq 1 - \frac{Var(X)}{\varepsilon^2}.$$

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$.

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$. Assume σ^2 is known, and μ is unknown, and we want to estimate it using the Sample.

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$. Assume σ^2 is known, and μ is unknown, and we want to estimate it using the Sample. We will construct a CI by the Chebyshev inequality, of given Confidence Level $1 - \alpha$.

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$. Assume σ^2 is known, and μ is unknown, and we want to estimate it using the Sample. We will construct a CI by the Chebyshev inequality, of given Confidence Level $1 - \alpha$.

Clearly,

$$\mathbb{E}(\bar{X}) =$$

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$. Assume σ^2 is known, and μ is unknown, and we want to estimate it using the Sample. We will construct a CI by the Chebyshev inequality, of given Confidence Level $1 - \alpha$.

Clearly,

$$\mathbb{E}(\bar{X}) = \mu \quad \text{and} \quad \text{Var}(\bar{X}) =$$

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$. Assume σ^2 is known, and μ is unknown, and we want to estimate it using the Sample. We will construct a CI by the Chebyshev inequality, of given Confidence Level $1 - \alpha$.

Clearly,

$$\mathbb{E}(\bar{X}) = \mu \quad \text{and} \quad \text{Var}(\bar{X}) = \frac{\sigma^2}{n},$$

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$. Assume σ^2 is known, and μ is unknown, and we want to estimate it using the Sample. We will construct a CI by the Chebyshev inequality, of given Confidence Level $1 - \alpha$.

Clearly,

$$\mathbb{E}(\bar{X}) = \mu \quad \text{and} \quad \text{Var}(\bar{X}) = \frac{\sigma^2}{n},$$

so if we will plug \bar{X} in the Cheby Inequality, we will obtain

$$\mathbb{P}(|\bar{X} - \mu| \geq \varepsilon) \leq \frac{\sigma^2}{n \cdot \varepsilon^2}$$

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$. Assume σ^2 is known, and μ is unknown, and we want to estimate it using the Sample. We will construct a CI by the Chebyshev inequality, of given Confidence Level $1 - \alpha$.

Clearly,

$$\mathbb{E}(\bar{X}) = \mu \quad \text{and} \quad \text{Var}(\bar{X}) = \frac{\sigma^2}{n},$$

so if we will plug \bar{X} in the Cheby Inequality, we will obtain

$$\mathbb{P}\left(|\bar{X} - \mu| \geq \varepsilon\right) \leq \frac{\sigma^2}{n \cdot \varepsilon^2} \quad \text{or} \quad \mathbb{P}\left(|\bar{X} - \mu| < \varepsilon\right) \geq 1 - \frac{\sigma^2}{n \cdot \varepsilon^2}$$

for any $\varepsilon > 0$.

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$. Assume σ^2 is known, and μ is unknown, and we want to estimate it using the Sample. We will construct a CI by the Chebyshev inequality, of given Confidence Level $1 - \alpha$.

Clearly,

$$\mathbb{E}(\bar{X}) = \mu \quad \text{and} \quad \text{Var}(\bar{X}) = \frac{\sigma^2}{n},$$

so if we will plug \bar{X} in the Cheby Inequality, we will obtain

$$\mathbb{P}(|\bar{X} - \mu| \geq \varepsilon) \leq \frac{\sigma^2}{n \cdot \varepsilon^2} \quad \text{or} \quad \mathbb{P}(|\bar{X} - \mu| < \varepsilon) \geq 1 - \frac{\sigma^2}{n \cdot \varepsilon^2}$$

for any $\varepsilon > 0$. Now, take $\frac{\sigma^2}{n \cdot \varepsilon^2} = \alpha$.

CI for the Mean, Variance is given, Cheby Method

Example: Assume X_1, X_2, \dots, X_n are Independent r.v. with the same Mean $\mathbb{E}(X_k) = \mu$ and the same Variance $\text{Var}(X_k) = \sigma^2$. Assume σ^2 is known, and μ is unknown, and we want to estimate it using the Sample. We will construct a CI by the Chebyshev inequality, of given Confidence Level $1 - \alpha$.

Clearly,

$$\mathbb{E}(\bar{X}) = \mu \quad \text{and} \quad \text{Var}(\bar{X}) = \frac{\sigma^2}{n},$$

so if we will plug \bar{X} in the Cheby Inequality, we will obtain

$$\mathbb{P}(|\bar{X} - \mu| \geq \varepsilon) \leq \frac{\sigma^2}{n \cdot \varepsilon^2} \quad \text{or} \quad \mathbb{P}(|\bar{X} - \mu| < \varepsilon) \geq 1 - \frac{\sigma^2}{n \cdot \varepsilon^2}$$

for any $\varepsilon > 0$. Now, take $\frac{\sigma^2}{n \cdot \varepsilon^2} = \alpha$. Here, σ , n and α are known, so this equality will give us the value for ε :

$$\varepsilon = \frac{\sigma}{\sqrt{n \cdot \alpha}}.$$

CI for the Mean, Variance is given, Cheby Method

Plugging this value into the above Cheby Inequality, we will get

$$\mathbb{P}\left(|\bar{X} - \mu| < \frac{\sigma}{\sqrt{n \cdot \alpha}}\right) \geq 1 - \alpha,$$

CI for the Mean, Variance is given, Cheby Method

Plugging this value into the above Cheby Inequality, we will get

$$\mathbb{P}\left(|\bar{X} - \mu| < \frac{\sigma}{\sqrt{n \cdot \alpha}}\right) \geq 1 - \alpha,$$

or, in the expanded form,

$$\mathbb{P}\left(\bar{X} - \frac{\sigma}{\sqrt{n \cdot \alpha}} < \mu < \bar{X} + \frac{\sigma}{\sqrt{n \cdot \alpha}}\right) \geq 1 - \alpha$$

CI for the Mean, Variance is given, Cheby Method

Plugging this value into the above Cheby Inequality, we will get

$$\mathbb{P}\left(|\bar{X} - \mu| < \frac{\sigma}{\sqrt{n \cdot \alpha}}\right) \geq 1 - \alpha,$$

or, in the expanded form,

$$\mathbb{P}\left(\bar{X} - \frac{\sigma}{\sqrt{n \cdot \alpha}} < \mu < \bar{X} + \frac{\sigma}{\sqrt{n \cdot \alpha}}\right) \geq 1 - \alpha$$

This means, that the interval

$$\left(\bar{X} - \frac{\sigma}{\sqrt{n \cdot \alpha}}, \bar{X} + \frac{\sigma}{\sqrt{n \cdot \alpha}}\right)$$

is a CI for μ of Confidence Level $1 - \alpha$.

CI for the Mean, Variance is given, Cheby Method

Plugging this value into the above Cheby Inequality, we will get

$$\mathbb{P}\left(|\bar{X} - \mu| < \frac{\sigma}{\sqrt{n \cdot \alpha}}\right) \geq 1 - \alpha,$$

or, in the expanded form,

$$\mathbb{P}\left(\bar{X} - \frac{\sigma}{\sqrt{n \cdot \alpha}} < \mu < \bar{X} + \frac{\sigma}{\sqrt{n \cdot \alpha}}\right) \geq 1 - \alpha$$

This means, that the interval

$$\left(\bar{X} - \frac{\sigma}{\sqrt{n \cdot \alpha}}, \bar{X} + \frac{\sigma}{\sqrt{n \cdot \alpha}}\right)$$

is a CI for μ of Confidence Level $1 - \alpha$. URA!

Some Notes

Two notes about the obtained CI - in fact, these notes will work also for other cases too:

Some Notes

Two notes about the obtained CI - in fact, these notes will work also for other cases too:

The CI length obtained above is

$$\frac{2\sigma}{\sqrt{n \cdot \alpha}}.$$

Note: If we increase n , the CI gets narrower. This is intuitive: if we collect more data, we can estimate the parameter more precisely, we can enclose it in a smaller length interval.

Some Notes

Two notes about the obtained CI - in fact, these notes will work also for other cases too:

The CI length obtained above is

$$\frac{2\sigma}{\sqrt{n \cdot \alpha}}.$$

Note: If we increase n , the CI gets narrower. This is intuitive: if we collect more data, we can estimate the parameter more precisely, we can enclose it in a smaller length interval.

Note: If we increase the Confidence Level, i.e., if we decrease α , then the length of CI increases. This is intuitive too: if we want to be more sure where our unknown Parameter is lying, we will get a larger interval.

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model.

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

Again we use the Chebyshev Inequality for \bar{X} : for any $\varepsilon > 0$,

$$\mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X})}{\varepsilon^2}$$

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

Again we use the Chebyshev Inequality for \bar{X} : for any $\varepsilon > 0$,

$$\mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X})}{\varepsilon^2} \quad \text{or} \quad \mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| < \varepsilon) \geq 1 - \frac{\text{Var}(\bar{X})}{\varepsilon^2}.$$

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

Again we use the Chebyshev Inequality for \bar{X} : for any $\varepsilon > 0$,

$$\mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X})}{\varepsilon^2} \quad \text{or} \quad \mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| < \varepsilon) \geq 1 - \frac{\text{Var}(\bar{X})}{\varepsilon^2}.$$

For the *Bernoulli*(p) case, we have

$$\mathbb{E}(\bar{X}) =$$

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

Again we use the Chebyshev Inequality for \bar{X} : for any $\varepsilon > 0$,

$$\mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X})}{\varepsilon^2} \quad \text{or} \quad \mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| < \varepsilon) \geq 1 - \frac{\text{Var}(\bar{X})}{\varepsilon^2}.$$

For the *Bernoulli*(p) case, we have

$$\mathbb{E}(\bar{X}) = p \quad \text{and} \quad \text{Var}(\bar{X}) =$$

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

Again we use the Chebyshev Inequality for \bar{X} : for any $\varepsilon > 0$,

$$\mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X})}{\varepsilon^2} \quad \text{or} \quad \mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| < \varepsilon) \geq 1 - \frac{\text{Var}(\bar{X})}{\varepsilon^2}.$$

For the *Bernoulli*(p) case, we have

$$\mathbb{E}(\bar{X}) = p \quad \text{and} \quad \text{Var}(\bar{X}) = \frac{p(1-p)}{n}.$$

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

Again we use the Chebyshev Inequality for \bar{X} : for any $\varepsilon > 0$,

$$\mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X})}{\varepsilon^2} \quad \text{or} \quad \mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| < \varepsilon) \geq 1 - \frac{\text{Var}(\bar{X})}{\varepsilon^2}.$$

For the *Bernoulli*(p) case, we have

$$\mathbb{E}(\bar{X}) = p \quad \text{and} \quad \text{Var}(\bar{X}) = \frac{p(1-p)}{n}.$$

Hence,

$$\mathbb{P}(|\bar{X} - p| < \varepsilon) \geq 1 - \frac{p(1-p)}{n \cdot \varepsilon^2}.$$

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

Again we use the Chebyshev Inequality for \bar{X} : for any $\varepsilon > 0$,

$$\mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X})}{\varepsilon^2} \quad \text{or} \quad \mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| < \varepsilon) \geq 1 - \frac{\text{Var}(\bar{X})}{\varepsilon^2}.$$

For the *Bernoulli*(p) case, we have

$$\mathbb{E}(\bar{X}) = p \quad \text{and} \quad \text{Var}(\bar{X}) = \frac{p(1-p)}{n}.$$

Hence,

$$\mathbb{P}(|\bar{X} - p| < \varepsilon) \geq 1 - \frac{p(1-p)}{n \cdot \varepsilon^2}.$$

Here, on the RHS, we have the unknown parameter value p , which is not desirable.

CI for the Proportion, Cheby Method

Example: Now, let us construct a CI of CLevel $1 - \alpha$ for p in the *Bernoulli*(p) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

Again we use the Chebyshev Inequality for \bar{X} : for any $\varepsilon > 0$,

$$\mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| \geq \varepsilon) \leq \frac{\text{Var}(\bar{X})}{\varepsilon^2} \quad \text{or} \quad \mathbb{P}(|\bar{X} - \mathbb{E}(\bar{X})| < \varepsilon) \geq 1 - \frac{\text{Var}(\bar{X})}{\varepsilon^2}.$$

For the *Bernoulli*(p) case, we have

$$\mathbb{E}(\bar{X}) = p \quad \text{and} \quad \text{Var}(\bar{X}) = \frac{p(1-p)}{n}.$$

Hence,

$$\mathbb{P}(|\bar{X} - p| < \varepsilon) \geq 1 - \frac{p(1-p)}{n \cdot \varepsilon^2}.$$

Here, on the RHS, we have the unknown parameter value p , which is not desirable. To get rid of that, we use the estimate

$$p(1-p) \leq \frac{1}{4}, \text{ so } \mathbb{P}(|\bar{X} - p| < \varepsilon) \geq 1 - \frac{1}{4n \cdot \varepsilon^2}.$$

CI for the Proportion, Cheby Method, Cont'd

On the RHS, for the CI, we want to have $1 - \alpha$. So, as in the previous Example, we choose

$$\frac{1}{4n \cdot \varepsilon^2} = \alpha,$$

CI for the Proportion, Cheby Method, Cont'd

On the RHS, for the CI, we want to have $1 - \alpha$. So, as in the previous Example, we choose

$$\frac{1}{4n \cdot \varepsilon^2} = \alpha, \quad i.e. \quad \varepsilon = \frac{1}{2\sqrt{n \cdot \alpha}}.$$

CI for the Proportion, Cheby Method, Cont'd

On the RHS, for the CI, we want to have $1 - \alpha$. So, as in the previous Example, we choose

$$\frac{1}{4n \cdot \varepsilon^2} = \alpha, \quad i.e. \quad \varepsilon = \frac{1}{2\sqrt{n \cdot \alpha}}.$$

Plugging into the inequality above, this will give

$$\mathbb{P}(|\bar{X} - p| < \frac{1}{2\sqrt{n \cdot \alpha}}) \geq 1 - \alpha,$$

CI for the Proportion, Cheby Method, Cont'd

On the RHS, for the CI, we want to have $1 - \alpha$. So, as in the previous Example, we choose

$$\frac{1}{4n \cdot \varepsilon^2} = \alpha, \quad i.e. \quad \varepsilon = \frac{1}{2\sqrt{n \cdot \alpha}}.$$

Plugging into the inequality above, this will give

$$\mathbb{P}(|\bar{X} - p| < \frac{1}{2\sqrt{n \cdot \alpha}}) \geq 1 - \alpha,$$

or,

$$\mathbb{P}\left(\bar{X} - \frac{1}{2\sqrt{n \cdot \alpha}} < p < \bar{X} + \frac{1}{2\sqrt{n \cdot \alpha}}\right) \geq 1 - \alpha.$$

CI for the Proportion, Cheby Method, Cont'd

On the RHS, for the CI, we want to have $1 - \alpha$. So, as in the previous Example, we choose

$$\frac{1}{4n \cdot \varepsilon^2} = \alpha, \quad i.e. \quad \varepsilon = \frac{1}{2\sqrt{n \cdot \alpha}}.$$

Plugging into the inequality above, this will give

$$\mathbb{P}(|\bar{X} - p| < \frac{1}{2\sqrt{n \cdot \alpha}}) \geq 1 - \alpha,$$

or,


$$\mathbb{P}\left(\bar{X} - \frac{1}{2\sqrt{n \cdot \alpha}} < p < \bar{X} + \frac{1}{2\sqrt{n \cdot \alpha}}\right) \geq 1 - \alpha.$$

This means that the interval

$$\left(\bar{X} - \frac{1}{2\sqrt{n \cdot \alpha}}, \bar{X} + \frac{1}{2\sqrt{n \cdot \alpha}}\right)$$

is a CI for p of level $1 - \alpha$.


CI for Bernoulli, R Simulation



Sample Size = 50, $CL = 95\%$


CI for Bernoulli, R Simulation

Bernoulli Model, CI by Cheby



Sample Size = 150, $CL = 95\%$


CI for Bernoulli, R Simulation



Sample Size = 150, $CL = 99\%$

CI for Bernoulli, R Simulation

Bernoulli Model, CI by Cheby



Sample Size = 250, $CL = 90\%$

CI, R Simulation, Code

```
#CI Idea, Bernoulli Model
p <- 0.345
conf.level <- 0.9; a = 1 - conf.level
sample.size <- 250; no.of.intervals <- 150
ME <- 1/(2*sqrt(sample.size*a)) #Margin of Error
plot.new()
plot.window(xlim = c(0,no.of.intervals), ylim = c(0,1))
axis(1); axis(2)
title("Bernoulli Model, CI by Cheby")
for(i in 1:no.of.intervals){
  x <- rbinom(sample.size, size = 1, prob = p)
  lo <- mean(x) - ME
  up <- mean(x) + ME
  if(lo > p || up < p){
    segments(c(i), c(lo), c(i), c(up), col = "red")
  }
  else{
    segments(c(i), c(lo), c(i), c(up))
  }
}
abline(h = p, lwd = 2, col = "blue")
```