

AUA CS 108, Statistics, Fall 2019

Lecture 03

Michael Poghosyan

YSU, AUA

michael@ysu.am, mpoghosyan@aua.am

30 Aug 2019

Descriptive Statistics

Contents

- ▶ Different Types of Variables
- ▶ Measurement Levels
- ▶ Frequency Tables and Plots

- ▶ **Mane's PSS:** Thursdays, 3:30 - 5PM, room TBD
- ▶ **Mane's OH:** Thursdays, 5:10 - 7:10PM, room TBD

Last Lecture ReCap

- ▶ What is a **Population**?

Last Lecture ReCap

- ▶ What is a **Population**?
- ▶ What is a **Sample**?

Last Lecture ReCap

- ▶ What is a **Population**?
- ▶ What is a **Sample**?
- ▶ Can you give an example of a Statistical Question/Problem, describe the Population and a Sample?

Last Lecture ReCap

- ▶ What is a **Population**?
- ▶ What is a **Sample**?
- ▶ Can you give an example of a Statistical Question/Problem, describe the Population and a Sample?
- ▶ What is an **Observation**?

Last Lecture ReCap

- ▶ What is a **Population**?
- ▶ What is a **Sample**?
- ▶ Can you give an example of a Statistical Question/Problem, describe the Population and a Sample?
- ▶ What is an **Observation**?
- ▶ What is a **Variable**?

Last Lecture ReCap

- ▶ What is a **Population**?
- ▶ What is a **Sample**?
- ▶ Can you give an example of a Statistical Question/Problem, describe the Population and a Sample?
- ▶ What is an **Observation**?
- ▶ What is a **Variable**?
- ▶ What is a **Parameter**?

Last Lecture ReCap

- ▶ What is a **Population**?
- ▶ What is a **Sample**?
- ▶ Can you give an example of a Statistical Question/Problem, describe the Population and a Sample?
- ▶ What is an **Observation**?
- ▶ What is a **Variable**?
- ▶ What is a **Parameter**?
- ▶ What is a **Statistics** (not the Subject!)?

Last Lecture ReCap

- ▶ What is a **Population**?
- ▶ What is a **Sample**?
- ▶ Can you give an example of a Statistical Question/Problem, describe the Population and a Sample?
- ▶ What is an **Observation**?
- ▶ What is a **Variable**?
- ▶ What is a **Parameter**?
- ▶ What is a **Statistics** (not the Subject!)?
- ▶ What is a **Representative Sample**?

Few Sampling Methods

Let us recall the Definition of the **Representative Sample** (of size k):

Definition: We say that our Sample is *Representative* (obtained by a Simple Random Sampling), if it is obtained in the process where all Samples of size k have the same probability of being chosen.

Few Sampling Methods

Let us recall the Definition of the **Representative Sample** (of size k):

Definition: We say that our Sample is *Representative* (obtained by a Simple Random Sampling), if it is obtained in the process where all Samples of size k have the same probability of being chosen.

Simple Random Sampling is not so easy to perform, so people are using different simpler Sampling Strategies (although they are not always giving exactly Representative Samples):

Few Sampling Methods

- ▶ *Systematic (Interval) Sampling*, we fit the population into a list, enumerate it; choose n ; choose at random a starting element from the first n elements in the list; and from that element on, every n -th member of the population is selected;

Few Sampling Methods

- ▶ *Systematic (Interval) Sampling*, we fit the population into a list, enumerate it; choose n ; choose at random a starting element from the first n elements in the list; and from that element on, every n -th member of the population is selected;
- ▶ *Stratified Sampling*, where the total population is divided into subgroups (strata), that share similar characteristics, and then random sample (of corresponding size) is chosen from each strata.

Few Sampling Methods

- ▶ *Systematic (Interval) Sampling*, we fit the population into a list, enumerate it; choose n ; choose at random a starting element from the first n elements in the list; and from that element on, every n -th member of the population is selected;
- ▶ *Stratified Sampling*, where the total population is divided into subgroups (strata), that share similar characteristics, and then random sample (of corresponding size) is chosen from each strata.
- ▶ *Cluster Sampling*, where the total population is divided into subgroups (clusters), then some clusters are randomly chosen. Then we include all elements of chosen clusters into our Sample.

Classification of Data wrt its Dimension

Data can be

- ▶ **Univariate** (1D) - here the observations are on a single Variable
- ▶ **Bivariate** (2D) - here the observations are on two Variables
- ▶ **Multivariate** (n -D, $n \geq 2$) - when the observations are on more than a one Variable (usually, more than two)

Classification of Variables wrt its Type

In Statistics, we deal with 2 types of Variables:

Classification of Variables wrt its Type

In Statistics, we deal with 2 types of Variables:

- ▶ **Qualitative** or **Categorical** Variable: the value is a category, non-numerical

Classification of Variables wrt its Type

In Statistics, we deal with 2 types of Variables:

- ▶ **Qualitative** or **Categorical** Variable: the value is a category, non-numerical

Examples: *Gender* is a Categorical Variable, taking values *Male*, *Female*;

Classification of Variables wrt its Type

In Statistics, we deal with 2 types of Variables:

- ▶ **Qualitative** or **Categorical** Variable: the value is a category, non-numerical

Examples: *Gender* is a Categorical Variable, taking values *Male*, *Female*; Or *Color* and *Model* (of a car) are again Categorical

Classification of Variables wrt its Type

In Statistics, we deal with 2 types of Variables:

- ▶ **Qualitative** or **Categorical** Variable: the value is a category, non-numerical

Examples: *Gender* is a Categorical Variable, taking values *Male*, *Female*; Or *Color* and *Model* (of a car) are again Categorical

- ▶ **Quantitative** or **Numerical** Variable: the value is a number obtained from counting, measuring something etc.

Classification of Variables wrt its Type

In Statistics, we deal with 2 types of Variables:

- ▶ **Qualitative** or **Categorical** Variable: the value is a category, non-numerical

Examples: *Gender* is a Categorical Variable, taking values *Male*, *Female*; Or *Color* and *Model* (of a car) are again Categorical

- ▶ **Quantitative** or **Numerical** Variable: the value is a number obtained from counting, measuring something etc. In this case we differentiate between
 - ▶ **Discrete:** the range is finite or countably infinite

Classification of Variables wrt its Type

In Statistics, we deal with 2 types of Variables:

- ▶ **Qualitative** or **Categorical** Variable: the value is a category, non-numerical

Examples: *Gender* is a Categorical Variable, taking values *Male*, *Female*; Or *Color* and *Model* (of a car) are again Categorical

- ▶ **Quantitative** or **Numerical** Variable: the value is a number obtained from counting, measuring something etc. In this case we differentiate between
 - ▶ **Discrete:** the range is finite or countably infinite
 - ▶ **Continuous:** the range is some interval

Classification of Variables wrt its Type

In Statistics, we deal with 2 types of Variables:

- ▶ **Qualitative** or **Categorical** Variable: the value is a category, non-numerical

Examples: *Gender* is a Categorical Variable, taking values *Male*, *Female*; Or *Color* and *Model* (of a car) are again Categorical

- ▶ **Quantitative** or **Numerical** Variable: the value is a number obtained from counting, measuring something etc. In this case we differentiate between
 - ▶ **Discrete:** the range is finite or countably infinite
 - ▶ **Continuous:** the range is some interval

Examples:

- ▶ *No. of Children*, *No. of Customers*, ... are Discrete

Classification of Variables wrt its Type

In Statistics, we deal with 2 types of Variables:

- ▶ **Qualitative** or **Categorical** Variable: the value is a category, non-numerical

Examples: *Gender* is a Categorical Variable, taking values *Male*, *Female*; Or *Color* and *Model* (of a car) are again Categorical

- ▶ **Quantitative** or **Numerical** Variable: the value is a number obtained from counting, measuring something etc. In this case we differentiate between
 - ▶ **Discrete:** the range is finite or countably infinite
 - ▶ **Continuous:** the range is some interval

Examples:

- ▶ *No. of Children*, *No. of Customers*, ... are Discrete
- ▶ *Height*, *Weight*, *Age*, ... are Continuous

Remark

Remark: Of course, we can enumerate Categorical Data, say, instead of *Male*, *Female* we can just use 0 and 1. It seems that we have already a Numerical, Quantitative (Discrete) Data. But there is a difference:

Remark

Remark: Of course, we can enumerate Categorical Data, say, instead of *Male*, *Female* we can just use 0 and 1. It seems that we have already a Numerical, Quantitative (Discrete) Data. But there is a difference:

Let me give by an example: when talking about the number of children in the family, we can have the following data: 0, 2, 1, 2, 4, 6, and we can calculate, say, the average number of children in families, here 2.5.

Remark

Remark: Of course, we can enumerate Categorical Data, say, instead of *Male*, *Female* we can just use 0 and 1. It seems that we have already a Numerical, Quantitative (Discrete) Data. But there is a difference:

Let me give by an example: when talking about the number of children in the family, we can have the following data: 0, 2, 1, 2, 4, 6, and we can calculate, say, the average number of children in families, here 2.5.

But even if we are enumerating the Sex or the Color, the average Sex or the average Color is not meaningful, we cannot deal with the assigned numbers as above!

Classification of Data wrt Measurement Levels

Statistician differentiate between the Measurement Levels, make difference between different types of, say, Categorical Data.

Classification of Data wrt Measurement Levels

Statistician differentiate between the Measurement Levels, make difference between different types of, say, Categorical Data. This is because there are different Statistical approaches for each level.

Classification of Data wrt Measurement Levels

Statistician differentiate between the Measurement Levels, make difference between different types of, say, Categorical Data. This is because there are different Statistical approaches for each level.

Example: *Color* and *Stat Final Letter Grade* are Categorical Variables.

Classification of Data wrt Measurement Levels

Statistician differentiate between the Measurement Levels, make difference between different types of, say, Categorical Data. This is because there are different Statistical approaches for each level.

Example: *Color* and *Stat Final Letter Grade* are Categorical Variables. Or, *Sex* and *Year of University Study* (freshman, sophomore, junior, senior) are again Categorical.

Classification of Data wrt Measurement Levels

Statistician differentiate between the Measurement Levels, make difference between different types of, say, Categorical Data. This is because there are different Statistical approaches for each level.

Example: *Color* and *Stat Final Letter Grade* are Categorical Variables. Or, *Sex* and *Year of University Study* (freshman, sophomore, junior, senior) are again Categorical. What do you think, is there an essential difference between these Variables?

Classification of Data wrt Measurement Levels

Statistician differentiate between the Measurement Levels, make difference between different types of, say, Categorical Data. This is because there are different Statistical approaches for each level.

Example: *Color* and *Stat Final Letter Grade* are Categorical Variables. Or, *Sex* and *Year of University Study* (freshman, sophomore, junior, senior) are again Categorical. What do you think, is there an essential difference between these Variables?

Yeah, there is an **order** in the second Variables, *Stat Final Letter Grade* and *Year of University Study*.

Measurement Levels

The (commonly agreed) Measurement Levels are:

Measurement Levels

The (commonly agreed) Measurement Levels are:

- ▶ **Nominal Measurements** - This is when we assign categories to observations, and categories do not have a logical order.

Measurement Levels

The (commonly agreed) Measurement Levels are:

- ▶ **Nominal Measurements** - This is when we assign categories to observations, and categories do not have a logical order.

Say, the Variable *Gender* can take values “male”, “female”; or the Variables *Marital Status*; *blood group*; *Mobile Phone Manufacturer*;

...

Measurement Levels

The (commonly agreed) Measurement Levels are:

- ▶ **Nominal Measurements** - This is when we assign categories to observations, and categories do not have a logical order.

Say, the Variable *Gender* can take values “male”, “female”; or the Variables *Marital Status*; *blood group*; *Mobile Phone Manufacturer*;

...

- ▶ **Ordinal Measurements** - This is when we assign categories to observations, but this time we have an intrinsic order.

Measurement Levels

The (commonly agreed) Measurement Levels are:

- ▶ **Nominal Measurements** - This is when we assign categories to observations, and categories do not have a logical order.

Say, the Variable *Gender* can take values “male”, “female”; or the Variables *Marital Status*; *blood group*; *Mobile Phone Manufacturer*; ...

- ▶ **Ordinal Measurements** - This is when we assign categories to observations, but this time we have an intrinsic order. Say, *Year of study* (freshman, sopho, ...), *Letter Grade* or *Education* (HS, BS, MS, PhD) are on the Ordinal Scale

Measurement Levels

The (commonly agreed) Measurement Levels are:

- ▶ **Nominal Measurements** - This is when we assign categories to observations, and categories do not have a logical order.

Say, the Variable *Gender* can take values “male”, “female”; or the Variables *Marital Status*; *blood group*; *Mobile Phone Manufacturer*; ...

- ▶ **Ordinal Measurements** - This is when we assign categories to observations, but this time we have an intrinsic order. Say, *Year of study* (freshman, sopho, . . .), *Letter Grade* or *Education* (HS, BS, MS, PhD) are on the Ordinal Scale

Maybe one of the well-known Ordinal Scale Measurements is the **Likert Scale**:

Measurement Levels

The (commonly agreed) Measurement Levels are:

- ▶ **Nominal Measurements** - This is when we assign categories to observations, and categories do not have a logical order.

Say, the Variable *Gender* can take values “male”, “female”; or the Variables *Marital Status*; *blood group*; *Mobile Phone Manufacturer*; ...

- ▶ **Ordinal Measurements** - This is when we assign categories to observations, but this time we have an intrinsic order. Say, *Year of study* (freshman, sopho, . . .), *Letter Grade* or *Education* (HS, BS, MS, PhD) are on the Ordinal Scale

Maybe one of the well-known Ordinal Scale Measurements is the **Likert Scale**: This is our famous

Strongly Disagree | Disagree | Neither | Agree | Strongly Agree

Measurement Levels

- ▶ **Interval Measurements** - Are like Ordinal Measurements, but here the differences are meaningful, equal differences between the category levels mean equal differences between the characteristics measured. Also here zero is arbitrary (does not mean the absence of the characteristic), and ratios do not make sense.

Measurement Levels

- ▶ **Interval Measurements** - Are like Ordinal Measurements, but here the differences are meaningful, equal differences between the category levels mean equal differences between the characteristics measured. Also here zero is arbitrary (does not mean the absence of the characteristic), and ratios do not make sense. For example, in the Likert Scale, it is not clear if the difference between *Disagree* and *Strongly Disagree* is the same as the difference between *Strongly Agree* and *Agree*.

Measurement Levels

- ▶ **Interval Measurements** - Are like Ordinal Measurements, but here the differences are meaningful, equal differences between the category levels mean equal differences between the characteristics measured. Also here zero is arbitrary (does not mean the absence of the characteristic), and ratios do not make sense. For example, in the Likert Scale, it is not clear if the difference between *Disagree* and *Strongly Disagree* is the same as the difference between *Strongly Agree* and *Agree*. But, say, *dates*, *Temperature in C°* are on the Interval Scale.

Measurement Levels

- ▶ **Ratio Measurement** - Here, in addition to the Interval Scale (differences are meaningful), zero is not arbitrary (zero represents the absence of the characteristic measured), and the ratios are meaningful too.

Measurement Levels

- ▶ **Ratio Measurement** - Here, in addition to the Interval Scale (differences are meaningful), zero is not arbitrary (zero represents the absence of the characteristic measured), and the ratios are meaningful too. Say, *Weight, Height, Temperature in Kelvin* etc.

Measurement Levels

- ▶ **Ratio Measurement** - Here, in addition to the Interval Scale (differences are meaningful), zero is not arbitrary (zero represents the absence of the characteristic measured), and the ratios are meaningful too. Say, *Weight, Height, Temperature in Kelvin* etc. Say, the difference between 4KG and 2KG is the same as between 8KG and 6KG (equal differences are the same),

Measurement Levels

- ▶ **Ratio Measurement** - Here, in addition to the Interval Scale (differences are meaningful), zero is not arbitrary (zero represents the absence of the characteristic measured), and the ratios are meaningful too. Say, *Weight, Height, Temperature in Kelvin* etc. Say, the difference between 4KG and 2KG is the same as between 8KG and 6KG (equal differences are the same), and also 4KG is twice as heavy as 2KG.

Measurement Levels

- ▶ **Ratio Measurement** - Here, in addition to the Interval Scale (differences are meaningful), zero is not arbitrary (zero represents the absence of the characteristic measured), and the ratios are meaningful too. Say, *Weight, Height, Temperature in Kelvin* etc. Say, the difference between 4KG and 2KG is the same as between 8KG and 6KG (equal differences are the same), and also 4KG is twice as heavy as 2KG. 0KG means the absence of the weight.

Measurement Levels

- ▶ **Ratio Measurement** - Here, in addition to the Interval Scale (differences are meaningful), zero is not arbitrary (zero represents the absence of the characteristic measured), and the ratios are meaningful too. Say, *Weight, Height, Temperature in Kelvin* etc. Say, the difference between 4KG and 2KG is the same as between 8KG and 6KG (equal differences are the same), and also 4KG is twice as heavy as 2KG. 0KG means the absence of the weight. The same works also, say, for the *Temperature in Kelvin*.

Measurement Levels

- ▶ **Ratio Measurement** - Here, in addition to the Interval Scale (differences are meaningful), zero is not arbitrary (zero represents the absence of the characteristic measured), and the ratios are meaningful too. Say, *Weight, Height, Temperature in Kelvin* etc. Say, the difference between 4KG and 2KG is the same as between 8KG and 6KG (equal differences are the same), and also 4KG is twice as heavy as 2KG. 0KG means the absence of the weight. The same works also, say, for the *Temperature in Kelvin*. But, say, for the *Temperature in Celcius*, 0 does not mean the *absence of the temperature*, and 40°C is not twice as hot as 20°C .

Descriptive Statistics for a Univariate Data

Let me recall that we use the Descriptive Statistics at the beginning of our Statistical Study to examine, explore the dataset.

Descriptive Statistics for a Univariate Data

Let me recall that we use the Descriptive Statistics at the beginning of our Statistical Study to examine, explore the dataset.

This part of Statistics is sometimes called **Exploratory Data Analysis**, EDA.

Descriptive Statistics for a Univariate Data

Let me recall that we use the Descriptive Statistics at the beginning of our Statistical Study to examine, explore the dataset.

This part of Statistics is sometimes called **Exploratory Data Analysis**, EDA.

And we start by describing some of the *Graphical Summaries*.

Descriptive Statistics for a Univariate Data

Let me recall that we use the Descriptive Statistics at the beginning of our Statistical Study to examine, explore the dataset.

This part of Statistics is sometimes called **Exploratory Data Analysis**, EDA.

And we start by describing some of the *Graphical Summaries*.

Here, for the beginning, we will assume that we have a univariate (mostly numerical) data (dataset), x_1, x_2, \dots, x_n . In this case we will say that we are given a (univariate, 1D) dataset x .

Frequency Tables

Here we assume that we have a univariate *discrete* numerical or categorical data x_1, x_2, \dots, x_n .

Frequency Tables

Here we assume that we have a univariate *discrete* numerical or categorical data x_1, x_2, \dots, x_n .

Definition: The **frequency** of a value t in observations x_1, x_2, \dots, x_n is the number of times t occurs in observations:

Frequency of t = number of occurrences of t in data.

Frequency Tables

Here we assume that we have a univariate *discrete* numerical or categorical data x_1, x_2, \dots, x_n .

Definition: The **frequency** of a value t in observations x_1, x_2, \dots, x_n is the number of times t occurs in observations:

Frequency of t = number of occurrences of t in data.

Definition: The **relative frequency** (or percentage) of a value t in observations x_1, x_2, \dots, x_n is the ratio of frequency of t divided by the total number of observations, n :

$$\begin{aligned}\text{Relative Frequency of } t &= \frac{\text{Frequency of } t}{\text{Total Number of Observations}} = \\ &= \frac{\text{Frequency of } t}{n}.\end{aligned}$$

Frequency Tables, Example

Example: Given the following Dataset:

1, 2, 4, 7, 2, 3, 2, 1, 2, 1, 4, 1, -1

obtain the Frequency and Relative Frequency Tables.

Frequency Tables, Example

Example: Given the following Dataset:

1, 2, 4, 7, 2, 3, 2, 1, 2, 1, 4, 1, -1

obtain the Frequency and Relative Frequency Tables.

Example: Let's construct the Frequency Table of the above Dataset using **R**:

```
x <- c(1, 2, 4, 7, 2, 3, 2, 1, 2, 1, 4, 1, -1)
table(x)
```

```
## x
## -1  1  2  3  4  7
##  1  4  4  1  2  1
```

Frequency Tables, Example

Now, consider the *iris* dataset in **R**:

```
head(iris)
```


| ## | Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|------|--------------|-------------|--------------|-------------|---------|
| ## 1 | 5.1 | 3.5 | 1.4 | 0.2 | setosa |
| ## 2 | 4.9 | 3.0 | 1.4 | 0.2 | setosa |
| ## 3 | 4.7 | 3.2 | 1.3 | 0.2 | setosa |
| ## 4 | 4.6 | 3.1 | 1.5 | 0.2 | setosa |
| ## 5 | 5.0 | 3.6 | 1.4 | 0.2 | setosa |
| ## 6 | 5.4 | 3.9 | 1.7 | 0.4 | setosa |

Frequency Tables, Example

Now, consider the *iris* dataset in **R**:

```
head(iris)
```

| ## | Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|------|--------------|-------------|--------------|-------------|---------|
| ## 1 | 5.1 | 3.5 | 1.4 | 0.2 | setosa |
| ## 2 | 4.9 | 3.0 | 1.4 | 0.2 | setosa |
| ## 3 | 4.7 | 3.2 | 1.3 | 0.2 | setosa |
| ## 4 | 4.6 | 3.1 | 1.5 | 0.2 | setosa |
| ## 5 | 5.0 | 3.6 | 1.4 | 0.2 | setosa |
| ## 6 | 5.4 | 3.9 | 1.7 | 0.4 | setosa |



Frequency Tables, Example, Cont'd

To get the *Species* Variable of the iris Dataset, we use

```
iris$Species
```

Frequency Tables, Example, Cont'd

To get the *Species* Variable of the iris Dataset, we use

```
iris$Species
```

And to calculate the Frequency of each of the Species, we use

```
table(iris$Species)
```

```
##
```


```
##      setosa versicolor  virginica
```

```
##          50          50          50
```

LineGraph and BarPlot

Now, let us visualize our Frequency Table:

```
barplot(table(iris$Species))
```



LineGraph and Barplot

Another standard Dataset, *mtcars*, again about cars 😊:

```
head(mtcars, 3)
```

| ## | | mpg | cyl | disp | hp | drat | wt | qsec | vs | am | gear | c |
|----|---------------|------|-----|------|-----|------|-------|-------|----|----|------|---|
| ## | Mazda RX4 | 21.0 | 6 | 160 | 110 | 3.90 | 2.620 | 16.46 | 0 | 1 | 4 | |
| ## | Mazda RX4 Wag | 21.0 | 6 | 160 | 110 | 3.90 | 2.875 | 17.02 | 0 | 1 | 4 | |
| ## | Datsun 710 | 22.8 | 4 | 108 | 93 | 3.85 | 2.320 | 18.61 | 1 | 1 | 4 | |


LineGraph and Barplot

Another standard Dataset, *mtcars*, again about cars ☺:

```
head(mtcars, 3)
```

| ## | | mpg | cyl | disp | hp | drat | wt | qsec | vs | am | gear | c |
|----|---------------|------|-----|------|-----|------|-------|-------|----|----|------|---|
| ## | Mazda RX4 | 21.0 | 6 | 160 | 110 | 3.90 | 2.620 | 16.46 | 0 | 1 | 4 | |
| ## | Mazda RX4 Wag | 21.0 | 6 | 160 | 110 | 3.90 | 2.875 | 17.02 | 0 | 1 | 4 | |
| ## | Datsun 710 | 22.8 | 4 | 108 | 93 | 3.85 | 2.320 | 18.61 | 1 | 1 | 4 | |


```
barplot(table(mtcars$cyl))
```



LineGraph and Barplot

Now, with the Line Graph:


```
plot(table(mtcars$cyl))
```



LineGraph and Barplot

More sophisticated (titiz) version:

```
x <- mtcars$cyl; y <- as.data.frame(table(x))  
a <- as.numeric(as.character(y$x)); b <- y$Freq  
plot(a,b,type="h", lwd=3, xlab = "Cylinders",  
      ylab = "Frequency", main = "Line Graph")  
points(a,b, pch=16, cex=2, col="red")
```



The Frequency Polygon

Again, same cars, but now the *carb* Variable Frequencies:

```
plot(table(mtcars$carb), type = "l")
```

