

AUA CS108, Statistics, Fall 2020

Lecture 41

Michael Poghosyan

02 Dec 2020

Contents

- ▶ Z-Test
- ▶ t -Test
- ▶ Tests for the Normal Model Variance
- ▶ Large Sample Hypothesis Testing

Test for the Mean of the Normal, σ is known: Z-Test

Model: Our Data comes from $\mathcal{N}(\mu, \sigma^2)$, σ is known; Our (unknown) Parameter is μ

Hypothesis: We are given some μ_0 , and we want to Test:

- ▶ Case 1: $\mathcal{H}_0 : \mu = \mu_0$ *vs* $\mathcal{H}_1 : \mu \neq \mu_0$
- ▶ Case 2: $\mathcal{H}_0 : \mu = \mu_0$ *vs* $\mathcal{H}_1 : \mu > \mu_0$
- ▶ Case 3: $\mathcal{H}_0 : \mu = \mu_0$ *vs* $\mathcal{H}_1 : \mu < \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Random Sample: We take $X_1, X_2, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$;

Test for the Mean of the Normal, σ is known: Z-Test

Test Statistics: We take

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}.$$

Under \mathcal{H}_0 ,

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1).$$

Test for the Mean of the Normal, σ is known: Z-Test

Rejection Region: Now we choose the **RR**. The idea is:

If \mathcal{H}_0 is True, then Z is close to 0

We consider our 3 cases:

Case 1: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu \neq \mu_0$

$$RR = \{|Z| > c\}.$$

Case 2: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu > \mu_0$

$$RR = \{Z > c\}.$$

Case 3: for Testing $\mathcal{H}_0 : \mu = \mu_0$ vs $\mathcal{H}_1 : \mu < \mu_0$

$$RR = \{Z < c\}.$$

Here the Critical Value c is yet to be determined.

Choosing the Critical Value

Now, let us choose c .

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha =$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha = \mathbb{P}(\text{Type I Error}) =$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) =$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\begin{aligned}\alpha &= \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) = \\ &= \mathbb{P}(|Z| > c \mid \mu = \mu_0) =\end{aligned}$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\begin{aligned}\alpha &= \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) = \\ &= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).\end{aligned}$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\alpha = \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) =$$

$$= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).$$

This means, geometrically, that

$$c =$$

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\begin{aligned}\alpha &= \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) = \\ &= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).\end{aligned}$$

This means, geometrically, that

$$c = z_{1-\alpha/2},$$

the $1 - \frac{\alpha}{2}$ -level quantile of the Standard Normal Distribution.

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\begin{aligned}\alpha &= \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) = \\ &= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).\end{aligned}$$

This means, geometrically, that

$$c = z_{1-\alpha/2},$$

the $1 - \frac{\alpha}{2}$ -level quantile of the Standard Normal Distribution.

So, finally, we have the Test for the Case 1: given μ_0 , σ , Observations and Significance Level α , calculate $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$.

Choosing the Critical Value

Now, let us choose c . We consider only **Case 1**:

$$RR = \{|Z| > c\}.$$

We choose c from the requirement to have a Test with Significance Level α :

$$\begin{aligned}\alpha &= \mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 | \mathcal{H}_0 \text{ is True}) = \\ &= \mathbb{P}(|Z| > c \mid \mu = \mu_0) = \mathbb{P}(|Z| > c \mid Z \sim \mathcal{N}(0, 1)).\end{aligned}$$

This means, geometrically, that

$$c = z_{1-\alpha/2},$$

the $1 - \frac{\alpha}{2}$ -level quantile of the Standard Normal Distribution.

So, finally, we have the Test for the Case 1: given μ_0 , σ , Observations and Significance Level α , calculate $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$.

- ▶ If $|Z| > z_{1-\alpha/2}$, **Reject** \mathcal{H}_0 ;
- ▶ If $|Z| \leq z_{1-\alpha/2}$, **Do Not Reject** \mathcal{H}_0 .

Z-Test, Complete Version

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ **is known**, the Parameter (our unknown) is μ ;

Z-Test, Complete Version

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ **is known**, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Z-Test, Complete Version

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ **is known**, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Z-Test, Complete Version

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ **is known**, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

Z-Test, Complete Version

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ is known, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $Z \sim \mathcal{N}(0, 1)$;

Z-Test, Complete Version

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ is known, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $Z \sim \mathcal{N}(0, 1)$;

| \mathcal{H}_1 is | RR is |
|--------------------|--------------------------------|
| $\mu \neq \mu_0$ | $ Z > z_{1-\frac{\alpha}{2}}$ |
| $\mu > \mu_0$ | $Z > z_{1-\alpha}$ |
| $\mu < \mu_0$ | $Z < z_{\alpha}$ |

t -Test

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ **is unknown**, the Parameter (our unknown) is μ ;

t -Test

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ **is unknown**, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

t -Test

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ **is unknown**, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

t -Test

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ is unknown, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where

$$S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n - 1}.$$

t -Test

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ is **unknown**, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where

$$S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n - 1}.$$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $t \sim t(n - 1)$;

t -Test

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, σ is **unknown**, the Parameter (our unknown) is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where

$$S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}.$$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $t \sim t(n-1)$;

| \mathcal{H}_1 is | RR is |
|--------------------|-------------------------------------|
| $\mu \neq \mu_0$ | $ t > t_{n-1, 1-\frac{\alpha}{2}}$ |
| $\mu > \mu_0$ | $t > t_{n-1, 1-\alpha}$ |
| $\mu < \mu_0$ | $t < t_{n-1, \alpha}$ |

t-test Example

Example: I have generated in **R** a Sample of Size 20 from $\mathcal{N}(3.12, 2^2)$ and made some rounding:

```
set.seed(20112019)
n <- 20; sigma <- 2
obs <- rnorm(n, mean = 3.12, sd = sigma)
obs <- round(obs, digits = 2); obs
```

```
##  [1]  1.80  5.60  1.10  3.20  4.91  5.15  1.76  2.47  0.
## [13]  3.98  4.79  1.98  4.50  3.52  4.13 -0.08  3.87
```

t-test Example

Example: I have generated in **R** a Sample of Size 20 from $\mathcal{N}(3.12, 2^2)$ and made some rounding:

```
set.seed(20112019)
n <- 20; sigma <- 2
obs <- rnorm(n, mean = 3.12, sd = sigma)
obs <- round(obs, digits = 2); obs
```

```
## [1] 1.80 5.60 1.10 3.20 4.91 5.15 1.76 2.47 0.
## [13] 3.98 4.79 1.98 4.50 3.52 4.13 -0.08 3.87
```

Now, let us forget about the fact that the actual value of μ is 3.12 and that $\sigma = 2$, and do some Testing, just assuming that our Observation is coming from a Normal Distribution.

t-test Example

Example: I have generated in **R** a Sample of Size 20 from $\mathcal{N}(3.12, 2^2)$ and made some rounding:

```
set.seed(20112019)
n <- 20; sigma <- 2
obs <- rnorm(n, mean = 3.12, sd = sigma)
obs <- round(obs, digits = 2); obs
```

```
## [1] 1.80 5.60 1.10 3.20 4.91 5.15 1.76 2.47 0.
## [13] 3.98 4.79 1.98 4.50 3.52 4.13 -0.08 3.87
```

Now, let us forget about the fact that the actual value of μ is 3.12 and that $\sigma = 2$, and do some Testing, just assuming that our Observation is coming from a Normal Distribution. Say, let us Test, at the 5% Significance Level, the following Hypothesis:

$$\mathcal{H}_0 : \mu = 4 \quad \text{vs} \quad \mathcal{H}_1 : \mu \neq 4.$$

Example, Cont'd

First, we calculate t -statistic:

```
mu0 <- 4;  
x.bar <- mean(obs); s <- sd(obs);  
t <- (x.bar - mu0)/(s/sqrt(n)); t  
  
## [1] -1.795358
```


Example, Cont'd

First, we calculate t -statistic:

```
mu0 <- 4;  
x.bar <- mean(obs); s <- sd(obs);  
t <- (x.bar - mu0)/(s/sqrt(n)); t
```

```
## [1] -1.795358
```

Now, we calculate the critical value, the quantile $t_{n-1, 1-\alpha/2}$:

```
a <- 0.05  
c <- qt(1-a/2, df = n-1); c
```

```
## [1] 2.093024
```

Example, Cont'd

First, we calculate t -statistic:

```
mu0 <- 4;  
x.bar <- mean(obs); s <- sd(obs);  
t <- (x.bar - mu0)/(s/sqrt(n)); t
```

```
## [1] -1.795358
```

Now, we calculate the critical value, the quantile $t_{n-1, 1-\alpha/2}$:

```
a <- 0.05  
c <- qt(1-a/2, df = n-1); c
```

```
## [1] 2.093024
```

Finally, we check if t is in RR, i.e., if $|t| > t_{n-1, 1-\alpha/2}$:

```
abs(t) > c
```

```
## [1] FALSE
```

Example, Cont'd

First, we calculate t -statistic:

```
mu0 <- 4;  
x.bar <- mean(obs); s <- sd(obs);  
t <- (x.bar - mu0)/(s/sqrt(n)); t
```

```
## [1] -1.795358
```

Now, we calculate the critical value, the quantile $t_{n-1, 1-\alpha/2}$:

```
a <- 0.05  
c <- qt(1-a/2, df = n-1); c
```

```
## [1] 2.093024
```

Finally, we check if t is in RR, i.e., if $|t| > t_{n-1, 1-\alpha/2}$:

```
abs(t) > c
```

```
## [1] FALSE
```

So the decision is:

Example, Cont'd

First, we calculate t -statistic:

```
mu0 <- 4;  
x.bar <- mean(obs); s <- sd(obs);  
t <- (x.bar - mu0)/(s/sqrt(n)); t
```

```
## [1] -1.795358
```

Now, we calculate the critical value, the quantile $t_{n-1, 1-\alpha/2}$:

```
a <- 0.05  
c <- qt(1-a/2, df = n-1); c
```

```
## [1] 2.093024
```

Finally, we check if t is in RR, i.e., if $|t| > t_{n-1, 1-\alpha/2}$:

```
abs(t) > c
```

```
## [1] FALSE
```

So the decision is: **Fail to Reject** \mathcal{H}_0 at 5% level.

Example, Cont'd

Now, the same, but with an **R** built-in function `t.test`:

```
t.test(obs, mu = mu0, conf.level = 0.95)
```

```
##  
##  One Sample t-test  
##  
## data:  obs  
## t = -1.7954, df = 19, p-value = 0.08852  
## alternative hypothesis: true mean is not equal to 4  
## 95 percent confidence interval:  
##  2.524009 4.112991  
## sample estimates:  
## mean of x  
##      3.3185
```

Example, Cont'd

Now, for the same Data, let us Test, at the 10% Significance Level, the following Hypothesis:

$$\mathcal{H}_0 : \mu = 3 \quad \text{vs} \quad \mathcal{H}_1 : \mu > 3.$$

Example, Cont'd

Now, for the same Data, let us Test, at the 10% Significance Level, the following Hypothesis:

$$\mathcal{H}_0 : \mu = 3 \quad \text{vs} \quad \mathcal{H}_1 : \mu > 3.$$

```
t.test(obs, mu=3, alternative="greater", conf.level=0.9)
```

```
##  
## One Sample t-test  
##  
## data: obs  
## t = 0.83906, df = 19, p-value = 0.2059  
## alternative hypothesis: true mean is greater than 3  
## 90 percent confidence interval:  
## 2.814508 Inf  
## sample estimates:  
## mean of x  
## 3.3185
```

Note

Note: In **R** `t.test` command, the default values for parameters are:

- ▶ `mu = 0`
- ▶ `alternative = "two.sided"`
- ▶ `conf.level = 0.95`

Note

Note: In **R** `t.test` command, the default values for parameters are:

- ▶ `mu = 0`
- ▶ `alternative = "two.sided"`
- ▶ `conf.level = 0.95`

Important Note: There are different ways to test a Hypothesis: one is by Test Statistics, another one is by CIs, and the next, easiest one is by p -Values.

Note

Note: In **R** `t.test` command, the default values for parameters are:

- ▶ `mu = 0`
- ▶ `alternative = "two.sided"`
- ▶ `conf.level = 0.95`

Important Note: There are different ways to test a Hypothesis: one is by Test Statistics, another one is by CIs, and the next, easiest one is by p -Values. All Statistical Software are calculating p -Values when doing testing. And the Decision based on the p -Value is:

- ▶ If $p\text{-Value} < \alpha$, then we Reject \mathcal{H}_0
- ▶ If $p\text{-Value} \geq \alpha$, then we Fail to Reject \mathcal{H}_0

t -Test Example

Example: Recall the Etruscans-Italians Problem: Scientists have a data about 84 skull sizes (widths) of adult Etruscans, and the problem was to see if Etruscans were Italians.

Also Scientists believe that the skull size is not changing much through time, and modern adult Italians skull size is in average 132.4mm.

t-Test Example

Example: Recall the Etruscans-Italians Problem: Scientists have a data about 84 skull sizes (widths) of adult Etruscans, and the problem was to see if Etruscans were Italians.

Also Scientists believe that the skull size is not changing much through time, and modern adult Italians skull size is in average 132.4mm.

We make a Hypothesis:

$$\mathcal{H}_0 : \text{Etruscans were Italians} \quad \text{vs} \quad \mathcal{H}_1 : \text{They were not}$$

t-Test Example

Example: Recall the Etruscans-Italians Problem: Scientists have a data about 84 skull sizes (widths) of adult Etruscans, and the problem was to see if Etruscans were Italians.

Also Scientists believe that the skull size is not changing much through time, and modern adult Italians skull size is in average 132.4mm.

We make a Hypothesis:

$$\mathcal{H}_0 : \text{Etruscans were Italians} \quad \text{vs} \quad \mathcal{H}_1 : \text{They were not}$$

In mathematical terms,

$$\mathcal{H}_0 : \mu = 132.4 \quad \text{vs} \quad \mathcal{H}_1 : \mu \neq 132.4$$

t-Test Example

Example: Recall the Etruscans-Italians Problem: Scientists have a data about 84 skull sizes (widths) of adult Etruscans, and the problem was to see if Etruscans were Italians.

Also Scientists believe that the skull size is not changing much through time, and modern adult Italians skull size is in average 132.4mm.

We make a Hypothesis:

$$\mathcal{H}_0 : \text{Etruscans were Italians} \quad \text{vs} \quad \mathcal{H}_1 : \text{They were not}$$

In mathematical terms,

$$\mathcal{H}_0 : \mu = 132.4 \quad \text{vs} \quad \mathcal{H}_1 : \mu \neq 132.4$$

Now let's test, at 95%, this Hypo in **R**:

t-Test Example

```
library(Rlab)
data <- etruscan
x <- data$width[data$group == "ancient"]

t.test(x, mu = 132.4)
```

```
##
## One Sample t-test
##
## data: x
## t = 17.46, df = 83, p-value < 2.2e-16
## alternative hypothesis: true mean is not equal to 132.4
## 95 percent confidence interval:
## 142.4781 145.0695
## sample estimates:
## mean of x
## 143.7738
```

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n)$;

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n)$;

| \mathcal{H}_1 is | RR is |
|----------------------------|---|
| $\sigma^2 \neq \sigma_0^2$ | $\chi^2 \notin \left[\chi_{n, \frac{\alpha}{2}}^2, \chi_{n, 1 - \frac{\alpha}{2}}^2 \right]$ |
| $\sigma^2 > \sigma_0^2$ | $\chi^2 > \chi_{n, 1 - \alpha}^2$ |
| $\sigma^2 < \sigma_0^2$ | $\chi^2 < \chi_{n, \alpha}^2$ |

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n-1)$;

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n-1)$;

| \mathcal{H}_1 is | RR is |
|----------------------------|---|
| $\sigma^2 \neq \sigma_0^2$ | $\chi^2 \notin \left[\chi_{n-1, \frac{\alpha}{2}}^2, \chi_{n-1, 1-\frac{\alpha}{2}}^2 \right]$ |
| $\sigma^2 > \sigma_0^2$ | $\chi^2 > \chi_{n-1, 1-\alpha}^2$ |
| $\sigma^2 < \sigma_0^2$ | $\chi^2 < \chi_{n-1, \alpha}^2$ |

Large Sample Hypothesis Testing

aka

Asymptotic Testing

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D} \mathcal{N}(0, 1)$;

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D} \mathcal{N}(0, 1)$;

| \mathcal{H}_1 is | RR is |
|--------------------|--------------------------------|
| $\mu \neq \mu_0$ | $ t > z_{1-\frac{\alpha}{2}}$ |
| $\mu > \mu_0$ | $t > z_{1-\alpha}$ |
| $\mu < \mu_0$ | $t < z_{\alpha}$ |

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta;$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta;$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta;$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1);$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$;

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\theta_0)}}$ or $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\hat{\theta}^{MLE})}}$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$;

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\theta_0)}}$ or $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\hat{\theta}^{MLE})}}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D} \mathcal{N}(0, 1)$;

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$;

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\theta_0)}}$ or $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\hat{\theta}^{MLE})}}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D} \mathcal{N}(0, 1)$;

| \mathcal{H}_1 is | RR is |
|------------------------|--------------------------------|
| $\theta \neq \theta_0$ | $ W > z_{1-\frac{\alpha}{2}}$ |
| $\theta > \theta_0$ | $W > z_{1-\alpha}$ |
| $\theta < \theta_0$ | $W < z_\alpha$ |