

Spectral Analysis and POD of Turbulent Pipe

M. Raba
May 27, 2022

- take fft azimuthally
 - use half of θ data to avoid aliasing
- find correlation in t, t' described in Smits2017.below.eq.2.4.

$$\mathbf{R} \left(k; m; t, t' \right) = \int_r \mathbf{u} (k; m; r, t) \mathbf{u}^* \left(k; m; r, t' \right) r \, \mathrm{d} r \tag{1}$$

- take fft in x of the above correlation to get k modes.

- the crossspectra for the kernal of the pod

$$\lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_0^\tau \mathbf{R}(k; m; t, t') \alpha^{(n)}(k; m; t') dt' = \lambda^{(n)}(k; m) \alpha^{(n)}(k; m; t) \tag{2}$$

- Find the (sorted) eigenvalues $\alpha^{(n)}$ found in (2) to solve for $\Phi^{(n)}$,

$$\lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_0^\tau \mathbf{u}_T(k; m; r, t) \alpha^{(n)*}(k; m; t) dt = \Phi_T^{(n)}(k; m; r) \lambda^{(n)}(k; m)$$