

Calculus Summary

Mathematical Notes

October 19, 2025

Contents

1	Limits	3
1.1	Definition	3
1.2	Limit Laws	3
1.3	One-Sided Limits	3
1.4	Infinite Limits	3
2	Continuity	3
2.1	Definition	3
2.2	Continuity on an Interval	3
2.3	Intermediate Value Theorem	4
3	Derivatives	4
3.1	Definition	4
3.2	Notation	4
3.3	Basic Differentiation Rules	4
3.4	Product and Quotient Rules	4
3.5	Chain Rule	4
3.6	Implicit Differentiation	4
3.7	Derivatives of Common Functions	5
4	Applications of Derivatives	5
4.1	Tangent Lines	5
4.2	Related Rates	5
4.3	Linear Approximation	5
4.4	Newton's Method	5
5	Extreme Values	5
5.1	Critical Points	5
5.2	First Derivative Test	6
5.3	Second Derivative Test	6
5.4	Extreme Value Theorem	6
6	Concavity and Inflection Points	6
6.1	Concavity	6
6.2	Inflection Points	6

7	L'Hôpital's Rule	6
8	Antiderivatives and Indefinite Integrals	6
8.1	Definition	6
8.2	Indefinite Integral	7
8.3	Basic Integration Rules	7
9	Definite Integrals	7
9.1	Definition	7
9.2	Fundamental Theorem of Calculus	7
9.3	Properties of Definite Integrals	7
10	Integration Techniques	8
10.1	Substitution Rule	8
10.2	Integration by Parts	8
10.3	Partial Fractions	8
10.4	Trigonometric Substitution	8
11	Applications of Integration	8
11.1	Area Between Curves	8
11.2	Volume of Revolution	8
11.3	Arc Length	8
11.4	Surface Area	8
12	Sequences and Series	9
12.1	Sequences	9
12.2	Series	9
12.3	Geometric Series	9
12.4	Convergence Tests	9
13	Power Series	9
13.1	Definition	9
13.2	Radius of Convergence	9
13.3	Taylor Series	9
13.4	Maclaurin Series	10
13.5	Common Maclaurin Series	10
14	Multivariable Calculus	10
14.1	Partial Derivatives	10
14.2	Gradient	10
14.3	Directional Derivative	10
14.4	Multiple Integrals	10
14.5	Change of Variables	10
15	Vector Calculus	11
15.1	Line Integrals	11
15.2	Green's Theorem	11
15.3	Divergence Theorem	11
15.4	Stokes' Theorem	11

1 Limits

1.1 Definition

The limit of $f(x)$ as x approaches a is L if:

$$\lim_{x \rightarrow a} f(x) = L$$

For every $\epsilon > 0$, there exists $\delta > 0$ such that:

$$0 < |x - a| < \delta \Rightarrow |f(x) - L| < \epsilon$$

1.2 Limit Laws

If $\lim_{x \rightarrow a} f(x) = L$ and $\lim_{x \rightarrow a} g(x) = M$, then:

- $\lim_{x \rightarrow a} [f(x) + g(x)] = L + M$
- $\lim_{x \rightarrow a} [f(x) - g(x)] = L - M$
- $\lim_{x \rightarrow a} [cf(x)] = cL$
- $\lim_{x \rightarrow a} [f(x)g(x)] = LM$
- $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{L}{M}$ (if $M \neq 0$)

1.3 One-Sided Limits

- $\lim_{x \rightarrow a^-} f(x)$: limit from the left
- $\lim_{x \rightarrow a^+} f(x)$: limit from the right
- $\lim_{x \rightarrow a} f(x) = L$ if and only if $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$

1.4 Infinite Limits

- $\lim_{x \rightarrow a} f(x) = \infty$: $f(x)$ grows without bound
- $\lim_{x \rightarrow \infty} f(x) = L$: horizontal asymptote at $y = L$

2 Continuity

2.1 Definition

A function f is continuous at a if:

$$\lim_{x \rightarrow a} f(x) = f(a)$$

2.2 Continuity on an Interval

f is continuous on $[a, b]$ if it's continuous at every point in (a, b) and:

- $\lim_{x \rightarrow a^+} f(x) = f(a)$ (continuous from the right at a)
- $\lim_{x \rightarrow b^-} f(x) = f(b)$ (continuous from the left at b)

2.3 Intermediate Value Theorem

If f is continuous on $[a, b]$ and N is between $f(a)$ and $f(b)$, then there exists $c \in (a, b)$ such that $f(c) = N$.

3 Derivatives

3.1 Definition

The derivative of f at a is:

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

3.2 Notation

- $f'(x)$ or $\frac{df}{dx}$: first derivative
- $f''(x)$ or $\frac{d^2f}{dx^2}$: second derivative
- $f^{(n)}(x)$ or $\frac{d^n f}{dx^n}$: n -th derivative

3.3 Basic Differentiation Rules

- $\frac{d}{dx}[c] = 0$ (constant rule)
- $\frac{d}{dx}[x^n] = nx^{n-1}$ (power rule)
- $\frac{d}{dx}[cf(x)] = cf'(x)$ (constant multiple rule)
- $\frac{d}{dx}[f(x) + g(x)] = f'(x) + g'(x)$ (sum rule)
- $\frac{d}{dx}[f(x) - g(x)] = f'(x) - g'(x)$ (difference rule)

3.4 Product and Quotient Rules

- Product rule: $\frac{d}{dx}[f(x)g(x)] = f'(x)g(x) + f(x)g'(x)$
- Quotient rule: $\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$

3.5 Chain Rule

$$\frac{d}{dx}[f(g(x))] = f'(g(x)) \cdot g'(x)$$

3.6 Implicit Differentiation

To find $\frac{dy}{dx}$ when y is implicitly defined by $F(x, y) = 0$:

$$\frac{\partial F}{\partial x} + \frac{\partial F}{\partial y} \frac{dy}{dx} = 0$$

3.7 Derivatives of Common Functions

$$\frac{d}{dx}[\sin x] = \cos x \quad (1)$$

$$\frac{d}{dx}[\cos x] = -\sin x \quad (2)$$

$$\frac{d}{dx}[\tan x] = \sec^2 x \quad (3)$$

$$\frac{d}{dx}[\ln x] = \frac{1}{x} \quad (4)$$

$$\frac{d}{dx}[e^x] = e^x \quad (5)$$

$$\frac{d}{dx}[a^x] = a^x \ln a \quad (6)$$

$$\frac{d}{dx}[\arcsin x] = \frac{1}{\sqrt{1-x^2}} \quad (7)$$

$$\frac{d}{dx}[\arctan x] = \frac{1}{1+x^2} \quad (8)$$

4 Applications of Derivatives

4.1 Tangent Lines

The equation of the tangent line to $y = f(x)$ at $(a, f(a))$ is:

$$y - f(a) = f'(a)(x - a)$$

4.2 Related Rates

When two or more quantities are related by an equation, their rates of change are also related.

4.3 Linear Approximation

$$f(x) \approx f(a) + f'(a)(x - a)$$

for x near a .

4.4 Newton's Method

To approximate a root of $f(x) = 0$:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

5 Extreme Values

5.1 Critical Points

A critical point of f is a number c where $f'(c) = 0$ or $f'(c)$ doesn't exist.

5.2 First Derivative Test

If c is a critical point:

- If f' changes from positive to negative at c , then $f(c)$ is a local maximum
- If f' changes from negative to positive at c , then $f(c)$ is a local minimum

5.3 Second Derivative Test

If $f'(c) = 0$ and $f''(c)$ exists:

- If $f''(c) > 0$, then $f(c)$ is a local minimum
- If $f''(c) < 0$, then $f(c)$ is a local maximum
- If $f''(c) = 0$, the test is inconclusive

5.4 Extreme Value Theorem

If f is continuous on $[a, b]$, then f attains both an absolute maximum and absolute minimum on $[a, b]$.

6 Concavity and Inflection Points

6.1 Concavity

- f is concave upward on (a, b) if $f''(x) > 0$ for all $x \in (a, b)$
- f is concave downward on (a, b) if $f''(x) < 0$ for all $x \in (a, b)$

6.2 Inflection Points

An inflection point is a point where the concavity changes, i.e., where $f''(x) = 0$ or $f''(x)$ doesn't exist.

7 L'Hôpital's Rule

If $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{0}{0}$ or $\frac{\pm\infty}{\pm\infty}$, then:

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

provided the limit on the right exists.

8 Antiderivatives and Indefinite Integrals

8.1 Definition

An antiderivative of f is a function F such that $F'(x) = f(x)$.

8.2 Indefinite Integral

$$\int f(x) dx = F(x) + C$$

where C is the constant of integration.

8.3 Basic Integration Rules

$$\int k dx = kx + C \tag{9}$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad (n \neq -1) \tag{10}$$

$$\int \frac{1}{x} dx = \ln |x| + C \tag{11}$$

$$\int e^x dx = e^x + C \tag{12}$$

$$\int \sin x dx = -\cos x + C \tag{13}$$

$$\int \cos x dx = \sin x + C \tag{14}$$

$$\int \sec^2 x dx = \tan x + C \tag{15}$$

9 Definite Integrals

9.1 Definition

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$$

where $\Delta x = \frac{b-a}{n}$ and x_i^* is any point in the i -th subinterval.

9.2 Fundamental Theorem of Calculus

- Part 1: If $F(x) = \int_a^x f(t) dt$, then $F'(x) = f(x)$
- Part 2: $\int_a^b f(x) dx = F(b) - F(a)$ where F is any antiderivative of f

9.3 Properties of Definite Integrals

- $\int_a^a f(x) dx = 0$
- $\int_a^b f(x) dx = -\int_b^a f(x) dx$
- $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$
- $\int_a^b cf(x) dx = c \int_a^b f(x) dx$
- $\int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$

10 Integration Techniques

10.1 Substitution Rule

$$\int f(g(x))g'(x) dx = \int f(u) du$$

where $u = g(x)$.

10.2 Integration by Parts

$$\int u dv = uv - \int v du$$

10.3 Partial Fractions

For rational functions, decompose into simpler fractions:

$$\frac{P(x)}{Q(x)} = \frac{A}{x-a} + \frac{B}{(x-a)^2} + \frac{Cx+D}{x^2+bx+c} + \cdots$$

10.4 Trigonometric Substitution

- For $\sqrt{a^2 - x^2}$: use $x = a \sin \theta$
- For $\sqrt{a^2 + x^2}$: use $x = a \tan \theta$
- For $\sqrt{x^2 - a^2}$: use $x = a \sec \theta$

11 Applications of Integration

11.1 Area Between Curves

$$A = \int_a^b |f(x) - g(x)| dx$$

11.2 Volume of Revolution

- Disk method: $V = \pi \int_a^b [f(x)]^2 dx$
- Washer method: $V = \pi \int_a^b ([f(x)]^2 - [g(x)]^2) dx$
- Shell method: $V = 2\pi \int_a^b x f(x) dx$

11.3 Arc Length

$$L = \int_a^b \sqrt{1 + [f'(x)]^2} dx$$

11.4 Surface Area

$$S = 2\pi \int_a^b f(x) \sqrt{1 + [f'(x)]^2} dx$$

12 Sequences and Series

12.1 Sequences

A sequence $\{a_n\}$ converges to L if:

$$\lim_{n \rightarrow \infty} a_n = L$$

12.2 Series

The series $\sum_{n=1}^{\infty} a_n$ converges if the sequence of partial sums $\{S_n\}$ converges, where:

$$S_n = \sum_{k=1}^n a_k$$

12.3 Geometric Series

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \quad \text{if } |r| < 1$$

12.4 Convergence Tests

- Divergence test: If $\lim_{n \rightarrow \infty} a_n \neq 0$, then $\sum a_n$ diverges
- Integral test: If f is positive, continuous, and decreasing, then $\sum_{n=1}^{\infty} f(n)$ and $\int_1^{\infty} f(x) dx$ both converge or both diverge
- Comparison test: If $0 \leq a_n \leq b_n$ and $\sum b_n$ converges, then $\sum a_n$ converges
- Ratio test: If $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L < 1$, then $\sum a_n$ converges
- Root test: If $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L < 1$, then $\sum a_n$ converges

13 Power Series

13.1 Definition

A power series centered at a is:

$$\sum_{n=0}^{\infty} c_n(x-a)^n$$

13.2 Radius of Convergence

The radius of convergence R is such that the series converges for $|x-a| < R$ and diverges for $|x-a| > R$.

13.3 Taylor Series

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$

13.4 Maclaurin Series

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

13.5 Common Maclaurin Series

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad (16)$$

$$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} \quad (17)$$

$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} \quad (18)$$

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n \quad (|x| < 1) \quad (19)$$

$$\ln(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1} x^n}{n} \quad (|x| < 1) \quad (20)$$

14 Multivariable Calculus

14.1 Partial Derivatives

$$\frac{\partial f}{\partial x} = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$$

14.2 Gradient

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right)$$

14.3 Directional Derivative

$$D_{\mathbf{u}} f = \nabla f \cdot \mathbf{u}$$

where \mathbf{u} is a unit vector.

14.4 Multiple Integrals

$$\iint_R f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx$$

14.5 Change of Variables

$$\iint_R f(x, y) dA = \iint_S f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

15 Vector Calculus

15.1 Line Integrals

$$\int_C f(x, y) \, ds = \int_a^b f(x(t), y(t)) \sqrt{[x'(t)]^2 + [y'(t)]^2} \, dt$$

15.2 Green's Theorem

$$\oint_C P \, dx + Q \, dy = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \, dA$$

15.3 Divergence Theorem

$$\iint_S \mathbf{F} \cdot \mathbf{n} \, dS = \iiint_E \nabla \cdot \mathbf{F} \, dV$$

15.4 Stokes' Theorem

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, dS$$