

# qmrdivideNEWTON — right matrix division $B/A$ for quaternion matrices

Solve  $X*A = B$  for  $X$  using a real embedding. Convert to Live Script: File → Save As... → Live Script (\*.mlx)

## Setup

```
hasQuat = true;
try
    quaternion(0,0,0,0);
catch
    hasQuat = false;
end
if ~hasQuat
    disp('This toolbox requires MATLAB''s built-in quaternion class (quaternion(w,x,y,z)).');
    disp('Examples in this page are skipped.');
```

return;

end

```
if exist('qmrdivideNEWTON','file') ~= 2
    thisFile = mfilename('fullpath');
    if ~isempty(thisFile)
        rootGuess = fileparts(fileparts(fileparts(thisFile))); % ../docs/source ->
        toolbox root
        if exist(fullfile(rootGuess,'qmrdivideNEWTON.m'),'file')
            addpath(rootGuess);
            rehash toolboxcache
        end
    end
end
if exist('qmrdivideNEWTON','file') ~= 2
    error('qmrdivideNEWTON not found on the MATLAB path. Add the toolbox root folder.');
```

end

## Syntax

- $X = \text{qmrdivideNEWTON}(B,A)$

## Example

```
rng(1);
n = 3;
A = quaternion(randn(n),randn(n),randn(n),randn(n));
% Make A better conditioned for the demo:
A = A + quaternion(5,0,0,0)*eye(n);

B = quaternion(randn(n),randn(n),randn(n),randn(n));
```

```
X = qmrdivideNEWTON(B,A);
```

```
disp('A:'); disp(A);
```

A:

4.351 + 0.58644i + 1.1752j + 1.4022k	-1.1096 - 1.5094i + 0.60366j + 1.2708k	-0.55868 + 0.17838 - 4.8031 -
1.1812 - 0.85189i + 2.0292j - 1.3677k	4.1544 + 0.87587i + 1.7813j + 0.066009k	
-0.75845 + 0.80032i - 0.27516j - 0.29253k	-0.57266 - 0.24279i + 1.7737j + 0.45129k	

```
disp('B:'); disp(B);
```

B:

-0.49079 - 0.34563i - 2.0187j + 0.28297k	-0.63579 + 0.92622i - 1.27j + 0.42286k	-0.15508 - 0.61212 - -1.0443 +
1.7972 - 1.1714i + 0.19974j + 0.063561k	0.60335 - 1.4817i - 0.48522j + 1.2995k	
0.5907 - 0.68559i + 0.42586j + 0.43343k	-0.53525 - 0.55806i + 0.59431j - 1.0498k	

```
disp('X = B/A (computed):'); disp(X);
```

X = B/A (computed):

-0.30797 - 0.19433i - 0.17634j - 0.087305k	-0.42424 + 0.18695i - 0.3133j + 0.14351k	0.034737 - 0.4229 - -0.19188 -
0.12961 - 0.28642i + 0.080214j + 0.20383k	0.15925 - 0.17614i - 0.49225j + 0.38786k	
0.28911 - 0.10296i - 0.035079j + 0.067281k	-0.017417 - 0.17696i + 0.18939j - 0.16635k	

```
% Check residual: ||X*A - B|| (componentwise max-norm)
```

```
R = qmtimesNEWTON(X,A) - B;
```

```
[w,x,y,z] = parts(R);
```

```
maxAbsRes = max(abs([w(:);x(:);y(:);z(:)]));
```

```
fprintf('Residual max-abs component: %.3e\n', maxAbsRes);
```

Residual max-abs component: 6.661e-16

## See also

qmldivideNEWTON, qmtimesNEWTON