

Homework 3 - 193DS Spring

Michael Torbensen

2026-05-26

```
# attaching packages

library(tidyverse)
library(janitor)
library(here)
library(effectsize)
library(readxl)
library(ggeffects)
library(rstatix)

# reading in kelp data
kelp <- read.csv(
  here("data", "temp_kelp.csv"))

# reading in personal data
my_data <- read.csv(
  here("data", "personal_data.csv"))
```

Problem 1. Giant Kelp Fronds


a. An appropriate test

To start off, in order to measure the strength and direction of the relationship between kelp length and temperature I will use Pearson's Correlation Coefficient. In order to visualize and model the relationship, I will use a simple linear regression model. The Pearson's Correlation Coefficient provides a numerical value to describe the relationship between the two variables and the simple linear regression models this relationship with a line.

b. Create a visualization

```
# base layer: ggplot
ggplot(data = kelp,
       # x-axis: temperature
       # y-axis: kelp elongation rate (cm/day)
       mapping = aes(x = temp_c,
                     y = kelp_elong)) +
# first layer: adding data points
geom_point(color = "#2a9d8f",
           alpha = 0.7,
           size = 3) +
# second layer: adding trend line with color and standard error
geom_smooth(method = "lm",
            color = "#e76f51",
            se = TRUE) +
# labeling x- and y-axis
labs(
  x = "Temperature (°C)",
  y = "Frond Elongation Rate (cm/day)",
  title = "Average Giant Kelp Frond Growth in Different Temperatures") +
# adding different theme from default
theme_minimal()
```

Average Giant Kelp Frond Growth in Different Temperatures




c. Checking assumptions and running tests

```
# Fit the model first to extract residuals ----
kelp_model <- lm(
  kelp_elong ~ temp_c,
  data = kelp)

# Check normality of residuals: Q-Q plot ----
qqnorm(residuals(kelp_model))
qqline(residuals(kelp_model),
  col = "#e76f51")
```

Normal Q-Q Plot



```
# Pearson's correlation coefficient test ----  
cor.test(kelp$temp_c,  
         kelp$kelp_elong,  
         method = "pearson")
```

Pearson's product-moment correlation

```
data: kelp$temp_c and kelp$kelp_elong  
t = -5.189, df = 30, p-value = 1.366e-05  
alternative hypothesis: true correlation is not equal to 0  
95 percent confidence interval:  
 -0.8359665 -0.4460157  
sample estimates:
```

```
cor
-0.6877489
```

```
# Simple linear regression ----
summary(kelp_model)
```

Call:

```
lm(formula = kelp_elong ~ temp_c, data = kelp)
```

Residuals:

Min	1Q	Median	3Q	Max
-1.8643	-0.5017	0.1790	0.5659	1.2177

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	14.08645	1.49219	9.440	1.72e-10 ***
temp_c	-0.43179	0.08321	-5.189	1.37e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.8665 on 30 degrees of freedom

Multiple R-squared: 0.473, Adjusted R-squared: 0.4554

F-statistic: 26.93 on 1 and 30 DF, p-value: 1.366e-05

Linearity was assessed visually using a scatterplot of temperature vs. elongation rate with a fitted regression line. Normality of residuals was checked using a Q-Q plot, and homoscedasticity was assessed using a residuals vs. fitted values plot. The scatterplot suggests a reasonably linear relationship, the Q-Q plot shows residuals falling close to the reference line, and the residuals vs. fitted plot shows no obvious pattern. All assumptions appear reasonably satisfied too.

d. Results communication

To evaluate the strength of the relationship between temperature and giant kelp frond elongation rate, I used a Pearson's correlation coefficient and a simple linear regression. These tests were appropriate because both variables are continuous and the scatterplot suggested a linear relationship between them.

Temperature was a significant negative predictor of giant kelp frond elongation rate ($r = -0.69$, $p < 0.001$, $R^2 = 0.473$). As seawater temperature increased, kelp fronds grew more slowly, suggesting that warmer water conditions affect *Macrocystis pyrifera* and inhibit its growth.


This pattern is consistent with what we know about kelp ecology, giant kelp thrives in cold, nutrient-rich water, and elevated temperatures likely symbolize conditions such as reduced upwelling that limit the nutrients available for growth.

e. Test implications

Our analysis found a strong negative relationship between seawater temperature and giant kelp frond elongation rate, meaning kelp grows significantly slower as water temperatures rise. This is very concerning given ongoing ocean warming trends. If temperatures continue to increase, we should expect noticeable declines in kelp growth rates, which could have cascading effects on the broader kelp forest ecosystem. I'd recommend we prioritize monitoring sites where temperatures are already at the higher end of the range we observed, as those are likely the most at-risk locations for growth suppression.

f. Double check my own work

```
# Check homoscedasticity: residuals vs fitted values ----
plot(fitted(kelp_model),
     residuals(kelp_model),
     xlab = "Fitted values",
     ylab = "Residuals",
     pch = 19,
     col = "#2a9d8f")
abline(h = 0,
       col = "#e76f51",
       lty = 2)
```



Problem 2. Personal data

a. Updating visualizations

```
# creating new object called my_data_clean
my_data_clean <- my_data |>
# cleaning column names so only lowercase and underscores are used
clean_names() |>
mutate(duration_sec = as.numeric(ms(duration_mm_ss)))
```

```
# define mm:ss label format
mmss_format <- function(x) sprintf("%d:%02d", x %/% 60, x %% 60)
```

```

# base layer: ggplot
ggplot(data = my_data_clean,
       # x-axis: trial number
       # y-axis: duration
       mapping = aes(x = factor(trial_number),
                     y = duration_sec,
                     color = factor(trial_number))) +
# first-layer: jitter horizontally to organize visualization
geom_jitter(width = 0.13, height = 0) +
# mean points on graph
stat_summary(fun = mean, # calculate average
            geom = "point", # display as point
            color = "#f2f533", # change color of point
            shape = 13, # change shape of point
            size = 8) +
# standard deviation error bars
stat_summary(fun.data = mean_sdl, , # calculate standard deviation
            fun.args = list(mult = 0.5), # tighter view of variability
            geom = "errorbar", # adding error-bar
            width = 0.3, # changing width
            color = "#2b2b2b") + # changing color
# assigning colors to trial 1 and trial 2
scale_color_manual(values = c("1" = "#19e36a",
                              "2" = "#7836b5")) +
# assigning a new and organized scale for y-axis
scale_y_continuous(
  breaks = seq(600, max(my_data_clean$duration_sec, na.rm = TRUE), by = 60),
  minor_breaks = seq(600, max(my_data_clean$duration_sec, na.rm = TRUE), by = 30),
  labels = mmss_format
) +
# adjusting plot (decreasing space in between categories and plot borders)
scale_x_discrete(expand = expansion(add = 0.8, mult = 0.1)) +
# changing theme from default
theme_bw(base_family = "serif") +
# changing panel and plot background
theme(
  panel.background = element_rect(fill = "lightblue"),
  plot.background = element_rect(fill = "transparent", color = NA)
) +
# taking out the minor grid
theme(panel.grid.minor = element_blank()) +
# taking out legend


```



```

theme(legend.position = "none") +
# adjusting title position
theme(plot.title = element_text(hjust = 0.5)) +
# adjusting subtitle position
theme(plot.subtitle = element_text(hjust = 0.5)) +
# adding title and labels to x- and y-axis
labs(x = "Trial Number",
      y = "Duration (mm:ss)",
      title = "Travel Time Analysis: Trial # and Duration",
      subtitle = "Latest observation; 5/26/2026")

```



```

# base layer ggplot
ggplot(data = my_data_clean,
       # x-axis = wind speed

```

```

    ## y-axis = direction of trip
    ## color varies by direction of trip
    aes(x = wind_speed_mph,
        y = duration_sec,
        color = direction_of_trip)) +

# adding points
geom_point(size = 4, alpha = 0.8) +
# adding a regression line for both plots
geom_smooth(method = "lm", se = FALSE) +
# splitting into two graphs (one per category)
facet_wrap(~ direction_of_trip) +
# assigning a new and organized scale for the y-axis
scale_y_continuous(
  breaks = seq(600, max(my_data_clean$duration_sec, na.rm = TRUE), by = 60),
  minor_breaks = seq(600, max(my_data_clean$duration_sec, na.rm = TRUE), by = 30),
  labels = mmss_format
) +
# assigning colors for each categories
scale_color_manual(values = c("home to UCSB" = "#fc5538",
                              "UCSB to home" = "#7836b5")) +

# changing theme from default
theme_light(base_family = "serif") +
# changing panel and plot background
theme(
  panel.background = element_rect(fill = "#bdf7d7"),
  plot.background = element_rect(fill = "transparent", color = NA)
) +
# labeling x- and y-axis and adding title
labs(title = "Travel Time Analysis: Wind & Duration",
     subtitle = "Latest observation; 5/26/2026",
     x = "Wind Speed (mph)",
     y = "Duration (mm:ss)") +
# adjusting the look of major and minor y grids
theme(panel.grid.minor.y = element_line(color = "gray80",
                                         linewidth = 0.5),
      panel.grid.major.y = element_line(color = "gray80",
                                         linewidth = 0.5)) +

# taking out the minor grid
theme(panel.grid.minor = element_blank()) +
# taking out legend
theme(legend.position = "none") +
# adjusting title position

```


```

theme(plot.title = element_text(hjust = 0.5)) +
# adjusting subtitle position
theme(plot.subtitle = element_text(hjust = 0.5)) +
# adjusting facet_wrap text and color
theme(
strip.text = element_text(size = 12, color = "white"),
strip.background = element_rect(fill = "#8e9e95")) +
# adjusting x- and y-axis text
theme(axis.title = element_text(size = 12))

```

Travel Time Analysis: Wind & Duration

Latest observation; 5/26/2026



b. Captions

Figure 1. Scatter plot with mean travel duration for trial number 1 and 2 and error bars displaying ± 0.5 standard deviations around the mean. Individual observation points in each

trial are denoted by x-axis (1,2) as well as colors (trial 1: green, trial 2: purple). Trial number averages are denoted by the yellow circles with X's going through. Error bars display the 95% confidence interval around each regional mean. Data source: Michael Torbensen. 2026. Average bike duration spreadsheet.

Figure 2. Two separate panels for travel direction show scatter plots with a regression line modeling the average relationship between duration (y-axis) and wind speed (x-axis). The first panel (home to UCSB) features red points indicating individual observations as well as a red regression line. The second panel (UCSB to home) features purple points indicating observations as well as a purple regression line. Data source: Michael Torbensen. 2026. Average bike duration spreadsheet.