

Stopping Power of Protons and Alpha particles in Ambient Matter (SPAM)

Michaël TOUATI

Centro de Láseres Pulsados de Salamanca (CLPU),
Edificio M5, Parque Científico, C/ Adaja 8, 37185 Villamayor,
Salamanca, Spain

5 August 2021

Abstract

SPAM (Stopping Power of Protons and Alpha particles in Ambient Matter) is a Python tool using Tkinter, PIL, Numpy and Matplotlib packages that allows for visualizing, printing and saving the stopping power and/or the Bragg's peak of protons or alpha particles in ambient matter. The code has been benchmarked against the PSTAR and ASTAR NIST databases. SPAM allows for a rapid visualization, data and image generations of the stopping power and continuous slowing down range of protons and alpha particles in ambient matter using a Python Tkinter Graphical User Interface (GUI); cf. Figure 1. New target materials at ambient conditions can be easily added. The next version release will also take into account the angular scattering of protons and alpha particles in ambient matter by using a Monte-Carlo approach of a multiple binary collision model based on [Molière, 1947] and/or [Lewis, 1950].

Tags— Python, Protons, Alpha particles, Stopping power, Continuous collisional slowing down range




Figure 1: Screenshot of SPAM GUI concerning the stopping power of non-relativistic protons in a scintillator (vinyltoluene-based solid at room temperature) and the corresponding range of a proton with an initial kinetic energy of 10 MeV

The stopping power for a proton in a material at ambient conditions

$$\frac{d\varepsilon}{ds} = \left(\frac{d\varepsilon}{ds}\right)_{\text{ele}} + \left(\frac{d\varepsilon}{ds}\right)_{\text{nuc}} \quad (1)$$

is defined as the average energy loss $d\varepsilon$ per unit path length ds . Due to the huge mass of atom nuclei relative to the electron mass, the proton slowing down is mainly due to Coulomb interaction of the proton with bound atomic electrons. According to Bethe theory [Bethe, 1933] [Staub et al., 1953], the contribution of collisions with atomic electrons can be written [Berger et al., 2016]

$$\left(\frac{d\varepsilon}{ds}\right)_{\text{ele}} = 4\pi \frac{n_e e^4 L}{m_e v^2}. \quad (2)$$

Here, e is the elementary charge, m_e is the electron mass. n_e is the atomic electron density and v is the proton velocity. The main contribution to the stopping number

$$L = L_0 + L_1 + L_2 \quad (3)$$

is

$$L_0 = \frac{1}{2} \ln \left(\frac{2m_e c^2 \beta^2 \Delta\varepsilon_m}{I(1-\beta^2)} \right) - \beta^2 - \frac{C}{Z} - \frac{\delta}{2} \quad (4)$$

due to the mean excitation energy I of the material where the proton is propagating through. Here,

$$\Delta\varepsilon_m = \frac{2m_e c^2 \beta^2}{1-\beta^2} \left[1 + \frac{2m_e}{m_p} (1-\beta^2)^{-\frac{1}{2}} + \left(\frac{m_e}{m_p} \right)^2 \right]^{-1} \quad (5)$$

is the largest possible energy loss by the proton in a single collision with a free electron, m_p the proton mass, $\beta = v/c$ and c the velocity of light in vacuum. However, as the proton kinetic energy decreases while propagating in the material, the contribution to the stopping power from interactions with bound atomic electrons in the K, L, M, ...-shells decreases and a correction term C/Z must be taken into account; see [Walske, 1952] for K-shell corrections, [Khandelwal, 1968] for L-shell corrections and [Bichsel, 1991] [Bichsel, 1992] [Bichsel, 1983] for M-shell corrections and above. Also, for relativistic proton kinetic energies, the stopping power is reduced due to the resulting electrical polarization of the medium [Fermi, 1940] [Sternheimer, 1952] [Sternheimer et al., 1982]. It is called the density effect correction because it increases with the electron density. However, considering only non-relativistic protons, this term can be neglected in all the following. The stopping number correction L_1 is the Barkas correction accounting for discrepancies between negatively and positively charged projectiles [Barkas et al., 1956] [Barkas et al., 1963]. Finally, the second stopping number correction L_2 provides the valid electronic stopping power expression when the proton velocity is large compared to the velocity of bound atomic electrons [Bloch, 1933] [Bohr, 1948]. The contribution of collisions with atomic nuclei

$$\left(\frac{d\varepsilon}{ds}\right)_{\text{nuc}} = n_{\text{nuc}} \int \Delta\varepsilon d\sigma_{\text{nuc}} \quad (6)$$

to the proton slowing down is much smaller since the recoil energy

$$\Delta\varepsilon = \frac{4\varepsilon\mu^2}{m_p m_{\text{nuc}}} \sin^2 \left(\frac{\theta}{2} \right) \quad (7)$$

received by the target atom nucleus is small. Here, $\mu = m_p m_{\text{nuc}} / (m_{\text{nuc}} + m_p)$ and θ are respectively the effective proton mass and its deflection angle in the collision center-of-mass frame, $d\Omega = 2\pi \sin\theta d\theta$ and

$$\left(\frac{d\sigma}{d\Omega}\right)_{\text{nuc}} = \frac{Z^2 e^4}{4\mu^2 v^4 \sin^4 \left(\frac{\theta}{2} \right)} \left[1 - \beta^2 \sin^2 \left(\frac{\theta}{2} \right) \right] \quad (8)$$

is the differential cross section obtained by [Mott, 1932]. However, the Bethe theory breaks down when the proton velocity is much lower than the orbital electron velocities. [Varelas and Biersack, 1970] compiled many experimental and theoretical results [Lindhard and Winther, 1964] [Newton et al., 1975] [Andersen and Ziegler, 1977] and provide a fitting formula for the electronic stopping power contribution in this low velocity regime.

In a compound, the stopping power for a proton can be approximated by a linear combination of stopping powers in each element constituents taken separately. If we note ρ the compound mass density, this so-called Bragg's additivity rule [Bragg and Kleeman, 1905] reads

$$\frac{d\varepsilon}{ds} = \sum_j \omega_j \left(\frac{d\varepsilon}{ds} \right)_j. \quad (9)$$

Here $(d\varepsilon/ds)_j$ is the stopping power for a proton in the element constituent j at ambient conditions and each linear coefficient $\omega_j = w_j \rho / \rho_j$ depends on its fraction by weight w_j and its density ρ_j .

References

- [Andersen and Ziegler, 1977] Andersen, H. H. H. H. and Ziegler, J. F. (James F.), j. a. (1977). Hydrogen stopping powers and ranges in all elements. Includes bibliographical references.
- [Barkas et al., 1956] Barkas, W. H., Birnbaum, W., and Smith, F. M. (1956). Mass-ratio method applied to the measurement of l -meson masses and the energy balance in pion decay. *Phys. Rev.*, 101:778–795.
- [Barkas et al., 1963] Barkas, W. H., Dyer, J. N., and Heckman, H. H. (1963). Resolution of the Σ^- -mass anomaly. *Phys. Rev. Lett.*, 11:26–28.
- [Berger et al., 2016] Berger, M. J., Inokuti, M., Andersen, H. H., Bichsel, H., Powers, D., Seltzer, S. . M., Thwaites, D. ., and Watt, D. E. (2016). Report 49. *Journal of the International Commission on Radiation Units and Measurements*, os25(2):NP–NP.
- [Bethe, 1933] Bethe, H. (1933). Quantenmechanik der Ein- und Zwei-Elektronen Probleme. *Handbuch der Physik*, 24/1:273P.
- [Bichsel, 1983] Bichsel, H. (1983). Stopping power of m -shell electrons for heavy charged particles. *Phys. Rev. A*, 28:1147–1150.
- [Bichsel, 1991] Bichsel, H. (1991). Stopping power of fast charged particles in heavy elements.
- [Bichsel, 1992] Bichsel, H. (1992). Stopping power and ranges of fast ions in heavy elements. *Phys. Rev. A*, 46:5761–5773.
- [Bloch, 1933] Bloch, F. (1933). Zur bremsung rasch bewegter teilchen beim durchgang durch materie. *Annalen der Physik*, 408(3):285–320.
- [Bohr, 1948] Bohr, N. (1948). The penetration of atomic particles through matter. *Matt.-Fys. Medd.*, 18.
- [Bragg and Kleeman, 1905] Bragg, W. H. and Kleeman, R. (1905). Xxxix. on the α particles of radium, and their loss of range in passing through various atoms and molecules. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 10(57):318–340.
- [Fermi, 1940] Fermi, E. (1940). The ionization loss of energy in gases and in condensed materials. *Phys. Rev.*, 57:485–493.
- [Khandelwal, 1968] Khandelwal, G. S. (1968). Shell corrections for K- and L-electrons. *Nucl. Phys.*, A116:97–111.
- [Lewis, 1950] Lewis, H. W. (1950). Multiple scattering in an infinite medium. *Phys. Rev.*, 78:526–529.
- [Lindhard and Winther, 1964] Lindhard, J. and Winther, A. (1964). Stopping power of electron gas and equipartition rule. *Kgl. Danske Videnskab. Selskab Mat.-fys. Medd.*, 34(4).
- [Molière, 1947] Molière, G. (1947). Theorie der Streuung schneller geladener Teilchen I. Einzelstreuung am abgeschirmten Coulomb-Feld. *Zeitschrift Naturforschung Teil A*, 2(3):133–145.
- [Mott, 1932] Mott, N. F. (1932). The polarisation of electrons by double scattering. *Proc. R. Soc. Lond. A*, 135:429–458.
- [Newton et al., 1975] Newton, M. D., Lucas, L. L., and Root, J. W. (1975). Proton stopping powers: Binary encounter calculations based on accurate speed distributions for target electrons. *Chemical Physics Letters*, 34(3):552 – 556.
- [Staub et al., 1953] Staub, H., Bethe, H. A., Ashkin, J., Ramsey, N. F., and Bainbridge, K. T. (1953). Experimental nuclear physics. I. Page 166, Part II, Passage of Radiations through Matter.
- [Sternheimer, 1952] Sternheimer, R. M. (1952). The density effect for the ionization loss in various materials. *Phys. Rev.*, 88:851–859.
- [Sternheimer et al., 1982] Sternheimer, R. M., Seltzer, S. M., and Berger, M. J. (1982). Density effect for the ionization loss of charged particles in various substances. *Phys. Rev. B*, 26:6067–6076.
- [Varelas and Biersack, 1970] Varelas, C. and Biersack, J. (1970). Reflection of energetic particles from atomic or ionic chains in single crystals. *Nuclear Instruments and Methods*, 79(2):213 – 218.
- [Walske, 1952] Walske, M. C. (1952). The stopping power of k -electrons. *Phys. Rev.*, 88:1283–1289.