
The magnetron (monopole) Planck's law

We start with the supposition: The magnetic monopole is a pion with a lower rest mass thus a meson π^+ with a rest mass of $1.369 \cdot 10^{-8}$ smaller than the electron mass. The electron responds with meson π^- for the reaction with a proton responds with π^+ in the magnetic bound.
The magnetic field lines never begin nor end but form loops or extend to infinity with the magnetic field due to a ring of current.



Thus, the magnetrons are force-mediating particles of the magnetic field. The magnetron has a similar onduscular equation as the photon with cylindrical symmetry and a rest mass of $0.007 eV/c^2$.
It has 3 informational quarks light = 1 (Up), gravity = 1 (Charm) & magnetism = - 2.73 (Down).
The fact that the equations of photons and magnetrons are similar and the magnetic field lines never begin nor end but form loops or extend to infinity suggests that the monopoles π^+, π^- exist standalone as in reaction.

$$e + \check{\tau} \xrightarrow{trans} e + \pi^+ + \pi^- + 2\mu\hat{} + \Delta E \qquad (1)$$

And τ neutrino will collide with a proton generating the process of continuous migration of protons into neurons generating gravity magnetism (π^+, π^-) and the μ neutrino that generates the hologram of space:

$$p + \check{\tau} \xrightarrow{transf} n + \nu\hat{} + \pi^+ + \mu\hat{} \xrightarrow{decay} (p + e + \nu\hat{}) + \mu\hat{} + \check{\pi}^- + \pi^+ = p + (e + \pi^+) = p + \mu + \Delta E \qquad (2)$$

In the nucleus, Hydrogen atoms collide τ every ~10^{-12} sec with an energy apport. In equation (1) introducing the rest mass of electron "e" and μ neutrino we obtain for the magnetron a rest mass of $0.007 eV/c^2$. Thus the magnetron I make the supposition magnetron is a pion with a lower rest muss thus a meson π^+ with a rest mass of $1.369 \cdot 10^{-8}$ smaller than the electron mass. The electron responds with meson π^- for the reaction with a proton responds with π^+ in the magnetic bound. Thus according to equations in certain conditions, a constant magnetic field B(r,z,t) = constant => is possible to generate electric field E(r,t) < > 0, not null.

The Dirac Quantization Condition implies that elementary magnetic charge: $\pi^- = \frac{\hbar}{2e}$ where e is the electron's electric charge (π^+, π^- = +/-$3.291059786 \cdot 10^{-16}$ Tesla·m^2 =Weber).
http://www.michaelvio.byethost8.com/grating.pdf

Thus we have all the elements for detecting the magnetron best to have a foldable antenna loop that can be stretched to line with 0 surfaces and remodeled into a circular surface of 1 m^2 and measure a low intensity of magnetron of a coil counting the number of monopoles from the Sun in a shielded from radiation place (in cosmic space). Ultra-low-noise SQUIDs have achieved sensitivities approaching 1 aT (atto-Tesla)= 10^{-18}T in laboratory conditions detecting monopoles one by one. The only obstacle is that there are usually billions of monopoles in the magnetic field created by a coil.

For Einstein-Planck's equation for monopole, we start with the energy of the electromagnetic field $B^2/2\mu$ And the quanta energy of the monopole RadioFrequency with cylindrical symmetry $E = h_m \cdot \nu$ with Planck "constant" depending on frequency ~2Khz÷18Ghz $h_m = h(\nu)$ so:

$$E_m = diff\left(T_q \cdot 8 \cdot \pi^2 \cdot \nu^2 \cdot \frac{h\nu}{e^{h\nu\left(\frac{2\mu_0}{B^2}\right) r_b^3} - 1}, \nu\right) \quad (3)$$

For Einstein-Planck's equation for monopole microwave, we start with the energy of the electromagnetic field $B^2/2\mu$ and the quanta energy of the monopole microwave with spherical simmetry $E = h_{mw} \cdot \nu$ with Planck "constant" depending on frequency ~950Mhz÷1.6Thz $h_m = h(\nu)$ so:

$h_{mw} = 7.890480167 \cdot 10^{-11}$ eV·s and $\Rightarrow E_{mw} = diff\left(\frac{4 \cdot T_q \cdot 8 \cdot \pi^2 \nu^2}{3} \frac{h\nu}{e^{h\nu\left(\frac{2\mu_0}{B^2}\right) \cdot r_b^3} - 1}, \nu\right)$ (4)

http://www.michaelvio.byethost8.com/Microwave.pdf

In general, for any spinning particle B the inner magnetic field of the electron the magnetic monopole $h/2 \cdot ec = 3.291 \cdot 10^{-16}$ Web thus the inner magnetic field of the electron we divide with sphere surface/$4\pi r_e^2 = 6.596 \cdot 10^{12}$ T the same formula for protons (r_e=electron radius $2.818 \cdot 10^{-15}$m). Thus $B_{elec} = 6.596 \cdot 10^{12}$ T

Property: / Particle: / Quark:	Mass Bottom	Mag.Field Down	ElectCharge Strange	Gravity Charm	Light Up	Time Top
$Monopole_M$ (π^- , π^+)	0	(+/-)2.793	0	1	1	0
$Monopole_{Mw}$ (π^- , π^+)	0	(+/-)2.793	0	0	1	1

19 Oct. 2025