The magnetron (monopole) in the interaction between the electron and proton

We start from the supposition that all the fundamental particles: proton, electron, neutron, photon, magnetron, thermal particle, and phonon interact with τ neutrino and receive a constant amount of energy when colliding with τ neutrino, increasing energy for every predetermined period of around ~$6\cdot10^{-11}$sec. For a Hydrogen atom at room temperature,1 S orbital, there is a reaction that an electron collides with a τ neutrino, as in the equation below, where mag is an abbreviation for magnetron:

$$e + \lceil\tau \xrightarrow{trans} e + \pi^+ + \pi^- + 2\mu\rceil + \Delta E \qquad (1)$$

And τ neutrino will collide with a proton, generating the process of continuous migration of protons into neurons, generating gravity magnetism (π^+, π^-) and the μ neutrino that generates the hologram of space:

$$p + \lceil\tau \xrightarrow{transf} n + \nu\rceil + \pi^+ + \mu\rceil \xrightarrow{decay} \left(p + e + \nu\rceil\right) + \mu\rceil + \lceil\pi^- + \pi^+ = p + (e + \pi^+) = p + \mu + \Delta E \quad (2)$$

In the nucleus Hydrogen atom, where the magnetron is the monopole π^+ & π^-.

Every $6{,}22\cdot10^{-11}$ sec, the electron charge is changing, forming a dipole and rearranging charge in the atom, thus forming a pair of π^+ and π^- exchanging charge between the electron and the proton.

In equation (1), introducing the rest mass e and μ, we obtain that the magnetron, which is a monopole, has a rest mass of 0.007eV/c^2. Thus, the magnetron I make the supposition magnetron is a pion with a lower rest mass, thus a meson π^+ with a rest mass of $1.369\cdot 10^{-8}$ smaller than the electron mass. The electron responds with a meson π^-, for the reaction with proton responds with π^+ in the magnetic bound.

https://michaelvio.github.io/ElecFall.pdf

The Dirac Quantization Condition implies that the elementary magnetic charge: $\pi^- = \frac{\hbar}{2e}$ where e is the electron's electric charge (π^+, π^- = +/-$3.291059786\cdot10^{-16}$ Tesla·m^2 =Weber). With the energy of the magnetron $E_0 = k\pi r_P^3 B^2/2\mu_0$ where $\mu = 4\pi\cdot10^{-7}$ H/m is the vacuum permeability, $r_P = 1.5347\cdot10^{-18}$ m is the classical radius of the proton, k ~ $6.75\cdot10^4$ is a form factor and B ~ 10^{13} T is the internal magnetic field of the proton ($E_0 \approx 190.36$ eV for radio frequency). The magnetic flux density of the magnetron $_{RadioFrequency}$ is $B_{mon} = \sqrt{k}\cdot B_m$ where $B_m = 1.000\cdot10^{13}$ T is the internal magnetic field of the proton ($B_{mon}^2 = k\cdot B_m^2$). The magnetic field of the magnetron Radiofrequency (monopole) is => $2.598\cdot10^{15}$ Tesla, with a rest mass of 0.007eV/c^2.

The magnetic flux density of the magnetron$_{microwave}$ is B_{mw} =$1.494\cdot10^{13}$T, the internal magnetic field of the proton ($E_{0mw} \approx 424.98$ eV); the microwave monopole has $B_{mon} = \sqrt{k}\cdot B_{mw} \approx 3.882\cdot10^{15}$ Tesla. This is because the ratio between the top and bottom cover is half the induction for the surface of the sphere, more precisely ($4\cdot\pi\cdot r^2 => 2\cdot\pi\cdot r^2$). The electric charge of the monopole (Radio & Microwave) is the same = ħ/(2e), but the induction flux for the surface of the cylinder or radius rb, top and bottom cover is almost half the induction for the surface of the sphere, thus the B_{mw} =$1.494\cdot10^{13}$Tesla.

Thus, we have all the elements for detecting the magnetron best to have a foldable antenna loop that can be stretched to line with 0 surfaces and remodeled into a circular surface of 1 m^2 and measure a low intensity of magnetron of a coil, counting the number of monopoles from the Sun in a shielded from radiation place (in cosmic space). Ultra-low-noise SQUIDs have achieved sensitivities approaching 1 aT (atto-Tesla) = 10^{-18}T in laboratory conditions, detecting monopoles one by one. The only obstacle is that there are usually billions of monopoles in the magnetic field created by a coil.

For Einstein-Planck's equation for monopole Radiofrequency, **h_{me}= $2.831857430\cdot10^{-9}$eV·s (3.259 eV at 1.150835 GHz),** we start with the energy of the electromagnetic field $E_0 = k\pi r_P^3 B^2/(2\mu_0)$ and the quanta energy of the monopole Radiofrequency with cylindrical symmetry $E = h_m\cdot\nu$ with Planck "constant" depending on frequency ~2Khz÷18Ghz $h_m = h(\nu)$ so:

$$E_m = diff\left(T_q\cdot 8\cdot\pi^2\cdot \quad \nu^2\cdot \frac{h\cdot \nu}{e^{\frac{h\cdot \nu}{k\cdot\pi\cdot r_P^3 B^2}\left(\frac{2\mu_0}{}\right)} - 1}, \nu\right) \qquad (3)$$

For Einstein-Planck's equation for monopole microwave, we start with the energy of the electromagnetic field $E_0 = k\pi r_P^3 B^2/(2\mu_0)$ and the quanta energy of the monopole microwave with spherical symmetry E =

$h_{mw}\cdot\nu$ with Planck "constant" depending on frequency ~6Ghz÷5.85Thz $h_m = h(\nu)$ so**:** $\mathbf{h_{mw} \sim 4.46\cdot 10^{-11}\ eV\cdot s}$ and =>

$$E_{mw} = diff\left(\frac{4\cdot T_q\cdot 8\cdot\pi^2\,\nu^2}{3}\frac{h\cdot\ \nu}{e^{\frac{h\cdot\ \nu}{\left(\frac{2\mu_0}{k\cdot\pi\cdot r_P^3 B^2}\right)}}-1},\nu\right) \quad (4)$$

Consider a point charge $q = \frac{h}{2\pi\epsilon}$(with $h = R$) located at the center of both the cylinder and the sphere. Here, $R$is the radius, the cylinder has height $h = R$, and $\epsilon$is the permittivity (assumed to be $\epsilon_0$in vacuum).
By Gauss's law, the **total electric flux** through the closed surface of the sphere of radius $R$is:

$$\Phi_{\text{sphere}} = \frac{q}{\epsilon} = \frac{h}{2\pi\epsilon^2} = \frac{R}{2\pi\epsilon^2}.$$

For the **cylinder** (radius R, height $h = R$), the flux through the **two end caps** (bases) is calculated using the solid angle subtended by the caps at the charge (located at the midpoint of the axis). The distance from the charge to each end cap is $d = h/2 = R/2$. The solid angle subtended by one circular end cap (disc of radius Rat axial distance $d = R/2$) is:

$$\Omega_1 = 2\pi(1 - \frac{d}{\sqrt{d^2 + R^2}}) = 2\pi(1 - \frac{R/2}{\sqrt{(R/2)^2 + R^2}}) = 2\pi(1 - \frac{1}{\sqrt{5}}).$$

For both end caps (symmetric):

$$\Omega_{\text{total caps}} = 2\Omega_1 = 4\pi(1 - \frac{1}{\sqrt{5}}).$$

The flux through the two end caps is:

$$\Phi_{\text{cylinder ends}} = \frac{q}{4\pi\epsilon}\cdot\Omega_{\text{total caps}} = \frac{q}{\epsilon}(1 - \frac{1}{\sqrt{5}}) = \frac{R}{2\pi\epsilon^2}(1 - \frac{1}{\sqrt{5}}).$$

The ratio of the flux through the cylinder's end caps to the flux through the sphere is:

$$\frac{\Phi_{\text{cylinder ends}}}{\Phi_{\text{sphere}}} = 1 - \frac{1}{\sqrt{5}}.$$

Numerically, $\sqrt{5} \approx 2.236$, so $1 - 1/\sqrt{5} \approx 0.5528$(or exactly $(\sqrt{5} - 1)/\sqrt{5}$).
This assumes the flux is the electric flux due to the enclosed charge, and the lateral surface of the cylinder contributes the remaining flux. $1.494170657\cdot B_m = B_{mw}$
It's quite likely that the microwave monopole has a gravitational toroidal topology, and the radiofrequency monopole has a light toroidal topology. See the informational quark table below and the paper of J.G. Williamson and Glasgow University, Department of Electronics & Electrical Engineering, Glasgow, Scotland, and M.B. van der Mark Philips Research Laboratories, Prof. Holstlaan 4, Eindhoven, The Netherlands.

https://michaelvio.github.io/MARK.TEX2.pdf

Particle: Property: Quark:	Mass Bottom	Mag. Field Down	ElectCharg eStrange	Gravity Charm	Light Up	Time Top
Monopole$_M$ (π^-, π^+)	0	(+/-)2.793	0	0	1	1
Monopole$_{Mw}$ (π^-, π^+)	0	(+/-)2.793	0	1	0	1

https://michaelvio.github.io/grating.pdf

17 Nov. 2025