Magnetic Monopole: Wave Equation with Variable Coefficient in Cylindrical Coordinates **Author:** Viorel Popescu
Framework: Teoria Onduscularǎ (TauTimp) 9Apr2026**Parameters:** $R = 4.2355 \times 10^{22}$ m, $c = 2.998 \times 10^{8}$ m/s

1. Problem Statement

We solve the wave equation with variable coefficient under axial cylindrical symmetry (f independent of azimuthal angle θ):

$$\frac{R}{r}\,\Delta f(r,z,t) = \frac{1}{c^2}\frac{\partial^2 f(r,z,t)}{\partial t^2}$$

where the cylindrical Laplacian with axial symmetry ($\partial/\partial\theta = 0$) is:

$$\Delta f = \frac{1}{r}\frac{\partial}{\partial r}\left(r\frac{\partial f}{\partial r}\right) + \frac{\partial^2 f}{\partial z^2} = \frac{\partial^2 f}{\partial r^2} + \frac{1}{r}\frac{\partial f}{\partial r} + \frac{\partial^2 f}{\partial z^2}$$

Boundary / initial conditions:

$$f(2R,\,0,\,0) = 0, \qquad \frac{\partial f}{\partial r}\Big|_{(0,0,0)} = -R$$

with $R = 4.2355 \times 10^{22}$ m and $c = 2.998 \times 10^{8}$ m/s.

2. Separation of Variables

Assume a product solution:

$$f(r,z,t) = H_3(r)\,H_4(z)\,u(t)$$

Substituting into the PDE:

$$\frac{R}{r}\left[H_4 u\left(H_3'' + \frac{1}{r}H_3'\right) + H_3 u\,H_4''\right] = \frac{1}{c^2}H_3 H_4\,\ddot{u}$$

Dividing both sides by $H_3 H_4 u$ and multiplying by r/R:

$$\frac{1}{H_3}\left(H_3'' + \frac{1}{r}H_3'\right) + \frac{H_4''}{H_4} = \frac{r}{Rc^2}\frac{\ddot{u}}{u}$$

Because the left side depends on r and z while the right side mixes r and t, consistent separation requires:

$$\frac{\ddot{u}}{u} = -\omega^2, \qquad \frac{H_4''}{H_4} = -k_z^2, \qquad \frac{1}{H_3}\left(H_3'' + \frac{1}{r}H_3'\right) = -k_r^2$$

with the dispersion-like constraint (from the variable coefficient R/r):

$$k_r^2 + k_z^2 = \frac{\omega^2 R}{r\,c^2}$$

3. Temporal Equation

$$\ddot{u}(t) + \omega^2 u(t) = 0$$

General solution:

$$\boxed{u(t) = A\cos(\omega t) + B\sin(\omega t)}$$

4. Axial (z) Equation

$$H_4''(z) + k_z^2\,H_4(z) = 0$$

General solution:

$$\boxed{H_4(z) = C\cos(k_z z) + D\sin(k_z z)}$$

5. Radial Equation — Bessel Equation of Order Zero

$$H_3'' + \frac{1}{r}H_3' + k_r^2 H_3 = 0$$

This is **Bessel's equation of order zero**. Under the substitution $x = k_r r$:

$$\frac{d^2 H_3}{dx^2} + \frac{1}{x}\frac{dH_3}{dx} + H_3 = 0$$

General solution:

$$\boxed{H_3(r) = E\,J_0(k_r r) + F\,Y_0(k_r r)}$$

where J_0 is the Bessel function of the first kind (order 0) and Y_0 is the Bessel function of the second kind (Neumann function).

Key properties:

$$J_0'(x) = -J_1(x), \qquad Y_0(0) \to -\infty \text{ (logarithmically singular)}$$

$$Y_0(x) \sim \frac{2}{\pi}\ln\frac{x}{2} + \frac{2\gamma}{\pi} \quad \text{as } x \to 0$$

6. Boundary Conditions and Quantisation

Condition 1: $f(2R, 0, 0) = 0$

At $t = 0$, $z = 0$:

$$f(2R,0,0) = H_3(2R)\cdot 1 \cdot A = 0$$

This requires $J_0(2k_r R) = 0$ (non-trivial solution), giving the quantisation:

$$\boxed{k_{r,n} = \frac{j_{0,n}}{2R}}$$

where $j_{0,n}$ are the positive zeros of J_0: $j_{0,1} \approx 2.4048$, $j_{0,2} \approx 5.5201$, …

Condition 2: $\partial_r f|_{(0,0,0)} = -R$

The derivative of Y_0 near the origin satisfies:

$$Y_0'(k_r r) = -k_r Y_1(k_r r) \approx \frac{-2}{\pi r} \quad \text{as } r \to 0$$

Thus:

$$\frac{\partial f}{\partial r}|_{r\to 0} \approx F \cdot \frac{-2}{\pi r} \cdot H_4(0) \cdot u(0) = -R$$

Setting $H_4(0) = C = 1$ and $u(0) = A = 1$ (normalisation):

$$F = \frac{\pi R^2 k_r}{2}$$

For the first mode $k_{r,1} = j_{0,1}/(2R)$:

$$\boxed{F_1 = \frac{\pi R\, j_{0,1}}{4} \approx 8.00 \times 10^{22}\ \text{m}}$$

7. Full General Solution

$$\boxed{f(r,z,t) = \sum_{m,n} \left[E_{mn} J_0(k_{r,m} r) + F_{mn} Y_0(k_{r,m} r)\right] \left[C_{mn}\cos(k_{z,n} z) + D_{mn}\sin(k_{z,n} z)\right] \left[A_{mn}\cos(\omega_{mn} t) + B_{mn}\sin(\omega_{mn} t)\right]}$$

Dominant single-mode solution (first radial mode, $k_z = 0$, z-independent):

$$\boxed{f(r,z,0) = \left[R\, J_0\left(\frac{j_{0,1} r}{2R}\right) + F_1\, Y_0\left(\frac{j_{0,1} r}{2R}\right)\right] \cos(k_z z)}$$

with $E = R$ (amplitude in metres, consistent with magnetic field interpretation).

8. Probability Density System and Construction of $H(r,\theta,z,t)$ 8.1 Probability Density PDE

The probability density of the magnetic monopole in cylindrical symmetry satisfies:

$$\frac{\partial^2 P(r,z,t)}{\partial t^2} - c^2 \frac{\partial^2 P(r,z,t)}{\partial r^2} = 0$$

This is a **1D wave equation in** r (for fixed z) with general d'Alembert solution:

$$P(r,z,t) = P_+(r + ct, z) + P_-(r - ct, z)$$

Boundary condition: $P(2R, 0,0) = 0$.

8.2 Squared-Amplitude Identification

We identify the probability density with the squared field amplitude ($|\psi|^2$ formalism):

$$P(r,z,t) = |f(r,z,t)|^2$$

At $t = 0$, $z = 0$:

$$P(r,0,0) = f(r,0,0)^2, \qquad P(2R,0,0) = f(2R,0,0)^2 = 0 \checkmark$$

8.3 Polynomial Ansatz for $P(r,0,0)$

A degree-2 polynomial satisfying the boundary condition $P(2R,0,0) = 0$ and normalised over $[0,2R]$:

$$P(r,0,0) = C_0(r - 2R)(r - r_{\min})$$

where $r_{\min}$ is determined by the normalization constraint:

$$\int_0^{2R} P\,(r,0,0)\, r\, dr = 1$$

This gives a bell-shaped interior maximum physically preferred for the monopole probability density.

8.4 Full Field Reconstruction

The magnetic monopole field in cylindrical coordinates (axial symmetry, θ-independent):

$$\boxed{H(r,z,t) = \sqrt{P(r,z,t)} = |f(r,z,t)|}$$

The vector field:

$$\boldsymbol{H}(r,z,t) = H(r,z,t)\, \hat{e}_r = f(r,z,t)\, \hat{e}_r$$

The full complex representation including phase:

$$H(r,\theta,z,t) = \sqrt{P(r,z,t)} \cdot e^{i\phi(r,z,t)}$$

where $\phi(r,z,t) = \omega t$ is the temporal phase from $u(t)$.

9. Force at $t = 0$ and Maxwell Equations 9.1 Definition of Force

From the cylindrical monopole geometry, the force is defined as:

$$F(r,z) = \frac{f(r,z,0)}{4\pi r z}$$

This is analogous to the Coulomb-type force $F = q_m B/(4\pi r^2)$ adapted to cylindrical geometry where $r \cdot z$ replaces r^2.

9.2 Evaluation at $t = 0$

$$\boxed{F(r,z) = \frac{R\,J_0\left(\frac{j_{0,1}r}{2R}\right) + F_1\,Y_0\left(\frac{j_{0,1}r}{2R}\right)}{4\pi r z}\cos(k_z z)}$$

9.3 Equality with Magnetic Force — Maxwell's Equations

The extended Maxwell equations including magnetic monopoles are:

$$\nabla \cdot \boldsymbol{E} = \frac{\rho_e}{\varepsilon_0}, \qquad \nabla \cdot \boldsymbol{B} = \mu_0\,\rho_m$$

$$\nabla \times \boldsymbol{E} = -\frac{\partial \boldsymbol{B}}{\partial t} - \mu_0 \boldsymbol{J}_m, \qquad \nabla \times \boldsymbol{B} = \mu_0 \boldsymbol{J}_e + \mu_0 \varepsilon_0 \frac{\partial \boldsymbol{E}}{\partial t}$$

For a monopole of magnetic charge g in the absence of electric sources ($\rho_e = 0, \boldsymbol{J}_e = 0$), the magnetic force per unit volume in cylindrical geometry is:

$$F_r = \frac{\mu_0\, g^2}{4\pi r^2}$$

Setting $F(r,z) = F_r$:

$$\frac{f(r,z,0)}{4\pi r z} = \frac{\mu_0 g^2}{4\pi r^2}$$

This gives:

$$f(r,z,0) = \frac{\mu_0 g^2\, z}{r}$$

Solving for the magnetic charge:

$$\boxed{g^2 = \frac{r}{\mu_0 z}\left[R\,J_0\left(\frac{j_{0,1}r}{2R}\right) + F_1\,Y_0\left(\frac{j_{0,1}r}{2R}\right)\right]\cos(k_z z)}$$

Physical interpretation: The magnetic charge g is a function of position (r,z), reflecting the spatially distributed nature of the monopole field in the TauTimp framework. The Dirac quantisation condition $g = n \cdot g_0$ with $g_0 = hc/(2\mu_0 e)$ can be applied as a quantisation constraint on the spatial modes.

9.4 Connection to Wave Equation for $\boldsymbol{H}$

In the absence of electric sources, the magnetic field satisfies the d'Alembertian:

$$\Box\, \boldsymbol{H} = \frac{1}{c^2}\frac{\partial^2 \boldsymbol{H}}{\partial t^2} - \nabla^2 \boldsymbol{H} = 0$$

With the TauTimp variable coefficient R/r from the monopole geometry:

$$\frac{R}{r}\left(\frac{\partial^2 H}{\partial r^2} + \frac{1}{r}\frac{\partial H}{\partial r} + \frac{\partial^2 H}{\partial z^2}\right) = \frac{1}{c^2}\frac{\partial^2 H}{\partial t^2}$$

This is exactly the governing equation of Section 1. ✓

10. Magnetic Field $B(r,z,t)$ — Modified Wave Equation 10.1 New Governing Equation

We now derive and solve the magnetic field equation with damping/source term:

$$\frac{\partial^2 B}{\partial r^2} = \frac{1}{c^2}\frac{\partial^2 B}{\partial t^2} + \frac{\partial B}{\partial t}\cdot\frac{1}{Rc}$$

Boundary conditions:

$$B(2R,z,t) = 0, \qquad \frac{\partial B}{\partial r}\Big|_{r=0} = R$$

This is a **1D hyperbolic PDE in** (r,t) (for fixed z) with a first-order time-derivative damping/dissipation term.

10.2 Physical Origin of the Damping Term

The term $\frac{1}{Rc}\frac{\partial B}{\partial t}$ arises from the cosmological boundary condition: the factor $1/(Rc)$ is the inverse of the light-travel time across the cosmological scale R, giving a characteristic dissipation rate:

$$\gamma = \frac{1}{2Rc} = \frac{1}{2 \times 4.2355 \times 10^{22} \times 2.998 \times 10^{8}} \approx 3.94 \times 10^{-32}\ \mathrm{s}^{-1}$$

This is an extremely slow dissipation, consistent with cosmological time scales.

10.3 Separation of Variables

Assume $B(r,z,t) = \mathcal{R}(r)\,\mathcal{T}(t)$. Substituting:

$$\mathcal{T}\,\mathcal{R}'' = \frac{1}{c^2}\mathcal{R}\,\ddot{\mathcal{T}} + \frac{1}{Rc}\mathcal{R}\,\dot{\mathcal{T}}$$

Dividing by $\mathcal{R}\mathcal{T}$:

$$\frac{\mathcal{R}''}{\mathcal{R}} = \frac{1}{c^2}\frac{\ddot{\mathcal{T}}}{\mathcal{T}} + \frac{1}{Rc}\frac{\dot{\mathcal{T}}}{\mathcal{T}} = -\lambda^2$$

10.4 Temporal Equation with Damping

$$\frac{1}{c^2}\ddot{\mathcal{T}} + \frac{1}{Rc}\dot{\mathcal{T}} + \lambda^2 \mathcal{T} = 0$$

Multiply by c^2:

$$\ddot{\mathcal{T}} + \frac{c}{R}\dot{\mathcal{T}} + \lambda^2 c^2 \mathcal{T} = 0$$

This is a **damped harmonic oscillator** with:

$$2\zeta\omega_n = \frac{c}{R} \equiv \frac{1}{\tau_R}, \qquad \omega_n = \lambda c$$

The damping ratio:

$$\zeta = \frac{c}{2R\omega_n} = \frac{c}{2R\lambda c} = \frac{1}{2\lambda R}$$

Characteristic equation:

$$s^2 + \frac{c}{R}s + \lambda^2 c^2 = 0$$

$$s = -\frac{c}{2R} \pm \sqrt{\frac{c^2}{4R^2} - \lambda^2 c^2} = -\frac{c}{2R} \pm c\sqrt{\frac{1}{4R^2} - \lambda^2}$$

Three regimes:
Underdamped ($\lambda > 1/(2R)$, i.e. $\zeta < 1$):

$$s = -\frac{c}{2R} \pm i\,c\sqrt{\lambda^2 - \frac{1}{4R^2}} \equiv -\alpha \pm i\Omega$$

$$\boxed{\mathcal{T}(t) = e^{-\alpha t}[A\cos(\Omega t) + B\sin(\Omega t)]}$$

with $\alpha = \frac{c}{2R}$ and $\Omega = c\sqrt{\lambda^2 - \frac{1}{4R^2}}$.
Overdamped ($\lambda < 1/(2R)$):

$$\boxed{\mathcal{T}(t) = e^{-\alpha t}\left[Ae^{\beta t} + Be^{-\beta t}\right], \quad \beta = c\sqrt{\frac{1}{4R^2} - \lambda^2}}$$

Critically damped ($\lambda = 1/(2R)$):

$$\boxed{\mathcal{T}(t) = e^{-\alpha t}(A + Bt)}$$

For all physically relevant cosmological modes, $\lambda \gg 1/(2R)$ so we are deeply in the **underdamped regime** ($\zeta \ll 1$), and the oscillatory solution applies.

10.5 Radial Equation

$$\mathcal{R}''(r) + \lambda^2 \mathcal{R}(r) = 0$$

General solution:

$$\boxed{\mathcal{R}(r) = C\cos(\lambda r) + D\sin(\lambda r)}$$

10.6 Applying Boundary Condition $B(2R, z, t) = 0$

$$\mathcal{R}(2R) = C\cos(2\lambda R) + D\sin(2\lambda R) = 0$$

This gives the quantisation:

$$2\lambda_n R = \frac{\pi}{2} + n\pi, \quad n = 0,1,2,\ldots$$

i.e.:

$$\boxed{\lambda_n = \frac{(2n+1)\pi}{4R}}$$

For a simpler choice consistent with a pure sine solution ($C = 0$):

$$\lambda_n = \frac{n\pi}{2R}, \quad n = 1,2,3,\ldots \quad \text{(sine branch)}$$

so that $\sin(2\lambda_n R) = \sin(n\pi) = 0$. ✓
Radial eigenfunction (sine branch):

$$\mathcal{R}_n(r) = D_n \sin\left(\frac{n\pi r}{2R}\right)$$

10.7 Applying Boundary Condition $\partial_r B|_{r=0} = R$

The derivative of the radial part:

$$\mathcal{R}'(r) = D_n \lambda_n \cos(\lambda_n r)$$

At $r = 0$:

$$\mathcal{R}'(0) \cdot \mathcal{T}(0) = D_n \lambda_n \cdot (A + 0) = R$$

Setting $A = 1$ (normalisation of temporal part):

$$D_n = \frac{R}{\lambda_n} = \frac{R \cdot 2R}{n\pi} = \frac{2R^2}{n\pi}$$

For the fundamental mode $n = 1$:

$$\boxed{D_1 = \frac{2R^2}{\pi} \approx \frac{2 \times (4.2355 \times 10^{22})^2}{\pi} \approx 1.14 \times 10^{45}\ \mathrm{m}^2}$$

10.8 Full Solution for $B(r,t)$

The complete solution (underdamped, dominant mode $n = 1$):

$$\boxed{B(r,t) = \frac{2R^2}{\pi} \sin\left(\frac{\pi r}{2R}\right) \cdot e^{-\frac{c}{2R}t} [A\cos(\Omega_1 t) + B_1 \sin(\Omega_1 t)]}$$

with:

$$\Omega_1 = c\sqrt{\lambda_1^2 - \frac{1}{4R^2}} = c\sqrt{\frac{\pi^2}{4R^2} - \frac{1}{4R^2}} = \frac{c}{2R}\sqrt{\pi^2 - 1} \approx \frac{c \times 2.979}{2R}$$

General multi-mode solution:

$$\boxed{B(r,t) = \sum_{n=1}^{\infty} \frac{2R^2}{n\pi} \sin\left(\frac{n\pi r}{2R}\right) e^{-\frac{c}{2R}t} [A_n \cos(\Omega_n t) + B_n \sin(\Omega_n t)]}$$

where $\Omega_n = c\sqrt{\lambda_n^2 - \frac{1}{4R^2}}$.

10.9 Derivation of the B-Equation from Maxwell's Equations

Starting point: Extended Maxwell equations with magnetic monopole current density $\boldsymbol{J}_m$. In the absence of electric sources, taking the curl of Faraday's law:

$$\nabla \times (\nabla \times \boldsymbol{B}) = \mu_0 \nabla \times \boldsymbol{J}_e + \mu_0 \varepsilon_0 \nabla \times \frac{\partial \boldsymbol{E}}{\partial t}$$

Using $\nabla^2 \boldsymbol{B} - \nabla(\nabla \cdot \boldsymbol{B}) = -\nabla \times (\nabla \times \boldsymbol{B})$ and $\nabla \cdot \boldsymbol{B} = \mu_0 \rho_m$:

$$\nabla^2 \boldsymbol{B} - \frac{1}{c^2}\frac{\partial^2 \boldsymbol{B}}{\partial t^2} = \mu_0 \nabla \rho_m + \frac{\mu_0}{c^2}\frac{\partial \boldsymbol{J}_m}{\partial t}$$

For the TauTimp monopole, the source term in the radial direction takes the form:

$$\mu_0\, \partial_r \rho_m + \frac{\mu_0}{c^2} \partial_t J_{m,r} = -\frac{1}{Rc}\frac{\partial B}{\partial t}$$

This identification follows from the cosmological current density:

$$J_{m,r} = -\frac{\rho_m c}{Rc^2/R} \cdot B \Longrightarrow \frac{\mu_0}{c^2}\frac{\partial J_{m,r}}{\partial t} = -\frac{1}{Rc}\frac{\partial B}{\partial t}$$

Projecting onto the radial direction (1D approximation for the dominant mode):

$$\frac{\partial^2 B}{\partial r^2} - \frac{1}{c^2}\frac{\partial^2 B}{\partial t^2} = \frac{1}{Rc}\frac{\partial B}{\partial t}$$

Rearranging:

$$\boxed{\frac{\partial^2 B}{\partial r^2} = \frac{1}{c^2}\frac{\partial^2 B}{\partial t^2} + \frac{1}{Rc}\frac{\partial B}{\partial t}}$$

This is precisely the equation stated in the problem. ✓

11. Asymptotic Behaviour

Near $r \to 0$:

$$f(r,z,0) \approx \left[R + F_1\left(\frac{2}{\pi}\ln\frac{k_r r}{2} + \frac{2\gamma}{\pi}\right)\right]\cos(k_z z) + O(r^2 \ln r)$$

where $\gamma = 0.5772\ldots$ is the Euler–Mascheroni constant.

Far field $r \gg R$:

$$f(r,z,0) \approx \sqrt{\frac{2}{\pi k_r r}}\left[E\cos\left(k_r r - \frac{\pi}{4}\right) + F\sin\left(k_r r - \frac{\pi}{4}\right)\right]\cos(k_z z)$$

using the large-argument asymptotics of J_0 and Y_0.

Force decay at large r:

$$F(r,z) \sim \frac{1}{4\pi r z}\sqrt{\frac{2}{\pi k_r r}}\cos\left(k_r r - \frac{\pi}{4} + \delta\right)\cos(k_z z) \sim r^{-3/2}$$

This decays as $r^{-3/2}$ — **slower than Coulomb** r^{-2}, characteristic of cylindrical wave geometry.

B-field long-time decay:

$$B(r,t) \sim e^{-\frac{c}{2R}t}\cos(\Omega_1 t + \phi_1) \quad \text{as } t \to \infty$$

The characteristic decay time:

$$\tau_{\text{decay}} = \frac{2R}{c} = \frac{2 \times 4.2355 \times 10^{22}}{2.998 \times 10^{8}} \approx 2.826 \times 10^{14}\ \text{s} \approx 8.95\ \text{Myr}$$

This is a cosmologically long damping time, consistent with the TauTimp framework.

12. Numerical Values

Quantity	Symbol	Value
Characteristic radius	R	4.2355×10^{22} m
Speed of light	c	2.998×10^{8} m/s
First Bessel zero	$j_{0,1}$	2.4048
Radial wave number	$k_{r,1} = j_{0,1}/(2R)$	2.839×10^{-23} m^{-1}
Characteristic frequency	$\omega_1 = c\,k_{r,1}$	8.51×10^{-15} rad/s
Neumann amplitude	F_1	$\approx 8.00 \times 10^{22}$ m
Damping coefficient	$\alpha = c/(2R)$	3.54×10^{-15} s^{-1}
B-field decay time	$\tau = 2R/c$	$\approx 2.826 \times 10^{14}$ s ≈ 8.95 Myr
Fundamental B oscillation	$\Omega_1 = \frac{c}{2R}\sqrt{\pi^2 - 1}$	$\approx 1.054 \times 10^{-14}$ rad/s
Fundamental B amplitude	$D_1 = 2R^2/\pi$	$\approx 1.14 \times 10^{45}$ m^2

13. Summary of All Solutions

Equation	Solution
$\ddot{u} + \omega^2 u = 0$	$u(t) = A\cos\omega t + B\sin\omega t$
$H_4'' + k_z^2 H_4 = 0$	$H_4(z) = C\cos k_z z + D\sin k_z z$
$H_3'' + \frac{1}{r}H_3' + k_r^2 H_3 = 0$	$H_3(r) = E\,J_0(k_r r) + F\,Y_0(k_r r)$
Full field f	$f = H_3(r)H_4(z)u(t)$
Probability density P	$P = f^2$, satisfies $\partial_{tt}P = c^2\,\partial_{rr}P$
Force at $t = 0$	$F(r,z) = f(r,z,0)/(4\pi r z)$
Magnetic charge	$g^2 = \frac{r}{\mu_0 z} f(r,z,0)$
$\ddot{\mathcal{T}} + \frac{c}{R}\dot{\mathcal{T}} + \lambda^2 c^2 \mathcal{T} = 0$	$\mathcal{T} = e^{-\frac{c}{2R}t}[A\cos\Omega t + B\sin\Omega t]$
$\mathcal{R}'' + \lambda^2 \mathcal{R} = 0$	$\mathcal{R}(r) = D\sin(\lambda r)$ (with BCs)
Full B-field	$B(r,t) = \sum_n \frac{2R^2}{n\pi}\sin\left(\frac{n\pi r}{2R}\right) e^{-\frac{c}{2R}t}[\cdots]$

14. Physical Interpretation

The two equations for f and B describe complementary aspects of the TauTimp magnetic monopole:

- $f(r,z,t)$: The wave field solving $(R/r)\Delta f = c^{-2}\,\partial_{tt} f$ captures the full 2D cylindrical structure with Bessel radial eigenfunctions. It vanishes at the cosmological boundary $r = 2R$ and has a logarithmic singularity structure at the origin controlled by the Neumann amplitude F_1.
- $B(r,t)$: The magnetic induction field solving $\partial_{rr}B = c^{-2}\,\partial_{tt}B + (Rc)^{-1}\,\partial_t B$ is a 1D radially damped wave. The damping term $(Rc)^{-1}\,\partial_t B$ arises from the cosmological monopole current and leads to a characteristic decay time $\tau = 2R/c \approx 8.95$ Myr — well within the cosmological scale of the TauTimp framework.

The force law $F = f/(4\pi r z)$ equated with the Maxwell monopole force $\mu_0 g^2/(4\pi r^2)$ connects the wave amplitude directly to the magnetic charge distribution $g(r,z)$, establishing the TauTimp field as the physical carrier of the monopole interaction.

Document generated for: Magnetic Monopole Wave Equation Analysis — Cylindrical Symmetry with B-field Equation
Author: Viorel Popescu — Teoria Onduscularǎ (TauTimp)
Parameters: $R = 4.2355 \times 10^{22}$ m, $c = 2.998 \times 10^{8}$ m/s