Quantum physics postulates that the electron in the stationary state doesn't lose energy, and it rotates around the nucleus forever. Is it quite so? Without receiving some energy of any kind?
The energy is coming from the Sun (more generally any star) into quanta of energy within a particle τ neutrino that provides the continuous movement and even the existence of matter itself. It has a bigger energy between the neutrinos and generates movement, gravity, and time in the reality that we experience, and that is non-created light. The energy absorbed by an electron in the hydrogen we consider nonrelativistic Newton's law with "V" the speed of Hydrogen's electron, "ec" the charge, "c" the speed of light, "ε" the electrical permittivity: $m\frac{V^2}{r}=\frac{1}{4\pi\varepsilon}\frac{ec^2}{r^2}\Rightarrow V^2=\frac{ec^2}{4\pi\varepsilon mr}$ (The centripetal force is equal to the electrostatic force). Total energy $E=m\frac{V^2}{2}-\frac{1}{4\pi\varepsilon}\frac{ec^2}{r}=\frac{m}{2}\frac{ec^2}{4\pi\varepsilon mr}-\frac{1}{4\pi\varepsilon}\frac{ec^2}{r}=\frac{1}{8\pi\varepsilon}\frac{ec^2}{r}$ Power radiated, with Larmour's formula. $\frac{dE}{dt}=-\frac{ec^2a^2}{6\pi\varepsilon c^3}=-\frac{ec^2}{6\pi\varepsilon c^3}\left(\frac{V^2}{r}\right)^2=-\frac{ec^6}{96(\pi\varepsilon c)^3m^2r^4}$ Where "a" is centripetal acceleration, $1=-\frac{96(\pi\varepsilon c)^3m^2r^4}{ec^6}\frac{dE}{dt}=-\frac{96(\pi\varepsilon c)^3m^2r^4}{ec^6}\frac{d}{dt}\left(-\frac{1}{8\pi\varepsilon}\frac{ec^2}{r}\right)=-3c^3\left(\frac{4\pi\varepsilon mr}{ec^2}\right)^2\frac{dr}{dt}$ and integrating

$$T_0=\int_0^{t0}dt=-3c^3\left(\frac{4\pi\varepsilon m}{ec^2}\right)^2\int_{t0}^{0}r^2\frac{dr}{dt}dt=-3c^3\left(\frac{4\pi\varepsilon m}{ec^2}\right)^2\int_{r0}^{0}r^2dr$$

$T_0=\left(\frac{4\pi\varepsilon m}{ec^2}\right)^2\cdot(rb\cdot c)^3$ as in https://michaelvio.github.io/Tint.pdf
Thus, the Hydrogen's electron must receive energy every $T_0=6.22\cdot10^{-11}$ sec, so the Hydrogen's electron won't fall on the nucleus. There are 25 quanta of (tau) τ neutrinos' energy that fully charge a Hydrogen's electron with energy per/s when the hole τ's kinetic energy, about 1.975 eV, is transferred to the electron. Hydrogen's electron won't fall on the nucleus. *Every $6{,}22\cdot10^{-11}$ sec, the electron charge is changing, forming a dipole and rearranging charge in the atom, thus forming a pair of π^+ and π^- exchanging charge between the electron and the proton.*

But probably the reaction $e+\lceil\tau\xrightarrow{trans}e+\tau\rceil+\pi^++\pi^-+\mu\rceil+\Delta E$ (1) with Eτ=rest mas 0.1889eV/c^2 and kinetic energy of 1.975eV is more complex in the equation supplying energy generated by the graviton $e+\lceil\nu\xrightarrow{trans}e+\Delta E_1$ (2) With a rest mass of 0.00038eV/c^2 & Kinetic energy of about 1.6295eV. In the nucleus, we have a τ neutrino that will collide with a pair of a proton and neutron, generating the process of continuous migration of protons, generating gravity, and the μ neutrino:
$p+\lceil\nu\xrightarrow{transf}n+posit\rceil\xrightarrow{decay}(p+e+\nu\rceil)+posit=p+(e+posit)=p$ and equation:

$$p+\lceil\tau\xrightarrow{transf}n+\nu\rceil+\pi^++\mu\rceil\xrightarrow{decay}(p+e+\nu\rceil)+\mu\rceil+\lceil\pi^-+\pi^+=p+(e+\pi^+)=p+\mu+\Delta E_2$$

I guess that they receive quanta of energy for every predetermined period of around ~0.62 picosecond for a Hydrogen atom at room temperature, 1-S orbital, there is a reaction in an electron collides with a τ neutrino as in the equation below where monopole is meson, the lightest pion π^+, π^- with a lower rest mass of 0.007eV/c^2. And the τ neutrino will collide with a pair of protons and neutrons, generating the process of continuous migration of protons, generating gravity, and the μ neutrino that generates the hologram of space:

$$p+\lceil\tau\xrightarrow{transf}n+\nu\rceil+\pi^++\mu\rceil\xrightarrow{decay}(p+e+\nu\rceil)+\mu\rceil+\lceil\pi^-+\pi^+=p+(e+\pi^+)+\Delta E \qquad (2)$$

in the Hydrogen nucleus. All the fundamental particles: proton, electron, neutron, photon, magnetron, $Termal_{Particle}$, and phonon interact with τ neutrino and receive a constant amount of energy and temporal ionization. https://www.youtube.com/watch?v=frg9rJvFoA0 The 16' minute

Correct Aspects: The classical force balance ($mv^2/r = k\,e^2/r^2$) and Larmor integration are standard for illustrating the classical instability. The ~25 collisions for energy transfer is a rough estimate but aligns with the calculated timescale. *Incorrect Aspects:* The document treats the hydrogen electron as non-relativistic (valid approximation, v << c) but ignores that relativity isn't the issue—quantum mechanics is. Relativistic corrections (e.g., fine structure) don't resolve the classical radiation problem. *Overall Correctness*: Math is approximately right for the classical case, but irrelevant to real atomic stability. 8 Nov. 2025