# Theory of Information (Quarks)

There are 6-informational quarks that mean: Mass (M), Magnetic field ($M_g$), Electric charge (E), Gravity (G), Light (L) & Time (T). A particle always consists of three informational quarks.
Two particles may interact if bought are in the boundary (neighboring) for at list a Quanta of time.
The informational quarks correspond with the known quarks as follows:
Light = Up Quark; Magnetic field = Down Quark
Gravity = Charm Quark; Electric charge = Strange Quark
Time = Top Quark; Mass = Bottom Quark; As you can see Gravity and Electric charge have a similar potential equation (Charm & Strange) as Light and Magnetic field (Up & Down); Time and Mass thus (Top & Bottom) have τ and μ neutrino as the interaction fundamental particle and the same onduscular equation with a similar solution.

| Property: Quark: / Particle: | Mass Bottom | Mag.Field Down | ElectCharge Strange | Gravity Charm | Light Up | Time Top |
|---|---|---|---|---|---|---|
| Electron | 1 | 1838 | -1 | 0 | 0 | 0 |
| Anti-electron (positron) | 1 | -1838 | 1 | 0 | 0 | 0 |
| Neutron | 1838 | -1.913 | 0 | -1 | 0 | 0 |
| Proton | 1836 | 2.793 | 1 | 0 | 0 | 0 |
| $Monopole_{mR}$ ($\pi^-$ , $\pi^+$) | 0 | (+/-)2.793 | 0 | 0 | 1 | 1 |
| $Monopole_{Mw}$ ($\pi^-$ , $\pi^+$) | 0 | (+/-)2.793 | 0 | 1 | 0 | 1 |
| Graviton = ν Electronic Neutrino | 0 | -1838 | 0 | 1 | 1 | 0 |
| Particle $_{\gamma}$ (Gama-ray) | 0 | 0 | 0 | 1 | 1 | -1 |
| Particle $_{Ex}$ | 0 | 0 | 1 | 1 | 0 | -1 |
| Photon (Light) | 0 | 1 | 0 | 0 | 1 | 1 |
| μ Neutrino bound with Higgs Boson | 0 | 0 | 0 | -1 | 1 | 1 |
| Tau τ Neutrino (Time) | 0 | 0 | 0 | 1 | 1 | 1 |
| Entanglement-Torsion | 0 | -1.913 | 0 | 0 | 1 | -1 |
| Particle S (Syncron) | 0 | 0 | 0 | 1 | -1 | -1 |
| Activate | 0 | 0 | 0 | -1 | 1 | -1 |
| Audion | 0 | 0 | 0 | -1 | 1 | 1 |
| Particle $_{M}$ (Mass resonance) | 1 | 0 | 1 | 0 | 1 | 0 |
| Optoelectron | 1 | 0 | -1 | 0 | 1 | 0 |
| Anti-Neutron | 1838 | -1.913 | 0 | -1 | 0 | 0 |
| Thermal Particle | 1 | 2.793 | 0 | 0 | 1 | 0 |
| Virtual Particle $_{En}$ (Physical Entropy) | 0 | 1 | 0 | 1 | 0 | 1 |

Not yet produced or detected. Particle $_{Mx}$; Particle $_{E}$; Particle $_{En}$ (Physical Entropy Virtual particle); Part $_{S;}$
The meaning of a negative magnetic field is that the magnetic field is opposite to the spin.
The mass is always positive in the local universe.
The meaning of positive time is that the propagation speed is the light speed ($3\cdot10^8$ m/s) and the negative time means that the transfer is ~$3\cdot10^{18}$m/s. The communications with γ-ray are suitable for long-distance between planets in several Time Qanta. For details see:
https://michaelvio.github.io/T&SQ.pdf
The meaning of positive gravity is attraction and negative gravity is rejection.
The meaning of positive light is its emission and negative light is its absorption.
The wave function of the graviton for the stationary state is:

https://michaelvio.github.io/grav.pdf
https://michaelvio.github.io/Sun.pdf

Where _C1 is the normed constant R space constant ($3.9228\cdot10^{22}$ meter) and in spherical coordinates for _C1=1; R=49 we have the following plot for r=0 to 2R, for stationary state $t_0$=0. As you can see the Newtonian force decreases parabolically. This is the Newton law explicit, annulated in 2R with sign-changing. It goes to infinity when r->-∞ (repulsion see the plot below) and is extremely high but not ∞ in origin because of the granularity of space-time.

Thus, the force between two nucleons that change ν neutrino equation (21):

$$F_{g1} \sim \frac{3R^2}{14 \cdot \pi} \cdot \frac{1}{r^2} - \frac{R}{4 \cdot \pi} \cdot \frac{1}{r} + \frac{1}{14 \cdot \pi}$$

As in the equation into FluxG.mw calculus.

Thus, the quantic gravitational forces in Newtons between 2 nucleons at distance r that change ν neutrino (gravitons) are:

$$\mathrm{F_{g1}} = \frac{4.09 \cdot 10^{-64}}{r^2} - 1.34 \cdot \frac{10^{-86}}{r} + 1.07 \cdot 10^{-109} \sim \frac{4.09 \cdot 10^{-64}}{r^2}$$

Thus, the quantic gravitational attraction force in Newton between 2 nucleons at radius r (in meters) may be approximate: $Fgq \sim 2 \div 4 \cdot 10^{-64} / r^2$ (Newton). According to the flux conservation equation.

21 Oct. 2025