

# CSC418 Computer Graphics Assignment 1

---

Michael Vu 1002473272

## 1. Curves

---

### 1.1 Convert Parametric to Implicit

$$\begin{aligned}x(t)^2 &= 4 \sin^2(t) \\y(t)^2 &= 25 \sin^2(t) \cos^2(t) \\y(t)^2 &= \frac{25}{16} x^2 (4 - x^2) \\0 &= y^2 + \frac{25}{16} x^4 - \frac{25}{4} x^2\end{aligned}$$

### 1.2 Tangent Vector

$$\begin{aligned}\vec{T}(t) &= \frac{\langle \frac{dx}{dt}, \frac{dy}{dt} \rangle}{\|\langle \frac{dx}{dt}, \frac{dy}{dt} \rangle\|} \\ \frac{dx}{dt} &= 2 \cos(t) \\ \frac{dy}{dt} &= 10 \cos(2t) \\ \langle \frac{dx}{dt}, \frac{dy}{dt} \rangle &= 2 \langle \cos(t), 5 \cos(2t) \rangle \\ \vec{T}(t) &= \frac{\langle \cos(t), 5 \cos(2t) \rangle}{\sqrt{\cos^2(t) + 25 \cos^2(2t)}}\end{aligned}$$

### 1.3 Normal Vector

$$\begin{aligned}\vec{N}(t) &= \frac{\frac{d\vec{T}}{dt}}{\|\frac{d\vec{T}}{dt}\|} \\ \vec{N}(t) &= \frac{\langle -\sin(t), -10 \sin(2t) \rangle}{\|\frac{d\vec{T}}{dt}\|} \\ &= \frac{\langle -\sin(t), -10 \sin(2t) \rangle}{\sqrt{\sin^2(t) + 100 \sin^2(2t)}}\end{aligned}$$

### 1.4 Symmetry

#### 1.4.1 X-axis Symmetry

If the function is symmetrical about the X-axis, then  $f(x, y) = f(x, -y)$ .

$$\begin{aligned}
 f(x, y) &= f(x, -y) \\
 y^2 + \frac{25}{16}x^4 - \frac{25}{4}x^2 &= (-y)^2 + \frac{25}{16}x^4 - \frac{25}{4}x^2 \\
 y^2 &= (-y)^2 \\
 \text{left side} &= \text{right side}
 \end{aligned}$$

Therefore the function is symmetric about the X-axis.

### 1.4.2 Y-axis Symmetry

If the function is symmetrical about the Y-axis, then  $f(x, y) = f(-x, y)$ .

$$\begin{aligned}
 f(x, y) &= f(-x, y) \\
 y^2 + \frac{25}{16}x^4 - \frac{25}{4}x^2 &= y^2 + \frac{25}{16}(-x)^2 - \frac{25}{4}(-x)^2 \\
 \frac{25}{16}x^4 - \frac{25}{4}x^2 &= \frac{25}{16}(-x)^4 - \frac{25}{4}(-x)^2 \\
 \text{left side} &= \text{right side}
 \end{aligned}$$

Therefore the function is symmetric about the Y-axis.

## 1.5 Area

Since the function is symmetric about the X and Y axis, it is sufficient to find the area of the function in one quadrant and multiply by 4.

$$\begin{aligned}
 f(x) &= \sqrt{\frac{25}{4}x^2 - \frac{25}{16}x^4} = \frac{5}{4}x\sqrt{4 - x^2} \\
 \frac{1}{4}A &= \frac{5}{4} \int_0^2 x\sqrt{4 - x^2} dx \\
 \text{let } j &= 4 - x^2 \\
 dx &= \frac{1}{-2x} dj \\
 A &= \frac{5}{2} \int_0^4 \sqrt{j} dj \\
 A &= \frac{5}{2} \cdot \frac{16}{3} = \frac{40}{3}
 \end{aligned}$$

Therefore the area under the bowtie is  $\frac{40}{3}$ .

## 1.6 Perimeter

TODO in polar coordinates

## 2. Transformations

### 2.a Translation and Translation

Two sequential  $N \times N$  translations have the form

$$\begin{aligned}
T_1 &= \begin{bmatrix} 1 & 0 & \dots & t_{x1} \\ 0 & 1 & \dots & t_{y1} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} \\
T_2 &= \begin{bmatrix} 1 & 0 & \dots & t_{x2} \\ 0 & 1 & \dots & t_{y2} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} \\
T_1 T_2 &= \begin{bmatrix} 1 & 0 & \dots & t_{x1} \\ 0 & 1 & \dots & t_{y1} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & \dots & t_{x2} \\ 0 & 1 & \dots & t_{y2} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & \dots & t_{x1} + t_{x2} \\ 0 & 1 & \dots & t_{y1} + t_{y2} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix}
\end{aligned}$$

By inspection, the summation in the rightmost column does not change value based on the order of  $t_{i1}$  and  $t_{i2}$ . Changing the order of  $T_1$  and  $T_2$  does not affect the resulting transformation matrix, therefore **translation and translation is commutative**.

## 2.b Translation and Rotation

Assume that translations and rotations are commutative. The form of an  $\mathbb{R}^2$  rotation in homogenous form is

$$T = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

It follows that a translation followed by a rotation has the form

$$T_1 = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta & t_x \cos \theta - t_y \sin \theta \\ \sin \theta & \cos \theta & t_x \sin \theta + t_y \cos \theta \\ 0 & 0 & 1 \end{bmatrix}$$

Conversely, a rotation followed by a rotation has the form

$$T_2 = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta & t_x \\ \sin \theta & \cos \theta & t_y \\ 0 & 0 & 1 \end{bmatrix}$$

$T_1 \neq T_2$  which is a contradiction, therefore **translation and rotation are not commutative**.

## 2.c Scaling and Rotation, with different fixed points

Assume towards a contradiction that scaling and rotation with differing fixed points is commutative.

Consider the following transformations:

$$T_{scale} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & -1 \\ 0 & 0 & 1 \end{bmatrix} \text{ (fixed point at (0,1))}$$

$$T_{rotate} = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ (90 degrees, fixed point at (0,0))}$$

If scaling and rotation are commutative, then in this example:

$$T_{scale}T_{rotate} = T_{rotate}T_{scale}$$

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & -1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & -1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 0 & -2 & 0 \\ 2 & 0 & -1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & -2 & 1 \\ 2 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Which is a contradiction, therefore **scaling and rotation around different fixed points is not commutative.**

## 2.d Scaling and Scaling, with the same fixed point.

A scaling transformation about a fixed point  $\mathbf{p}$  can be expressed as a series of transformations:

1. Translate such that  $\mathbf{p}$  is at the new origin
2. Scale (denoted  $\mathbf{S}$ ) about the origin
3. Translate the inverse of 1.

Let Two scaling transformations (denoted  $\mathbf{T}_a, \mathbf{T}_b$ ) about the same fixed point has the form

$$\begin{aligned} T_a T_b &= (T_1 S_a T_3)(T_1 S_b T_3) \\ &= T_1 S_a T_3 T_3^{-1} S_b T_3 \\ &= T_1 (S_a S_b) T_3 \end{aligned}$$

And reversing the order of  $\mathbf{a}, \mathbf{b}$  will yield a similar result

$$\begin{aligned} T_b T_a &= (T_1 S_b T_3)(T_1 S_a T_3) \\ &= T_1 S_b T_3 T_3^{-1} S_a T_3 \\ &= T_1 (S_b S_a) T_3 \end{aligned}$$

We then observe that

$$T_a T_b = T_b T_a \iff S_a S_b = S_b S_a$$

The implication here is that since both  $\mathbf{T}_a$  and  $\mathbf{T}_b$  share the same  $\mathbf{T}_1$  and  $\mathbf{T}_3$ , the overall transformation will be commutative if arbitrary origin-centred scale transformations  $\mathbf{S}_a$  and  $\mathbf{S}_b$  are also commutative.

Two  $\mathbf{N} \times \mathbf{N}$  origin-scale transformations will take the following form

$$T = \begin{bmatrix} S_{x1} & 0 & \dots & 0 \\ 0 & S_{y1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} \begin{bmatrix} S_{x2} & 0 & \dots & 0 \\ 0 & S_{y2} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} = \begin{bmatrix} S_{x1}S_{x2} & 0 & \dots & 0 \\ 0 & S_{y1}S_{y2} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix}$$

which is clearly commutative. Therefore, since scaling about the origin is commutative, **scaling transformations with the same fixed point are commutative.**

## 3. Homography

---

### 3.1 Affine Transform Derivation

$$\begin{bmatrix} (1,0) \\ (0,1) \\ (1,1) \\ (0,0) \end{bmatrix} \Rightarrow \begin{bmatrix} (6,2) \\ (7,3) \\ (6,3) \\ (7,2) \end{bmatrix}$$

A trace of the transformation reveals that it is a x-scaling by a factor of  $-1$  about the origin, and a translation by  $\langle 7, 2 \rangle$ . The combined transformation matrix is

$$T = \begin{bmatrix} 1 & 0 & 7 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 0 & 7 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$$

### 3.2 Map (2,5) Under the Transformation

$$\vec{p} = \begin{bmatrix} -1 & 0 & 7 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 5 \\ 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \\ 1 \end{bmatrix} \rightarrow (5, 7)$$

The point  $(2, 5)$  maps to  $(5, 7)$  under this transformation.

## 4. Polygons

---

TODO