

A New Paradigm in X-ray Spectral Analysis

Deconvolving Spectra using Machine Learning

Carter Rhea Ph.D Candidate,¹ Jutie Hlavacek-Larrondo¹,
Akos Bogdan², & Ralph Kraft²


May 17, 2022

¹ L'Université de Montréal

² Harvard Smithsonian Center for Astrophysics

Fonds de recherche
Nature et
technologies

Québec









IVADO

Université
de Montréal

CRAQ
Comprendre l'Univers
Understanding the Universe




X-ray Astronomy




Observations vs Reality

The Source's Intrinsic Spectrum



What we observe




Why do we need this?


1. Initial Parameter Estimation for Traditional Fitting Methods

LevMar

Fit Space




Monte Carlo




Why do we need this?

1. Initial Parameter Estimation for Traditional Fitting Methods
2. Stacking of X-ray Data


Individual Spectra



Combine Spectra



Stacked Spectra



Outline of Presentation


- Introduction
 - Formulation of observed spectrum
 - Effect of Response Matrix in Observations
 - Previously Attempted Solutions
- Recurrent Inference Machine
 - Introduction
 - Basic Example
 - In the Context of X-ray Spectra
- Next Steps

Introduction



Potential Solutions for the Response Matrix

E = Photon Energy Space
 E' = Detector Energy Space

$$S_{obs}(E) = \int_0^\infty R(E', E) S_{true}(E) dE$$



=




Potential Solutions for the Response Matrix



i = Photon Energy Space

j = Detector Energy Space


$$S_{obs_i} = \sum_{ij} R_{ij} S_{true_j}$$



=



Effects of Response Files



Solving the Matrix Equation

$$R S_{true} = S_{obs}$$

Pseudo-Inverse

Tikhonov Regularization

DOES NOT WORK

$$S_{true} = R^\dagger S_{obs}$$

$$\|R S_{true} - S_{obs}\|_{L_2} - \mu \|D\|_{L_2}$$


$$S_{true} = [R R^T + \mu I]^\dagger [R^T S_{obs}]$$

Recurrent Inference Machine

Recurrent Inference Machines

How does a Recurrent Inference Machine work:

Solve the linear equation $Ax=b$ iteratively by using an RNN to update solution



Putzky & Welling 2017
arxiv.org/abs/1706.04008


Morningstar et al. 2018
arxiv.org/abs/1808.00011

Morningstar et al. 2019
arxiv.org/pdf/1901.01359.pdf



A quick reminder about our problem

i = Photon Energy Space
j = Detector Energy Space


$$S_{obs_i} = \sum_{ij} R_{ij} S_{true_j}$$



=



Application to X-ray Spectra



What can we do with this?

- Train a convolutional neural network to estimate the underlying parameters (Rhea et al. in prep.)
- Explore the transiency of X-ray sources (Rhea et al. in prep.)
- Study the calibration of the Chandra X-ray Observatory (Prunier & Rhea et al. in prep.)
- Investigate Metallicity in the outskirts of galaxies

A New Paradigm in X-ray Spectral Analysis

Deconvolving Spectra using Machine Learning

Carter Rhea ^{Ph.D Candidate, 1}, Julie

Hlavacek-Larrondo¹, Akos Bogdan², & Ralph Kraft²


May 17, 2022

¹ L'Université de Montréal

² Harvard Smithsonian Center for Astrophysics

Fonds de recherche
Nature et
technologies

Québec




IVADO

Université
de Montréal



CRAQ
Comprendre l'Univers
Understanding the Universe



Next Step: Estimate Temperature and Metallicity



Estimate Temperature and Metallicity



Putting it all together!



Preprocessing (RIM)


Deconvolve X-ray Spectra s/t we can stack them

Mixture Density Network

Train MDN on deconvolved spectra to obtain posterior distributions of temperature and metallicity

Bayesian Inference (BXA)

Use MDN results as priors to do a full Bayesian inference approach to X-ray Spectral Fitting




Rhea, C. et al, AJ 160, 5 (2020)

Buchner, J. et al, A&A 564, A125 (2014)

Next Steps

1. Converge RIM on larger dataset
 - a. Redo CNN analysis on larger dataset
2. Apply to real data (Self-Similarity)
 - a. Reduce Chandra Data
 - b. Use RIM to deconvolve
 - c. Apply CNN to estimate parameters
3. Potential other ideas?

Galaxy Cluster X-ray Luminosity-Temperature Relation



A New Paradigm in X-ray Spectral Analysis

Deconvolving Spectra using Machine Learning

Carter Rhea, Julie Hlavacek-Larrondo, Akos Bogdan, & Ralph Kraft
April XX, 2022

