

ooooo  
oooo

ooooooo



Eidgenössische Technische Hochschule Zürich  
Swiss Federal Institute of Technology Zurich



**Beauty**



**Truth**

## The Discovery of the Heavy Quarks

### Experimental Foundations of Particle Physics

**Michael Reichmann**

20th November 2017



# Table of contents

1 Introduction

2 The Beauty Quark

3 The Truth Quark


4 Conclusion

## Section 1

### Introduction

# Introduction

- in 1974 (with J/ $\psi$ ) 4 leptons and 4 quarks discovered



## Introduction


- in 1974 (with  $J/\psi$ ) 4 leptons and 4 quarks discovered
  - in 1973 6-plet proposed by Kobayashi and Maskawa to describe CP violation
  - in 1975 Perl et al. discovered  $\tau$ -lepton and its neutrino
  - induction of another pair of quarks

## Standard Model

	I	II
mass	$\approx 2.4 \text{ MeV}/c^2$	$\approx 1.275 \text{ GeV}/c^2$
charge	2/3	2/3
spin	1/2	1/2
quarks	up  down 	charm  strange 
leptons	electron  tau neutrino 	muon  tau neutrino 
	$\approx 4.8 \text{ MeV}/c^2$ -1/3 1/2	$\approx 95 \text{ MeV}/c^2$ -1/3 1/2
	$\approx 0.511 \text{ MeV}/c^2$ -1 1/2	$\approx 105.67 \text{ MeV}/c^2$ -1 1/2
	electron 	muon 
	$<2.2 \text{ eV}/c^2$ 0 1/2	$<1.7 \text{ MeV}/c^2$ 0 1/2
	electron neutrino 	muon neutrino 
		$<15.5 \text{ MeV}/c^2$ 0 1/2
		tau neutrino 

# Introduction

- in 1974 (with J/ $\psi$ ) 4 leptons and 4 quarks discovered
- in 1973 6-plet proposed by Kobayashi and Maskawa to describe CP violation
- in 1975 Perl et al. discovered  $\tau$ -lepton and its neutrino
- indiction of another pair of quarks
- in 1977 discovery of the bottom (beauty) quark
- postulation of a sixth quark



# Introduction

- in 1974 (with J/ $\psi$ ) 4 leptons and 4 quarks discovered
- in 1973 6-plet proposed by Kobayashi and Maskawa to describe CP violation
- in 1975 Perl et al. discovered  $\tau$ -lepton and its neutrino
- indiction of another pair of quarks
- in 1977 discovery of the bottom (beauty) quark
- postulation of a sixth quark
- in 1995 discovery of the top (truth) quark
- complete set of fermions until now

Standard Model			
three generations of matter (fermions)			
	I	II	
mass	$\approx 2.4 \text{ MeV}/c^2$	$\approx 1.275 \text{ GeV}/c^2$	$\approx 172.44 \text{ GeV}/c^2$
charge	2/3	2/3	2/3
spin	1/2	1/2	1/2
	u	c	t
	up	charm	top
mass	$\approx 4.8 \text{ MeV}/c^2$	$\approx 95 \text{ MeV}/c^2$	$\approx 4.18 \text{ GeV}/c^2$
charge	-1/3	-1/3	-1/3
spin	1/2	1/2	1/2
	d	s	b
	down	strange	bottom
mass	$\approx 0.511 \text{ MeV}/c^2$	$\approx 105.67 \text{ MeV}/c^2$	$\approx 1.7768 \text{ GeV}/c^2$
charge	-1	-1	-1
spin	1/2	1/2	1/2
	e	$\mu$	$\tau$
	electron	muon	tau
mass	$< 2.2 \text{ eV}/c^2$	$< 1.7 \text{ MeV}/c^2$	$< 15.5 \text{ MeV}/c^2$
charge	0	0	0
spin	1/2	1/2	1/2
	$\nu_e$	$\nu_\mu$	$\nu_\tau$
	electron neutrino	muon neutrino	tau neutrino

LEPTONS




## Section 2


### The Beauty Quark

# The $\Upsilon$ -Meson


- bound state of  $b\bar{b}$
- decay channels:



(a) 82 %



(b) 9 %



(c) 2 %

- mostly decay into gluons which hadronise  $\rightarrow$  signals mostly caused by hadrons
- leptonic decay splits up into 2.5 % to 3 % for each  $e$ ,  $\mu$  and  $\tau$

# Discovery Paper

## Observation of a Dimuon Resonance at 9.5 GeV in 400-GeV Proton-Nucleus Collisions

S. W. Herb, D. C. Hom, L. M. Lederman, J. C. Sens,<sup>(a)</sup> H. D. Snyder, and J. K. Yoh  
*Columbia University, New York, New York 10027*

and

J. A. Appel, B. C. Brown, C. N. Brown, W. R. Innes, K. Ueno, and T. Yamanouchi  
*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*


and

A. S. Ito, H. Jöstlein, D. M. Kaplan, and R. D. Kephart  
*State University of New York at Stony Brook, Stony Brook, New York 11974*  
 (Received 1 July 1977)

Accepted without review at the request of Edwin L. Goldwasser under policy announced 26 April 1976

Dimuon production is studied in 400-GeV proton-nucleus collisions. A strong enhancement is observed at 9.5 GeV mass in a sample of 9000 dimuon events with a mass  $m_{\mu^+\mu^-} > 5$  GeV.


We have observed a **strong enhancement at 9.5 GeV** in the mass spectrum of dimuons produced in **400-GeV proton-nucleus collisions**. Our conclusions are based upon an analysis of 9000 dimuon events with a reconstructed mass  $m_{\mu^+\mu^-}$  greater than 5 GeV corresponding to  $1.6 \times 10^{16}$  protons incident on Cu and Pt targets:




The produced muons are analyzed in a **double-arm magnetic-spectrometer system** with a mass resolution  $\Delta m/m$  (rms)  $\approx 2\%$ .

The experimental configuration (Fig. 1) is a modification of an earlier dilepton experiment in the Fermilab Proton Center Laboratory.<sup>1-3</sup> Narrow targets ( $\sim 0.7$  mm) with lengths corresponding to 30% of an interaction length are employed.

# Setup




## Setup




- 400 GeV proton beam shot on narrow target (Pt/Cu) with 30% interaction length

# Setup




- hadron filter out of Be with 18 interaction length ( $3^\circ$  to  $5^\circ$  horiz. and  $\pm 0.5^\circ$  vert.)

# Setup




- heavy metal (Steel, W) shielding to minimise particle leakage

# Setup




- tungsten beam dump

# Setup




- additional shielding out of polyethylene and more steel

# Setup




- spectrometer dipole magnets with horizontal field
- both arms are symmetric to drawing plane and detect  $\mu^+$  and  $\mu^-$

# Setup




- scintillation hodoscopes and wire chambers for tracking (limit of  $10 \times 10^7$  counts/s)

# Setup




- solid iron magnet to partially refocus and redetermine muon momentum

# Setup




- Čerenkov counter to prevent low momentum muon triggers

## Intermezzo: Sideband Fit




- typical shape of data in particle physics: continuous background with a small bump

## Intermezzo: Sideband Fit




- background extraction with a fit of the whole set does not work well

## Intermezzo: Sideband Fit




- divide the data set in a signal and two background parts ( $\rightarrow$  side bands)

## Intermezzo: Sideband Fit




- fitting just the side bands yields a much better result

## Intermezzo: Sideband Fit



- getting the signal distribution by subtracting the full data set by the fit

## Intermezzo: Sideband Fit



- get width and position of the signal by a fit

# Results

- statistically significant enhancement at 9.5 GeV  $\mu^+\mu^-$  mass
- solid line background fit using side band method


$$\frac{d^2\sigma}{dm dy} = Ae^{-bm}$$

- fit expects 350 events in excluded region but 770 events in data
- bump wider than resolution of detector (FWHM: 0.5(1) GeV)
- simple Gaussian fit of background subtracted data yields:

$$m = 9.54(4) \text{ GeV}$$

$$\text{FWHM} = 1.16(9) \text{ GeV}$$


- same goodness of fit with two Gaussians with fixed width of detector resolution  $\rightarrow$  later  $\Upsilon$  and  $\Upsilon'$



**Figure:** dimuon production cross section as a function of invariant mass

## Verification

- same sign dimuon spectrum ( $\mu^+ \mu^+$  and  $\mu^- \mu^-$ )
  - ▶ upper limit on combined effects of accidental coincidences and hadronic decays
  
- re-measurement of the muon momentum by the second magnet and by PCW at the centre of the first magnet
  - ▶ avoid misidentified  $\psi \rightarrow \mu^+ + \mu^-$  at high mass
  - ▶ confirmed by clear separation of the  $\psi$  and  $\psi'$  in the Figure
  
- study of various subsets of the data with different magnetic fields
  - ▶ check for apparatus bias
  
- study of data with and without target
  - ▶ rule out signal created from beam dump





# PLUTO Collaboration Paper

## OBSERVATION OF A NARROW RESONANCE FORMED IN $e^+e^-$ ANNIHILATION AT 9.46 GeV

PLUTO Collaboration

Ch. BERGER, W. LACKAS, F. RAUPACH, W. WAGNER

*I. Physikalisches Institut der RWTH Aachen, FRG*

G. ALEXANDER<sup>1</sup>, L. CRIEGEE, H.C. DEHNE, K. DERIKUM, R. DEVENISH, G. FLÜGGE, G. FRANKE

Ch. GERKE, E. HACKMACK, P. HARMS, G. HORLITZ, Th. KAHL<sup>2</sup>, G. KNIES, E. LEHMANN,

B. NEUMANN, R.L. THOMPSON<sup>3</sup>, U. TIMM, P. WALOSCHEK, G.G. WINTER,

S. WOLFF, W. ZIMMERMANN

*Deutsches Elektronen-Synchrotron DESY, Hamburg, FRG*

O. ACHTERBERG, V. BLOBEL, L. BOESTEN, H. DAUMANN, A.F. GARFINKEL<sup>4</sup>, H. KAPITZA,


B. KOPPITZ, W. LÜHRSEN, R. MASCHUW, H. SPITZER, R. van STAA, G. WETJEN

*II. Institut für Experimentalphysik der Universität Hamburg, FRG*

Received 9 May 1978


An experiment using the PLUTO detector has observed the formation of a narrow, high mass, resonance in  $e^+e^-$  annihilations at the DORIS storage ring. The mass is determined to be  $9.46 \pm 0.01$  GeV which is consistent with that of the Upsilon. The gaussian width  $\sigma$  is observed as  $8 \pm 1$  MeV and is equal to the DORIS energy resolution. This suggests that the resonance is a bound state of a new heavy quark-antiquark pair. An electronic width  $\Gamma_{ee} = 1.3 \pm 0.4$  keV was obtained. In standard theoretical models, this favors a quark charge assignment of  $-1/3$ .

# PLUTO Detector



## Further Experiments


## PLUTO Detector



- up to 5 GeV electron beams collided in the centre

## Further Experiments


## PLUTO Detector



- cylindrical proportional wire chamber with 92 % coverage
- tracking and momentum measurement

## Further Experiments


## PLUTO Detector



- cylindrical array of shower counters with 8.6 radiationlength and 94 % coverage
- calorimeter to measure the full energy


## Further Experiments

## PLUTO Detector



- magnet with 1.69 T

# PLUTO Detector




- muon chambers



## Results

- acquiring cross section by scanning  $\sqrt{s}$  in steps of either 5 GeV or 10 GeV
- only considering hadronic decay products
- remove  $1/s$  background
- reduce cosmic ray background by use of bunch crossing time (bunched beam structure vs continuous background)
- remove QED events with coplanarity cut and shower recognition
- remove beam gas interaction by cuts on visible energy and missing mass
- fitting data with Gaussian yields:



**Figure:** total cross section for hadron production

$$\mathbf{m = 9.46(1) \text{ GeV}}$$

$$\sigma_{\text{Gauss}} = 7.8(9) \text{ MeV}$$



## Further Experiments

## Charge

- relation of total hadronic cross section to resonance mass and the resonance width

$$\int \sigma_h dM = \frac{6\pi^2}{M_R^2} \frac{\Gamma_{ee}\Gamma_h}{\Gamma_{tot}}$$

- standard assumption:  $\Gamma_{tot} \approx \Gamma_h \rightarrow$  direct measurement of  $\Gamma_{ee}$

$$\Gamma_{ee} = 1.3(4) \text{ keV}$$

- theoretical predictions from the same model:

Decay Mode	Width [keV]	
	$e = \pm 1/3$	$e = \pm 2/3$
$\mu^+ \mu^-$	0.70(9)	2.8(4)

- implies charge of  $-1/3$  for the bottom quark

## Section 3

### The Truth Quark

## Introduction

- in early 90s all SM particles except H and t-quark discovered
- beauty discovery → weak isospin partner was undoubted
- quark masses are fundamental parameters in the SM
- early estimates:  $m_t \approx 3m_b \approx 15 \text{ GeV}$
- many new accelerator could only push limits higher:
- TRISTAN ( $e^+e^-$ ) at KEK (Tsukuba, Japan) with  $\sqrt{s} = 61.4 \text{ GeV} \rightarrow 30.2 \text{ GeV}$
- Sp $\bar{p}$ S at CERN with  $\sqrt{s} = 630 \text{ GeV} \rightarrow 69 \text{ GeV}$
- SLC ( $e^+e^-$ ) at Stanford and LEP ( $e^+e^-$ ) at CERN  $\rightarrow 1/2 m_t$
- hadron collider needed ( $\rightarrow$  Tevatron)




## Decay Channels (1)

- estimate on  $m_t$  in 1994:  $\sim 180 \text{ GeV}$
- prior to discovery: behaviour completely predicted by SM
- $m_t > m_W \rightarrow$  main decay channel ( $\sim 95\%$ ):  $t \rightarrow W^+ b$

$$\Gamma_t = \frac{G_F m_t^3}{8\pi\sqrt{2}} \left(1 - \frac{m_W^2}{m_t^2}\right)^2 \left(1 + 2\frac{m_W^2}{m_t^2}\right)$$


- width for the expected mass:  $\sim 1 \text{ GeV} \rightarrow$  decay before hadronisation



- leptonic decay equally splits up into  $e$ ,  $\mu$  and  $\tau$

## Decay Channels (2)

- top mostly pair produced via  $q\bar{q} \rightarrow t\bar{t}$  or gluon fusion:  $gg \rightarrow t\bar{t}$
- main decay of the top pair:  $t\bar{t} \rightarrow W^+ b W^- \bar{b}$
- b decay:




- typical signals:
  - ▶ 2 b-jets + dilepton ( $e^+e^-$ ,  $\mu^+\mu^-$ ,  $e^+\mu^-$ ,  $\mu^+e^-$ )
  - ▶ 2 b-jets + single lepton + two jets
  - ▶ 2 b-jets + 4 jets
- huge background on pure QCD process due to other more common QCD processes
- how to discriminate b-jets from other jets?

# B-Tagging

- how to discriminate b-jets from other jets?

## t̄t Cross Section



- cross section extracted from SM
- Tevatron Lumi in 1995:  $10 \times 10^{30} \text{ cm}^{-2} \text{ s}^{-1}$

$$R_{t\bar{t}} = \sigma_{t\bar{t}} \mathcal{L} = 0.1 \text{ Hz}$$

# Discovery Paper (1)

VOLUME 74, NUMBER 14

PHYSICAL REVIEW LETTERS

3 APRIL 1995

## Observation of Top Quark Production in $p\bar{p}$ Collisions with the Collider Detector at Fermilab

- F. Abe,<sup>14</sup> H. Akimoto,<sup>32</sup> A. Akoopman,<sup>27</sup> M. G. Albrow,<sup>5</sup> S. R. Amendolia,<sup>24</sup> D. Amidei,<sup>17</sup> J. Antos,<sup>29</sup> C. Anway-Wiese,<sup>4</sup> S. Aota,<sup>32</sup> G. Apollinari,<sup>27</sup> T. Asakawa,<sup>27</sup> W. Ashmanakis,<sup>13</sup> M. Atac,<sup>1</sup> P. Auchincloss,<sup>38</sup> F. Azfar,<sup>27</sup> P. Azz-Bacchetta,<sup>27</sup> N. Bacchetta,<sup>27</sup> W. Badgett,<sup>17</sup> S. Bagdasarov,<sup>27</sup> M. W. Bailey,<sup>19</sup> J. Bao,<sup>29</sup> P. de Barbaro,<sup>26</sup> A. Barbour,<sup>1</sup> J. Baur,<sup>1</sup> J. Baumgartner,<sup>1</sup> J. Beaman,<sup>34</sup> J. Belcher,<sup>1</sup> J. Benekos,<sup>1</sup> J. Bensinger,<sup>1</sup> D. Bernbach,<sup>14</sup> S. Beltramini,<sup>1</sup> G. Beltrami,<sup>24</sup> J. Bellinger,<sup>14</sup> S. Benoit,<sup>1</sup> J. Bentwich,<sup>1</sup> J. Bensinger,<sup>1</sup> D. Bernbach,<sup>14</sup> A. Beretvas,<sup>1</sup> J. P. Berge,<sup>1</sup> S. Bertlucci,<sup>1</sup> A. Bhatt,<sup>27</sup> K. Biery,<sup>27</sup> M. Binkley,<sup>1</sup> D. Bisello,<sup>27</sup> R. E. Blair,<sup>1</sup> C. Blocker,<sup>1</sup> A. Bodok,<sup>20</sup> W. Bokhari,<sup>19</sup> V. Bolognesi,<sup>24</sup> D. Bornstein,<sup>27</sup> J. Boureau,<sup>23</sup> G. Brindenburg,<sup>1</sup> L. Breccia,<sup>1</sup> C. Bremberg,<sup>1</sup> E. Buckley-Geer,<sup>1</sup> H. S. Budd,<sup>28</sup> K. Burkett,<sup>17</sup> J. Busetto,<sup>21</sup> A. Byon-Wagner,<sup>1</sup> K. L. Byrum,<sup>1</sup> J. Cammermeyer,<sup>3</sup> C. Campagnari,<sup>1</sup> M. Campbell,<sup>17</sup> A. Carter,<sup>1</sup> W. Cartiers,<sup>13</sup> D. Carlsmith,<sup>34</sup> A. Castro,<sup>21</sup> G. Cataz,<sup>24</sup> Y. Chen,<sup>27</sup> F. Cervelli,<sup>24</sup> H. Y. Chou,<sup>1</sup> J. Chapman,<sup>17</sup> M.-T. Cheng,<sup>27</sup> S. Chiarelli,<sup>34</sup> T. Chikamatsu,<sup>27</sup> C. N. Chiu,<sup>17</sup> P. Chrostek,<sup>18</sup> J. Chubang,<sup>1</sup> A.-G. Choi,<sup>27</sup> D. Cobal,<sup>24</sup> M. Contreras,<sup>2</sup> J. Conway,<sup>17</sup> J. Cooper,<sup>1</sup> M. Corcoran,<sup>1</sup> C. Courtney,<sup>1</sup> D. Currat,<sup>1</sup> J. Curran,<sup>1</sup> J. Cumming,<sup>1</sup> T. Daniels,<sup>16</sup> J. Dolough,<sup>1</sup> D. Dschung,<sup>1</sup> J. Donahue,<sup>1</sup> P. A. Dujardin,<sup>1</sup> L. Dutour,<sup>1</sup> M. Dzhaparidze,<sup>1</sup> P. Derwent,<sup>17</sup> T. Devlin,<sup>24</sup> M. Dickson,<sup>20</sup> J. R. Dittmann,<sup>8</sup> S. Donati,<sup>24</sup> R. B. Drucker,<sup>15</sup> A. Dunn,<sup>17</sup> N. Eddy,<sup>17</sup> K. Einwaller,<sup>17</sup> J. E. Elias,<sup>17</sup> R. Ely,<sup>17</sup> E. Engels,<sup>17</sup> D. Erede,<sup>17</sup> S. Erede,<sup>17</sup> Q. Fan,<sup>17</sup> L. Fiori,<sup>1</sup> B. Flaugher,<sup>1</sup> G. W. Foster,<sup>1</sup> M. Franklin,<sup>1</sup> M. Frantzis,<sup>17</sup> J. Freeman,<sup>17</sup> J. Friedman,<sup>19</sup> H. Frisch,<sup>1</sup> T. A. Fuess,<sup>1</sup> Y. Fukui,<sup>14</sup> S. Funki,<sup>32</sup> G. Gagliardi,<sup>23</sup> S. Galeotti,<sup>24</sup> M. Gallinaro,<sup>17</sup> M. Garcia-Sivvers,<sup>18</sup> A. G. Garfinkel,<sup>17</sup> C. Gay,<sup>17</sup> S. Geer,<sup>27</sup> D. W. Gerdes,<sup>17</sup> P. Giannetti,<sup>24</sup> N. Giokaris,<sup>27</sup> P. Giromini,<sup>17</sup> L. Gladney,<sup>17</sup> D. Glenzinski,<sup>13</sup> M. Gold,<sup>19</sup> J. Gonzalez,<sup>22</sup> A. Gorini,<sup>17</sup> A. T. Gorini,<sup>17</sup> P. Gosselain,<sup>17</sup> B. Handt,<sup>24</sup> R. M. Hans,<sup>17</sup> S. Han,<sup>17</sup> B. Han,<sup>17</sup> R. M. Harris,<sup>7</sup> S. A. Hauser,<sup>1</sup> J. Hauer,<sup>1</sup> C. Hawk,<sup>28</sup> E. Hayashi,<sup>12</sup> J. Heinrich,<sup>27</sup> M. Hohmann,<sup>13</sup> C. Holck,<sup>27</sup> R. Holtbeck,<sup>27</sup> L. Holloway,<sup>11</sup> A. Hölzler,<sup>32</sup> S. Hong,<sup>17</sup> G. Hu,<sup>27</sup> B. T. Huffman,<sup>27</sup> R. Hughes,<sup>27</sup> J. Huston,<sup>17</sup> J. Hurt,<sup>9</sup> J. Hylen,<sup>17</sup> H. Iieda,<sup>27</sup> M. Incagli,<sup>24</sup> J. Incandela,<sup>17</sup> J. Iwas,<sup>17</sup> Y. Iwata,<sup>19</sup> H. Jensen,<sup>17</sup> U. Joshi,<sup>1</sup> R. W. Kadel,<sup>15</sup> E. Kajase,<sup>17</sup> T. Kamon,<sup>17</sup> T. Kaneko,<sup>17</sup> K. Karr,<sup>17</sup> H. Kasba,<sup>17</sup> Y. Kata,<sup>20</sup> L. Keeble,<sup>8</sup> K. Kelley,<sup>17</sup> R. D. Kennedy,<sup>28</sup> R. Kephart,<sup>7</sup> T. Kesten,<sup>17</sup> D. Kestenbaum,<sup>9</sup> B. J. Kent,<sup>17</sup> F. Keutelian,<sup>17</sup> P. Keywan,<sup>17</sup> B. J. Kim,<sup>17</sup> D. H. Kim,<sup>7,27</sup> H. S. Kim,<sup>17</sup> S. B. Kim,<sup>17</sup> S. H. Kim,<sup>17</sup> Y. K. Kim,<sup>17</sup> J. Kirsch,<sup>17</sup> S. Koide,<sup>17</sup> J. Konigstberg,<sup>17</sup> S. Kopp,<sup>3</sup> K. Kuroda,<sup>17</sup> T. Kuroda,<sup>17</sup> S. Kuroda,<sup>17</sup> T. Kuroda,<sup>17</sup> T. Kuroda,<sup>17</sup> T. Kuroda,<sup>17</sup> T. Kuroda,<sup>17</sup> T. Kuroda,<sup>17</sup> T. Kuroda,<sup>17</sup> S. E. Kuhlmann,<sup>17</sup> E. Kunc,<sup>17</sup> A. J. Laasanen,<sup>17</sup> N. Labrecque,<sup>17</sup> J. Lai,<sup>17</sup> S. Lammel,<sup>17</sup> J. J. Lamoureux,<sup>17</sup> T. LeCompte,<sup>17</sup> S. Leone,<sup>24</sup> J. D. Lewis,<sup>17</sup> P. Limon,<sup>17</sup> M. Lindgren,<sup>17</sup> T. M. Liu,<sup>17</sup> N. Lockyer,<sup>17</sup> O. Long,<sup>22</sup> C. Loos,<sup>17</sup> M. Loos,<sup>17</sup> J. Lu,<sup>17</sup> D. Luccio,<sup>27</sup> A. Menzione,<sup>24</sup> S. Meschi,<sup>24</sup> S. Metzler,<sup>17</sup> C. Mian,<sup>17</sup> G. Micalilli,<sup>17</sup> S. Mikamo,<sup>17</sup> R. Miller,<sup>18</sup> H. Minato,<sup>27</sup> S. Missetti,<sup>3</sup> M. Mishima,<sup>14</sup> T. Mizushima,<sup>32</sup> T. Miyamoto,<sup>32</sup> S. Miyashita,<sup>32</sup> Y. Morita,<sup>17</sup> J. Mueller,<sup>27</sup> S. Mukherjee,<sup>17</sup> T. Murakami,<sup>17</sup> P. Murli,<sup>17</sup> H. Nakada,<sup>17</sup> M. Nakamura,<sup>17</sup> T. Nakazono,<sup>17</sup> T. Neuberg,<sup>17</sup> S. Nakanishi,<sup>17</sup> M. Ninomiya,<sup>27</sup> L. Nozaki,<sup>17</sup> S. Okada,<sup>17</sup> S. Okada,<sup>17</sup> K. E. Ohr,<sup>17</sup> T. Ohsawa,<sup>17</sup> R. Ohura,<sup>17</sup> S. Okubo,<sup>17</sup> S. Okuno,<sup>17</sup> R. Olivier,<sup>17</sup> J. Olsen,<sup>17</sup> C. Pagliarone,<sup>17</sup> R. Pauletta,<sup>17</sup> V. Papadimitriou,<sup>17</sup> S. Pappas,<sup>17</sup> S. Pappas,<sup>17</sup> S. Park,<sup>17</sup> J. Patrick,<sup>17</sup> G. Paulieta,<sup>24</sup> M. Paulini,<sup>17</sup> L. Pescina,<sup>21</sup> M. D. Peter,<sup>17</sup> T. J. Phillips,<sup>5</sup> G. Piacentino,<sup>2</sup> M. Pillai,<sup>26</sup> K. T. Pitts,<sup>17</sup> R. Plunkett,<sup>17</sup> L. Pondrom,<sup>17</sup> J. Proudfit,<sup>1</sup> F. Probos,<sup>17</sup> G. Purzy,<sup>17</sup> K. Ragan,<sup>17</sup> A. Ribon,<sup>27</sup> F. Rimondi,<sup>17</sup> L. Ristori,<sup>24</sup> W. J. Robertson,<sup>17</sup> T. Rodriguez,<sup>17</sup> J. Romano,<sup>17</sup> R. Roser,<sup>17</sup> W. K. Sakamoto,<sup>26</sup> D. Saltzberg,<sup>3</sup> A. Samson,<sup>8</sup> L. Sant,<sup>24</sup> H. Sato,<sup>32</sup> V. Scarpine,<sup>17</sup> R. E. Schmidt,<sup>17</sup> M. P. Schmidt,<sup>17</sup> G. F. Sciaccia,<sup>24</sup> A. Scribano,<sup>24</sup> S. Seglar,<sup>17</sup> S. Sheng,<sup>17</sup> Y. Miyata,<sup>17</sup> S. Saito,<sup>17</sup> S. Saito,<sup>17</sup> S. Saito,<sup>17</sup> S. K. Siwa,<sup>27</sup> P. Slepnev,<sup>17</sup> D. A. Smith,<sup>17</sup> F. D. Sandler,<sup>17</sup> J. Senter,<sup>17</sup> J. Spalding,<sup>17</sup> P. Sphicas,<sup>17</sup> L. Spiegel,<sup>17</sup> A. Spies,<sup>17</sup> L. Stanev,<sup>17</sup> J. Staelo,<sup>34</sup> A. Stefanini,<sup>24</sup> K. Stahl,<sup>17</sup> J. Strait,<sup>17</sup> D. Stuar,<sup>17</sup> G. Sullivan,<sup>17</sup> A. Szumanski,<sup>27</sup> K. Sunрок,<sup>16</sup> J. Suzuki,<sup>32</sup> T. Takada,<sup>32</sup> T. Takahashi,<sup>27</sup> T. Takami,<sup>17</sup> K. Takayama,<sup>17</sup> N. Tamura,<sup>33</sup> F. Tartarelli,<sup>27</sup> W. Taylor,<sup>17</sup> P. K. Teng,<sup>29</sup> Y. Teranomoto,<sup>20</sup> S. Tether,<sup>16</sup> D. Theriot,<sup>17</sup> L. T. Thomas,<sup>17</sup> R. Thun,<sup>17</sup> M. Timko,<sup>17</sup> P. Tipton,<sup>17</sup> A. Titov,<sup>27</sup> S. Tkaczyn,<sup>17</sup> J. Toback,<sup>20</sup> K. Tolleson,<sup>20</sup> A. Tolleson,<sup>20</sup> J. Tomison,<sup>27</sup> J. F. de Trocino,<sup>17</sup> S. Trout,<sup>17</sup> J. Tseng,<sup>17</sup> N. Turini,<sup>24</sup> T. Uchida,<sup>32</sup> N. Uemura,<sup>32</sup> K. Uegawa,<sup>27</sup> G. Umlat,<sup>17</sup> S. C. van der Brink,<sup>27</sup> S. Veijari,<sup>17</sup> G. Velez,<sup>24</sup> R. Vital,<sup>17</sup> M. Vonruck,<sup>17</sup> D. Vacinic,<sup>17</sup> R. G. Wagner,<sup>17</sup> R. L. Wagner,<sup>17</sup> J. Wahl,<sup>17</sup> R. C. Walker,<sup>26</sup> C. Wang,<sup>7</sup> C. H. Williams,<sup>17</sup> G. Williams,<sup>17</sup> W. Williams,<sup>17</sup> M. Williams,<sup>17</sup> Q. F. Williams,<sup>17</sup> W. Williams,<sup>17</sup> W. Williams,<sup>17</sup> R. Webb,<sup>17</sup> C. Wei,<sup>17</sup> C. Wendt,<sup>17</sup> W. C. Wester,<sup>17</sup> A. B. Wicklund,<sup>17</sup> E. Wicklund,<sup>17</sup> R. Wilkinson,<sup>17</sup> H. H. Williams,<sup>17</sup> P. Wilson,<sup>17</sup> B. L. Winer,<sup>17</sup> D. Wolinski,<sup>17</sup> J. Wolinski,<sup>30</sup> X. Wu,<sup>24</sup> J. Wyss,<sup>21</sup> A. Yagil,<sup>17</sup> W. Yao,<sup>15</sup>

## Discovery Paper (2)

K. Yasuoka,<sup>32</sup> Y. Ye,<sup>12</sup> G. P. Yeh,<sup>7</sup> P. Yeh,<sup>29</sup> M. Yin,<sup>6</sup> J. Yoh,<sup>7</sup> C. Yosef,<sup>18</sup> T. Yoshida,<sup>20</sup> D. Yovanovitch,<sup>7</sup> I. Yu,<sup>35</sup>  
 J. C. Yun,<sup>7</sup> A. Zanetti,<sup>24</sup> F. Zetti,<sup>24</sup> L. Zhang,<sup>34</sup> W. Zhang,<sup>22</sup> and S. Zucchelli<sup>2</sup>

(CDF Collaboration)

<sup>1</sup>Argonne National Laboratory, Argonne, Illinois 60439

<sup>2</sup>Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40126 Bologna, Italy

<sup>3</sup>Brandeis University, Waltham, Massachusetts 02254

<sup>4</sup>University of California at Los Angeles, Los Angeles, California 90024

<sup>5</sup>University of Chicago, Chicago, Illinois 60637

<sup>6</sup>Duke University, Durham, North Carolina 27708

<sup>7</sup>Fermi National Accelerator Laboratory, Batavia, Illinois 60510

<sup>8</sup>Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy

<sup>9</sup>Harvard University, Cambridge, Massachusetts 02138

<sup>10</sup>Hiroshima University, Higashi-Hiroshima 724, Japan

<sup>11</sup>University of Illinois, Urbana, Illinois 61801

<sup>12</sup>Institute of Particle Physics, McGill University, Montreal, Canada H3A 2T8

and University of Toronto, Toronto, Canada M5S 1A7

<sup>13</sup>The Johns Hopkins University, Baltimore, Maryland 21218

<sup>14</sup>National Laboratory for High Energy Physics (KEK), Tsukuba, Ibaraki 305, Japan

<sup>15</sup>Lawrence Berkeley Laboratory, Berkeley, California 94720

<sup>16</sup>Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

<sup>17</sup>University of Michigan, Ann Arbor, Michigan 48109

<sup>18</sup>Michigan State University, East Lansing, Michigan 48824

<sup>19</sup>University of New Mexico, Albuquerque, New Mexico 87131

<sup>20</sup>Osaka City University, Osaka 588, Japan

<sup>21</sup>Università di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy

<sup>22</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104

<sup>23</sup>University of Pittsburgh, Pittsburgh, Pennsylvania 15260

<sup>24</sup>Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy

<sup>25</sup>Purdue University, West Lafayette, Indiana 47907

<sup>26</sup>University of Rochester, Rochester, New York 14627

<sup>27</sup>Rockefeller University, New York, New York 10021

<sup>28</sup>Rutgers University, Piscataway, New Jersey 08854

<sup>29</sup>Academia Sinica, Taipei, Taiwan 11529, Republic of China

<sup>30</sup>Texas A&M University, College Station, Texas 77843

<sup>31</sup>Texas Tech University, Lubbock, Texas 79409

<sup>32</sup>University of Tsukuba, Tsukuba, Ibaraki 305, Japan


<sup>33</sup>Tufts University, Medford, Massachusetts 02155

<sup>34</sup>University of Wisconsin, Madison, Wisconsin 53706

<sup>35</sup>Yale University, New Haven, Connecticut 06511



(Received 24 February 1995)

We establish the existence of the top quark using a  $67 \text{ pb}^{-1}$  data sample of  $\bar{p}p$  collisions at  $\sqrt{s} = 1.8 \text{ TeV}$  collected with the Collider Detector at Fermilab (CDF). Employing techniques similar to those we previously published, we observe a signal consistent with  $t\bar{t}$  decay to  $WWb\bar{b}$ , but inconsistent with the background prediction by  $4.8\sigma$ . Additional evidence for the top quark is provided by a peak in the reconstructed mass distribution. We measure the top quark mass to be  $176 \pm 8(\text{stat}) \pm 10(\text{syst}) \text{ GeV}/c^2$  and the  $t\bar{t}$  production cross section to be  $6.8^{+1.4}_{-1.2} \text{ pb}$ .



## Discovery at CDF

## CDF Detector



ooooo  
oooo

oooo●oo


## Introduction

- paper from 1994 with estimate on mass and cross section
- using dataset of  $19 \text{ pb}^{-1} + 47 \text{ pb}^{-1}$  ( $\rightarrow \sim 400$  events)
- looking at two decay channels
  - ▶ dilepton
  - ▶ lepton + jets
- both data samples subsets of events with isolated leptons with high  $P_T > 20 \text{ GeV}$
- cut on invariant mass of dilepton  $75 \text{ GeV} < m_l < 105 \text{ GeV} \rightarrow$  exclude Z events
- main background reduction by b-tagging
  - ▶ reconstruction of secondary vertices from b decay in SVX  $\rightarrow$  SVX tag
  - ▶ finding additional leptons from b decay in ECAL  $\rightarrow$  SLT tag

## Lepton + Jets Channel


### SVX tagging:

- search for secondary vertices with three or more tracks
- then search for two or more tracks with more stringent track and vertex quality
- efficiency estimated by  $e, \mu$  samples with enriched b decays (96 % agreement to MC)
- tagging efficiency: 42(5) %
- backgrounds from recoil of heavy quark pairs against W and mistags
- for  $W+ \geq 3$  jets: observation of 27 tags with bg of 6.7(21) tags
- decay lifetime of SVX tags agrees well with MC



# Dilepton Channel

- major backgrounds:
  - ▶ Drell-Yan process
  - ▶  $Z \rightarrow \tau\tau$
  - ▶ misidentified hadrons
  - ▶  $WW, b\bar{b}$



## Section 4

### Conclusion

# The Truth Quark

- empty
- moreempty
- moremoreempty