

Prediction of amyloidogenicity based on the n-gram analysis

¹University of Wrocław, Department of Genomics, ²Wrocław University of Technology, Department of Mathematics and ³Wrocław University of Technology, Department of Biomedical Engineering, Faculty of Fundamental Problems of Technology

Associate Editor: XXXXXXXX

Abstract

Supplementary information: Supplementary data are available at *Bioinformatics* online.

1

| Category | Property |
|--------------------|---|
| Contactivity | Average flexibility indices (Bhaskaran and Ponnuswamy, 1988) |
| Contactivity | 14 Å contact number (Nishikawa and Ooi, 1986) |
| Contactivity | Accessible surface area (Radzicka and Wolfenden, 1988) |
| Contactivity | Buriability (Zhou and Zhou, 2004) |
| Contactivity | Values of Wc in proteins from class Beta, cutoff 12 Å, separation 5 (Wozniak and Kotulska, 2014) |
| Contactivity | Values of Wc in proteins from class Beta, cutoff 12 Å, separation 15 (Wozniak and Kotulska, 2014) |
| β -frequency | Average relative probability of inner beta-sheet (Kanehisa and Tsong, 1980) |
| β -frequency | Relative frequency in beta-sheet (Prabhakaran, 1990) |
| β -frequency | Thermodynamic beta sheet propensity (Kim and Berg, 1993) |
| Hydrophobicity | Hydrophobicity index (Argos <i>et al.</i> , 1982) |
| Hydrophobicity | Optimal matching hydrophobicity (Sweet and Eisenberg, 1983) |
| Hydrophobicity | Hydrophobicity-related index (Kidera <i>et al.</i> , 1985) |
| Hydrophobicity | Scaled side chain hydrophobicity values (Black and Mould, 1991) |
| Polarity | Polarizability parameter (Charton and Charton, 1982) |
| Polarity | Mean polarity (Radzicka and Wolfenden, 1988) |
| Size | Average volumes of residues (Pontius <i>et al.</i> , 1996) |
| Stability | Side-chain contribution to protein stability (kJ/mol) (Takano and Yutani, 2001) |

Table 1. Physicochemical properties used in the creation of reduced amino acid alphabets.




Fig. 1. The scheme of n-gram extraction. A) Input data - peptides with a known amyloidogenicity status. B) Each peptide sequence was divided into overlapping hexamers. The amyloidogenicity status of a source sequence was used as the amyloidogenicity status of a derived hexamer. C) From each hexamer we extracted continuous 1-, 2- and 3-grams. We selected also gapped 2-grams with the length of gap equal from 1 to 3 residues and gapped 3-grams with a single gap between the first and the second or the second and the third element of the n-gram.

a short part of the sequence is responsible for amyloidogenicity, our method might results in many false positives in the training data set. To evade this problem, we restricted the maximum length of peptides in training data set to fifteen amino acids to easy the extraction of probable hot-spots.

From each hexamer we extracted n-grams of the following length: 1, 2 and 3. In the case of 2- and 3-grams, we separately counted both gapped and continous n-grams. For 2-grams we counted n-grams with gaps of lengths from 1 to 3 and for 3-grams with a single gap between the first and the second or the second and the third element (see Fig. 1).

The study the length of amyloidogenicity signal, we trained three classifiers for each encoding on the sequences of different length. We considered separately six-residue sequences, shorter of equal to ten residues and shorter or equal to fifteen residues.

All n-grams extracted from the hexamers in the training data set were filtered using QuiPT, our own implementation of permutation test with the

information gain (mutual information) as the criterion of the importance of a specific n-gram. In the next step, we used n-grams with the p-value smaller than 0.05 to build a random forest (Breiman, 2001) classifier using ranger **R** package (Wright and Ziegler, 2015).

Furthermore, we repeated the procedure described above on two typical reduced alphabets of amino acids derived from the literature to check if the process of amyloidogenicity does require nonstandard groupings of amino acids. We also added the full amino acid alphabet to assess the advantages of the amino acid encoding.

3.4 Cross-validation and selection of the best-performing encoding

The ability to correctly predict amyloidogenicity was assessed during the five-fold cross-validation. Since the data set was very heterogeneous, we

repeated the cross-validation fifteen times for each classifier to obtain more precise estimates of performance measures for each classifier.

To evaluate if our classifiers are able to use decision rules extracted from sequences of given length to correctly classify longer or shorter sequences, we calculate performance measures separately for four ranges of lengths of sequences: 6, 7-10, 11-15 and 16-25. The number of sequences from the given length range was roughly comparable between folds of cross-validation.

To choose the most adequate reduced amino acid alphabet, we ranked the values of AUC (with rank 1 for the best AUC, rank 2 for the second best AUC and so on) for each range of sequence length in the testing data set. The encoding with the lowest sum of ranks from all sequence length categories was selected as the best one. For this encoding, we choose the range of peptides length in the training set providing the best AUC in cross-validation.

3.5 Benchmark of AmyloGram

The best-performing reduced amino acid alphabet chosen during the cross-validation was later used to train AmyloGram, n-gram based predictor of amyloidogenicity.

We used *pep424* data set (Walsh *et al.*, 2014) to compare the performance of AmyloGram and other predictors of amyloidogenicity. Since the model of AmyloGram does not covers peptides shorter than five amino acids, we removed them from the total benchmark data set. It should not affect the comparison as only five sequences were eliminated (around 1% of the original data set). Additionally, we also benchmarked three predictors learned on the n-grams extracted from sequences of different length ranges without any amino acid encoding.

All benchmarked classifiers were trained on sequences used during the cross-validation. Since some peptides were common for both *pep424* and AmyLoad, we removed them from the training data set. After purification, the learning data set had 269 positive sequences and 746 negative sequences longer than five residues and shorter than fifteen residues. Aside from the preparation of the training data, we exactly repeated the procedure of n-gram extraction as described above (see Fig. 1).

3.6 Encoding distance

The encoding distance is defined as the minimum number of amino acids that have to be moved between subgroups of encoding to make a identical to b (the order of subgroups in the encoding and amino acids in a group is unimportant). This measure may be further normalized by a factor reflecting how much moving amino acids between groups altered mean group properties.

4 Results and discussion

4.1 Selection of the best-performing encoding

The selected encoding performed better than other reduced alphabets considering all sequence length ranges in training and testing data set. It has the AUC always in the fourth quartile (Fig. 2). For the best-performing encoding the most problematic was correct prediction of the amyloidogenicity in the longest peptides 16-25 not when learned on very short sequences, but longer ones (6-10 and 6-15). Such behavior was typical for most of the analyzed encodings.

The highest values of AUC the chosen encoding reaches in the relatively the simplest cases of predicting amyloidogenicity of the shortest sequences (6 residues). Of course, the decision rules were the easiest to extract when also the learning set has only sequences of the same lengths, but results were quite comparable even if the training set contained longer sequences. Also in this situation, majority of reduced amino acid alphabets

behaved like the selected encoding and also have higher AUC than in other scenarios.

Furthermore, the best encoding has also very good sensitivity and specificity regardless of the length of sequences in training and testing set (Fig. 3). The classifiers trained on the peptides of length 6 tend to have the best specificity, while predictors learned on the long sequences have the best sensitivity. As the matter of fact, training on the sequences ranging from six to ten residues seem to yield most balanced classifiers with optimal sensitivity and specificity.

In addition to reduced alphabets, we also evaluated classifiers based on the full amino acid alphabet. In most cases they also places in the fourth quartile considering their AUC value (Fig. 2). Nevertheless, they never predict amyloidogenicity better than the selected encoding. It suggests that despite the predictive power of n-grams based on full alphabet, the encoding allows better generalization of the prediction rules and in the consequence better performance.

Similarly to the best-performing encoding, the sensitivity of classifiers based on the full amino acid alphabet seems to decrease with the length of sequences in the training data set (Fig. 3). Furthermore, these classifiers seem to always have one of the worst sensitivities among all analyzed predictors, especially when tested on longer amyloids. It means that using the full amino acid alphabet it easier to point the non-amyloidogenic sequences instead of recognizing amyloids.

The encodings found in the literature performs substantively worse than other analyzed amino acid alphabets in all categories. It indicates that classical divisions of amino acids do not create groups suitable for the recognition of amyloids. This observation is well-supported by the specificity-sensitivity plot (Fig. 3), where classifiers trained with this encodings of amino acids groups with the worst performers.

4.2 The best-performing encoding and important n-grams

The reduced amino acid alphabet chosen during the analysis has six elements (Tab. 2). Identical reduced amino acid alphabets were created through ten other combinations of physicochemical properties. The following features appeared in all eleven combinations: hydrophobicity index (Argos *et al.*, 1982), average flexibility indices (Bhaskaran and Ponnuswamy, 1988), polarizability parameter (Charton and Charton, 1982) and thermodynamic beta sheet propensity (Kim and Berg, 1993).

Two groups, 3 and 4 contain mostly hydrophobic amino acids, while 1 and 4 are mostly hydrophilic. In addition to this, group 1 and 4 have amino acid with the highest average flexibility. The least flexible amino acids are in group 5. The most polarizable amino acids belong to group 4, while the lowest polarizability is typical for glycine in group 1. The glycine has also the highest thermodynamic beta sheet propensity, which has the lowest values for groups 3 and 4.

Table 2. The best-performing encoding.

| ID | Groups |
|----|------------------|
| 1 | G |
| 2 | K, P, R |
| 3 | I, L, V |
| 4 | F, W, Y |
| 5 | A, C, H, M |
| 6 | D, E, N, Q, S, T |

We discovered that 65 n-grams have p-values smaller than 0.05 in the every fold of repetitions of cross-validation regardless of the training set.




Fig. 2. Distribution of AUC values of different reduced amino acid alphabets. Red circle: classifier employing best encoding of amino acid. Green square: classifier using full amino acid alphabet. Blue square: classifiers employing encodings from literature.

4.3 Benchmark of AmyloGram

The benchmark covered AmyloGram as well as two best-performing peer-reviewed predictors of amyloidogenicity (PASTA2 (Walsh *et al.*, 2014) and FoldAmyloid (Garbuzynskiy *et al.*, 2010)). We analyzed Area Under the Curve (AUC), Matthew’s Correlation Coefficient (MCC), Sensitivity and Specificity (see Tab. 3).

Interestingly, the n-gram extraction method was efficient enough to produce classifiers able to outperformed published methods. Two of three n-gram based classifiers trained on the full alphabet have AUC higher than PASTA2 and all three were more successful than FoldAmyloid. They also maintained they high Specificity as seen previously during cross-validation.

Although the proposed n-gram extraction creates efficient classifiers, the encoding of amino acids further increases the prediction of amyloidogenicity. AmyloGram has the highest AUC, MCC and Sensitivity among all tested classifiers. Is has lower specificity than two classifiers trained on the full alphabet, but still outperforms other published method in this category. It is important to highlight that AmyloGram is the most balanced of analyzed classifiers, having the best Specificity/Sensitivity balance, as indicated by the value of MCC.

Table 3. Results of benchmark on pep424 data set for AmyloGram, PASTA2, FoldAmyloid and random forest predictor learned on n-grams extracted without any amino acid encoding from the sequences of the length specified in the brackets.

| Classifier | AUC | MCC | Sensitivity | Specificity |
|----------------------|---------------|---------------|---------------|---------------|
| AmyloGram | 0.8972 | 0.6307 | 0.8658 | 0.7889 |
| PASTA2 | 0.8550 | 0.5227 | 0.7987 | 0.7444 |
| FoldAmyloid | 0.7351 | 0.4526 | 0.7517 | 0.7185 |
| full alphabet (6) | 0.8411 | 0.5427 | 0.4966 | 0.9593 |
| full alphabet (6-10) | 0.8581 | 0.5698 | 0.7517 | 0.8259 |
| full alphabet (6-15) | 0.8610 | 0.5490 | 0.8188 | 0.7519 |

5 Conclusion

Acknowledgments

Text Text Text Text Text Text Text Text.

Funding

This research was partially funded by the KNOW Consortium.





Fig. 3. Sensitivity and specificity of classifiers in cross-validation. Red circle: classifier employing best encoding of amino acid. Green square: classifier using full amino acid alphabet. Blue square: classifiers employing encodings from literature. The classifier based on the best encoding always have good specificity and sensitivity.

References

- Argos, P., Rao, J. K., and Hargrave, P. A. (1982). Structural prediction of membrane-bound proteins. *European journal of biochemistry / FEBS*, **128**(2-3), 565–575.
- Bhaskaran, R. and Ponnuswamy, P. (1988). Positional flexibilities of amino acid residues in globular proteins. *International Journal of Peptide and Protein Research*, **32**(4), 241–255.
- Black, S. D. and Mould, D. R. (1991). Development of hydrophobicity parameters to analyze proteins which bear post- or cotranslational modifications. *Analytical Biochemistry*, **193**(1), 72–82.
- Breiman, L. (2001). Random Forests. *Machine Learning*, **45**(1), 5–32.
- Charton, M. and Charton, B. I. (1982). The structural dependence of amino acid hydrophobicity parameters. *Journal of Theoretical Biology*, **99**(4), 629–644.
- Garbuzynskiy, S. O., Lobanov, M. Y., and Galzitskaya, O. V. (2010). FoldAmyloid: a method of prediction of amyloidogenic regions from protein sequence. *Bioinformatics (Oxford, England)*, **26**(3), 326–332.
- Jr, J. H. W. (1963). Hierarchical Grouping to Optimize an Objective Function. *Journal of the American Statistical Association*, **58**(301), 236–244.
- Kanehisa, M. I. and Tsong, T. Y. (1980). Local hydrophobicity stabilizes secondary structures in proteins. *Biopolymers*, **19**(9), 1617–1628.
- Kawashima, S., Pokarowski, P., Pokarowska, M., Kolinski, A., Katayama, T., and Kanehisa, M. (2008). AAindex: amino acid index database, progress report 2008. *Nucleic Acids Research*, **36**(suppl 1), D202–D205.
- Kidera, A., Konishi, Y., Oka, M., Ooi, T., and Scheraga, H. A. (1985). Statistical analysis of the physical properties of the 20 naturally occurring amino acids. *Journal of Protein Chemistry*, **4**(1), 23–55.
- Kim, C. A. and Berg, J. M. (1993). Thermodynamic beta-sheet propensities measured using a zinc-finger host peptide. *Nature*, **362**(6417), 267–270.
- Nishikawa, K. and Ooi, T. (1986). Radial locations of amino acid residues in a globular protein: correlation with the sequence. *Journal of Biochemistry*, **100**(4), 1043–1047.
- Pontius, J., Richelle, J., and Wodak, S. J. (1996). Deviations from Standard Atomic Volumes as a Quality Measure for Protein Crystal Structures. *Journal of Molecular Biology*, **264**(1), 121–136.
- Prabhakaran, M. (1990). The distribution of physical, chemical and conformational properties in signal and nascent peptides. *The Biochemical Journal*, **269**(3), 691–696.
- Radzicka, A. and Wolfenden, R. (1988). Comparing the polarities of the amino acids: side-chain distribution coefficients between the vapor phase, cyclohexane, 1-octanol, and neutral aqueous solution. *Biochemistry*, **27**(5), 1664–1670.
- Sweet, R. M. and Eisenberg, D. (1983). Correlation of sequence hydrophobicities measures similarity in three-dimensional protein structure. *Journal of Molecular Biology*, **171**(4), 479–488.
- Takano, K. and Yutani, K. (2001). A new scale for side-chain contribution to protein stability based on the empirical stability analysis of mutant proteins. *Protein Engineering*, **14**(8), 525–528.
- Walsh, I., Seno, F., Tosatto, S. C. E., and Trovato, A. (2014). PASTA 2.0: an improved server for protein aggregation prediction. *Nucleic Acids Research*, page gku399.
- Wozniak, P. P. and Kotulska, M. (2014). Characteristics of protein residue-residue contacts and their application in contact prediction. *Journal of Molecular Modeling*, **20**(11).
- Wozniak, P. P. and Kotulska, M. (2015). AmyLoad: website dedicated to amyloidogenic protein fragments. *Bioinformatics (Oxford, England)*.
- Wright, M. N. and Ziegler, A. (2015). ranger: A Fast Implementation of Random Forests for High Dimensional Data in C++ and R. *arXiv:1508.04409 [stat]*. arXiv: 1508.04409.



Zhou, H. and Zhou, Y. (2004). Quantifying the effect of burial of amino acid residues on protein stability. *Proteins*, **54**(2), 315–322.

Fig. 4. The encoding distance and AUC of reduced alphabets studied in the cross-validation. Red circle: classifier employing best encoding of amino acid. Green square: classifier using full amino acid alphabet. Blue square: classifiers employing encodings from literature. Classifiers having the smallest encoding distance to the best classifier have the highest AUC.