


signalHsmm: prediction of malarial signal peptides

Michał Burdukiewicz^{1*}, Piotr Sobczyk², Paweł Błazej¹, Paweł Mackiewicz¹

¹University of Wrocław, Department of Genomics,

²Wrocław University of Science and Technology, Faculty of Pure and Applied Mathematics,

Amyloids



- n-region: mostly basic residues (Nielsen and Krogh, 1998),
- h-region: strongly hydrophobic residues (Nielsen and Krogh, 1998),
- c-region: a few polar, uncharged residues (Jain et al., 1994).

Acknowledgements and funding

This research was partially funded by the KNOW Consortium and National Science Center (2015/17/N/NZ2/01845).

- Paweł Mackiewicz.
- **biogram** package
(<https://cran.r-project.org/package=biogram>):
 - Piotr Sobczyk,
 - Chris Lauber.

References

- Jain, R. G., Rusch, S. L., and Kendall, D. A. (1994). Signal peptide cleavage regions. functional limits on length and topological implications. *The Journal of Biological Chemistry*, 269(23):16305–16310.
- Nielsen, H. and Krogh, A. (1998). Prediction of signal peptides and signal anchors by a hidden markov model. *Proceedings / ... International Conference on Intelligent Systems for Molecular Biology ; ISMB. International Conference on Intelligent Systems for Molecular Biology*, 6:122–130.